
CONTROL OF ABSCISSION OF FLOWERS AND FRUITS OF PHASEOLUS VULGARIS L.

Dissertation for the Degree of Ph. D. MICHIGAN STATE UNIVERSITY SURANANT SUBHADRABANDHU 1976







This is to certify that the

thesis entitled

CONTROL OF ABSCISSION OF FLOWERS AND FRUITS OF PHASEOLUS VULGARIS L.

presented by

Suranant Subhadrabandhu

has been accepted towards fulfillment of the requirements for

Ph.D. degree in <u>Department</u> of Crop and Soil Sciences

Major professor

Date 7105.3, 1976

**O**-7639



LY Y



RETURNING MATERIALS: Place in book drop to remove this checkout from your record. FINES will be charged if book is returned after the date stamped below.













two dry l

"Seafare house an

differen tern. "

Produced abscissi

ferences signific

synthes

ment ap

decreas

respons

#### ABSTRACT

# CONTROL OF ABSCISSION OF FLOWERS AND FRUITS OF PHASEOLUS VULGARIS L.

By

## Suranant Subhadrabandhu

Abscission of flowers and fruits was studied in two dry bean (Phaseolus vulgaris L.) cultivars, "Black Turtle Soup," an indeterminate, semi-vine type, and "Seafarer," a determinate bush type. Under normal greenhouse and field conditions, these two cultivars exhibited differences in percent pod abscission and flowering pattern. "Black Turtle Soup" had a longer flowering period, produced fewer flowers and exhibited less flower and pods abscission than "Seafarer." In spite of the above differences, seed yields from the two cultivars were not significantly different.

Environmental conditions favorable for photosynthesis, such as long days and carbon dioxide enrichment applied during the reproductive phase in the greenhouse increased vegetative growth and leaf area, and decreased pod abscission. "Black Turtle Soup" was more responsive than "Seafarer."

"Bla starch in ve

ticularly in and starch i

potential we

Bean

decreasing c probabilitie between pod

groups withi abscisic aci

were determi pedicel, usi

treatments a

dihydrophase and dihydrop

"Seafarer" t

With high ab contain high

data obtaine

acid and its sion of dry

Long days and carbon dioxide enrichment promoted vegetative growth and seed yield, increased free sugar and starch in the vegetative parts, and reduced pod abscission.

"Black Turtle Soup" contained more sugar and starch in vegetative tissues than did "Seafarer," particularly in stem and leaf. The levels of free sugar and starch in seeds were similar in the two cultivars.

Bean populations having a low or high abscission potential were established by either increasing or decreasing competition between flowers. Abscission probabilities based on a predetermined relationship between pod size and abscission were assigned to subgroups within each population. Levels of endogenous abscisic acid, phaseic acid and dihydrophaseic acid were determined in methanol extracts of both pod and pedicel, using gas liquid chromatography. None of the treatments affected abscisic acid, phaseic acid or dihydrophaseic acid level in pedicels. Abscisic acid and dihydrophaseic acid levels in pods were higher in "Seafarer" than in "Black Turtle Soup." Populations with high abscission potential from both cultivars also contain higher abscisic acid levels in the pod. The data obtained here cannot explain the role of abscisic acid and its related compounds in flower and pod abscission of dry bean.

in

# CONTROL OF ABSCISSION OF FLOWERS AND FRUITS OF PHASEOLUS VULGARIS L.

Ву


Suranant Subhadrabandhu

## A DISSERTATION

Submitted to
Michigan State University
in partial fulfillment of the requirements
for the degree of

DOCTOR OF PHILOSOPHY

Department of Crop and Soil Sciences



# DEDICATION

To my parents, wife and sons.

I wis gratitude to and Dr. F. G.

valuable adv

course of the

experience o Dr. D. Penne A gr

Hanover who thank Dr. D.

study. Than

and encourage

I w

encourageme

study. Sin

her sacrifi

#### ACKNOWLEDGMENTS

I wish foremost to express my sincere thanks and gratitude to my advisers, Dr. M. W. Adams, Dr. D. Penner, and Dr. F. G. Dennis Jr., for their encouragement, valuable advice, guidance and interest throughout the course of this investigation and for their constructive criticism of this manuscript during its preparation.

I am most honored and grateful to have had the experience of working in both Dr. F. G. Dennis' and Dr. D. Penner's laboratories.

A grateful acknowledgment is extended to Dr. J. W. Hanover who served on the guidance committee. I wish to thank Dr. D. A. Reicosky and Dr. E. A. Mielke who also served on the guidance committee in some parts of this study. Thanks are also extended to Mr. J. L. Taylor for field assistance and to Dr. J. V. Wiersma for suggestions and encouragement. Special thanks to my Thai friends who kindly assisted in field planting in 1974.

I would like to thank my father for his kind encouragement and endless support during my graduate study. Sincere thanks to my loving wife, Pannee, for her sacrifices, understanding, and encouragement

this study and of this manus thanks for the Apinya, for a

especially in

Last:

out my stay

especially in time of frustration during the course of this study and also to her valuable help in preparation of this manuscript. To my sons, Dhanant and Dhanat, many thanks for their unsuspecting sacrifices and to my sister, Apinya, for assisting in preparation of this manuscript.

Lastly, I wish to acknowledge the financial support granted me by the Rockefeller Foundation throughout my stay in America.

LIST OF TABL LIST OF FIGU

INTRODUCTION

CHAPTER 1:

Introduct Genetics Absciss Types of Factors Absciss

Envir

Li Ph Te Ca Re So

Inter

Carbohyd Carbohyd

Extra Analy Role of Role of

REFERENCES.

## TABLE OF CONTENTS

																Page
LIST OF	TABI	LES														
LIST OF	FIGU	JRES														
INTRODUC	CTION	١.														1
CHAPTER	1:	LITE	ERAT	URE	RE	EVIE	W									2
Intro Gene	oduct	ion			· De											2
Ab	sciss	sion		. 01	• 110	·	·			· ·	•	·				4
Type	s of	Abso	ciss	ion												6
Fact																
Ab	sciss	sion		•		•	•	•	•			•	•		•	7
E	nviro	onmer	ntal	Fa	cto	ors				•						7
	Lig	ght i	inte	nsi	ty											7
	Pho	npera rbon lativ	erio	od												8
	Ter	npera	atur	re										•		8
	Car	rbon	dic	xic	le				•							9
	Re:	lativ	ze h	umi	.dit	tу										10
	So:	il mo	oist	ure	s s	tres	SS	•	•	•	•	•	•	•	•	10
I	nter	nal H	act	ors	3.											11
Carb	ohydi	rate	Ava	aila	bi:	lity	7.									13
Carb	ohyd:	rate	Det	ern	nina	atio	on	•	•	•						14
E	xtra	ction	n.													1.4
A	naly	sis														16
Role	of (	Carbo	ohvá	irat	es	in	Abs	cis	ssi	on						16
	of i														i	17
DEFEDEN	CEC															

Abstract.

Introduct Materials Green Field

Results

Green Field Discussi

REFERENCES
CHAPTER 3:

Abstract

Introduc Materia: Results

Pod : Leaf Plan Seed

REFERENCES CHAPTER 4:

> Abstrac Introdu

-----

																Page
CH.	APTER 2:	ABSCI	ss:	ION	OF	FL	OWE	RS	AND	FR	UIT	S II	N P	HA-		
		SEOLU														
		ENCE	IN	FL	OWE	RIN	G P	ATT	ERN	AN:	D A	BSC:	ISS	ION		31
																31
	Abstract.		•	•	•	•	•	•	•	•	•	•	•	•	•	31
	Introduct	tion														32
	Materials		Me	tho	ds											33
	Greenl								•	•	•	•	٠	•	•	33
	Field	Expe	cime	ent	•	٠	٠	•	٠	•	•	•	•	•	•	33
	Results .															34
	Results		•	•	•	•	•	•	•		•	•				
	Greenl	house	Ex	per	ime	nt										34
	Field												•	•		36
																39
	Discussion	on .	٠	•	•	٠	•	•	•	٠	•	•	•	•	•	39
שם	FERENCES									_						47
RE	FERENCES		•	•	•	•	•	•	•	•	•	•	•			
СН	APTER 3:	ABSC	ISS	ION	OF	FL	OWE	ERS	AND	FR	UIT	S I	N P	HA-		
		SECUL	IS '	VUL	GAR	IS	L.	II.	GR	OWT	H A	ND	POD	)		
		ABSC:				AF	FEC	CTEL	BY	DI	FFE	REN	т			49
		DAYL	ENG	THS		•	٠	•	•	•	٠	•	•	٠	•	49
	Abstract															49
	ADSTRACT		•	•	•	•	•	•	•	•						
	Introduc	tion														50
	Material	s and	Me	tho	ds										•	51
	Results	and D	isc	uss	ion					•	•	•	•	•	•	53
																53
	Pod A						•		•	٠	•	•	•	•	•	58
	Leaf	Devel	mqc	ent	•	•		•		•	•	•	•	•	•	58
	Plant	Deve:	lop	men	t.		•		•	•	•	•	•	•	•	60
	Seed '	Yield			•	•	•	•	•	•	•	•	•	•	•	60
																64
RE	FERENCES	-	•	•	•	•	•	•	•	•	•	-	•			
СН	APTER 4:	ABSC	TSS	TON	OF	FI	OWI	ERS	AND	FF	LIU	'S I	NE	HA-		
		anor!		TTTT	CAD	TC	Τ.	TTT	* R	ET.A	TIL	NSH	LP	Or		
		OF C	A DR	OHV	DRA	TES	PI	RODU	JCED	BY	V.	RIA	TIC	N		
		IN D	TVA	ENG	тH											66
		TIN D				•										66
	Abstract									٠	•	٠	•	•	•	66
																68
	Introduc	tion					•	•	•	•	٠	•	•	•	•	0.0

Materials

Sample Determ Determ hydr

Results a Free S Starc Relat

REFERENCES

Ava

CHAPTER 5:

Abstract

Introduc Material Results

Pod i Leaf Plan Seed

REFERENCES
CHAPTER 6:

Abstrac

Introdu Materia

Samp

															Page
Materia	ls an	d M	etho	ods											68
Samo	le Pr	ena	rati	i on										_	68
	rmina														69
	rmina										Ca	rbo	_ `		
	drate														70
11.4	arucc	٠,	1110,	•	•	•	•	•	•	•	•	•			
Results	and	Dis	cus	sion	ı.	•	•	•						•	71
Free	Suga	r.													71
	ch .			•	•	•									79
Dolo	tions	hin	o f	Po.	ו א			ion	ŧ.	T.e	ve 1	ຮັດ	f		
Rela	ailab	1111	Com	hobi	14 W	2+0	-						٠.		86
AV	arrab	те	Cari	DOIL	yur.	ace	٥.	•	•	•	•	•	•	•	00
REFERENCES															89
CHAPTER 5:	SEO	LUS	SION VUI SION	LGA	RIS	L.	IV	. G	ROW	TH .	AND	IN PC	PHA	-	
			E L										•	•	91
Abstrac	t.														91
Introdu	ction														92
Materia			io+h	ods		- 1									92
Results											•	•	•		93
Pod	Absci	ssi	on												93
T.o.a.f	Deve	lor	men	t.											99
Dlan	+ Det	2010	nme	n t											101
Cood	Deve t Dev Yiel	a	pine.												104
seed	1161	.u.	•	•	•										
REFERENCES										•	•	•	•	•	110
CHAPTER 6:	POD	) AE	SIO SCI SBOH	LGA SSI YDR	RIS ON ATE	TO S P	V. DIF ROD	RE FER UCE	LAT ENT D B	IAL Y V	SHI LE ARI	EVEI	LS	<u>:</u> -	111
	IN	CAF	BON	DI	OXI	DE	CON	CEN	TRA	TIC	N.	٠	•	•	111
Abstrac	t.										•	•	•	•	111
															113
Introdu	ction	٠.	. :.		•	•	•	•	•						113
Materia	ıs an	nd N	ieth	uas	•	•	•	•	•	•	•	-			
G	7 - D-		+	ion				_							113

Results a

Free Starc Relat

REFERENCES CHAPTER 7:

> Abstract Introduc Material

Plant Extra Fract Analy

Results

Pod ( Hormo

AI Pi Di

Discuss: REFERENCES

CHAPTER 8: APPENDICES

Appendix

A. GRO

																	Page
Re	sults	an	d D	isc	uss	ion	•										114
	Free	Su	gar														114
	Star			٠.	. ;		:	:	٠.	٠.	٠.	٠.					117
	Rela		n o hyd			ADS	cıs	sio.	n t	O A	vaı	lab	Ie				125
			1				•	•		•	•	•	•		•	•	
REFER	ENCES	٠	٠	٠	•	•	٠	٠	٠	٠	•	•	٠	•	٠	•	130
СНАРТ	ER 7:	S: B: A:	EOL ETW BSC	US EEN ISI	PO PO	GAR D A PHA	IS BSC SEI	OWE L. ISS C,	VI. ION AND	TH: AN: DI:	E R D E HYD	ELA NDO ROP	TIO GEN HAS	NSH OUS EIC	ΙP		101
		A	CID	5 1	N T	HE.	PED	ICE	LS .	AND	F.K	ULT.	S.	٠	٠	•	131
Ab	strac	t.	•	٠	•	•	•		•	•	•	•	•	•	•	•	131
In	trodu	cti	on														133
Ma	teria.	ls .	and	Me	tho	ds	•		•	•	•	•	•	•	٠	•	134
	Plan	t C	ult	ure													134
	Extra	act	ion														135
	Frac																136
	Anal														•		136
Re	sults																137
	Pod (	-ro	w+h	WS	. A	hsc	iss	ion									137
	Hormo									on							139
	Al	зА															139
	P.	A															139
		PA	:	:	÷										•	•	139
Di	scuss	ion															139
REFER	ENCES															•	143
CHAPT	ER 8:	S	UMM	ARY	AN	D C	ONC	LUS	ION	s.		•	•	•		•	145
APPEN	DICES																
Append	dix																
A.	GRO	VTH	AN	D P	OD A	ABS	CIS	sio		s A	FFE	CTE	D B	Y D	IF-		147

Appendix

B. RELAT LEV TIO

C. GROWI

D. RELAT

SELECTED BIE

Append	lix	P	age
В.	RELATIONSHIP OF POD ABSCISSION TO DIFFERENTIAL LEVELS OF CARBOHYDRATES PRODUCED BY VARIA-		
	TION IN DAYLENGTH	•	153
c.	GROWTH AND POD ABSCISSION AS AFFECTED BY CARBON DIOXIDE LEVELS		159
D.	RELATIONSHIP OF POD ABSCISSION TO DIFFERENTIAL LEVELS OF CARBOHYDRATES PRODUCED BY VARIATION IN CARBON DIOXIDE CONCENTRATION		. 165
SELECTE	D BIBLIOGRAPHY		. 171

# CHAPTER 2

1. Effective the

2. Mean Flore Ver Ab

CHAPTER 3

l. Effe Nu Pe Be di

2. Corr of Pa je Af

3. Effe Cu An Le Co

4. Effe Co So he 1

5. Eff P W

# LIST OF TABLES

CHAPTER 2	Page
1. Effects of Cultivar on Yield Components in the Greenhouse	35
<ol> <li>Mean Values per Space-planted Plants of Flowers Produced, Pods Retained at Har- vest and Percentage of Flower and Pod Abscission</li> </ol>	40
CHAPTER 3	
<ol> <li>Effects of Daylengths and Harvesting Time on Number of Flowers, Number of Pods, and Percentage of Pod Abscission of Two Dry Bean Cultivars Grown Under Greenhouse Con- ditions in 1975 and 1976</li> </ol>	5.4
<ol> <li>Correlation Coefficients Between Percentage of Pod Abscission or Leaf Area and Other Parameters in Two Dry Bean Cultivars Sub- jected to Different Daylength Treatments After Flowering.</li> </ol>	57
<ol> <li>Effects of Daylengths, Harvesting Times and Cultivars on Leaf Weight Per Plant, Leaf Area Per Plant and Number of Trifoliate Leaves Per Plant Grown Under Greenhouse Conditions in the Year 1975 and 1976</li> </ol>	59
<ol> <li>Effects of Daylengths, Harvesting Times and Cultivars on the Weight Per Plant of Root, Stem and Total Plant Grown Under Green- house Conditions in the Year 1975 and 1976</li></ol>	61
5. Effects of Daylength on Number of Seeds Per Plant, Total Seed Weight Per Plant and Weight Per Seed in Two Dry Bean Cultivars	62

# CHAPTER 4

l. Devel Eff Pla

2. Effective Fre

3. Effe

4. Effe on Ti

5. Deve

6. Effe

7. Effe St

8. Effe

9. Cor o d D G

CHAPTER 5

l. Sta

2. Eff

CHAPTER 4	
<ol> <li>Developmental Stages, Cultivars and Daylength Effects on Soluble Sugar Level in Various</li> </ol>	-
Plant Parts	72
<ol><li>Effects of Cultivar and Time of Sampling on Free Sugar Concentrations in Bean Tissues .</li></ol>	74
<ol> <li>Effects of Photoperiod and Cultivar on Free Sugar Concentration in Bean Tissues</li> </ol>	75
4. Effects of Photoperiod and Time of Sampling on Free Sugar Concentration of Bean	
Tissues	77
<ol><li>Developmental Stages, Cultivars and Daylength Effects on Starch Level in Various Plant</li></ol>	
Parts	. 80
6. Effects of Photoperiod and Cultivar on Starch Concentration in Bean Tissues	. 81
<ol> <li>Effects of Cultivar and Time of Sampling on Starch Concentration in Bean Tissues.</li> </ol>	. 83
<ol> <li>Effects of Photoperiod and Time of Sampling on Starch Concentration of Bean Tissues.</li> </ol>	. 85
9. Correlation Coefficients Between Percentage of Pod Abscission and Available Carbohydrate in the Vegetative Tissues of the Two Drybean (Phaseolus vulgaris L.) Cultivars Grown Under 3 Daylengths.	88
CHAPTER 5	
<ol> <li>Stages of Pod Development and Carbon Dioxide Levels Effect on Number of Flowers, Number of Pods and Percentage of Abscission in Two Dry Bean Cultivars Grown Under the Two Seasons</li></ol>	94
<ol> <li>Effects of Carbon Dioxide Levels on Number of Flowers, Number of Pods Retained, and Per- centage of Pod Abscission in Two Dry Bean Cultivars Grown Under the Two Seasons</li> </ol>	95

Page

3. Effect Weight Pla Gro

4. Effec Are and Cul

5. Stage ide Lea Cul

6. Corre

7. Effect of Pl. Be.

8. Corr of Ch je Af

CHAPTER 6

1. The

2. Effe Sa of

3. Effe D: in B

4. Eff

	Page
<ol> <li>Effects of Carbon Dioxide Levels on Dry Weight Per Plant of Root, Stem and Total Plant Weight in Two Dry Bean Cultivars Grown Under the Two Seasons</li> </ol>	96
<ol> <li>Effects of Carbon Dioxide Levels on Leaf Area Per Plant, Number of Leaf Per Plant and Leaf Weight Per Plant in Two Dry Bean Cultivars Grown Under the Two Seasons.</li> </ol>	97
<ol> <li>Stages of Pod Development and Carbon Diox- ide Levels Effect on Number of Leaf and Leaf Area Per Plant in Two Dry Bean Cultivars Grown Under the Two Seasons.</li> </ol>	100
6. Correlation Coefficients Between Percentage of Pod Abscission and Other Growth Char- acters in Two Dry Bean Cultivars Sub- jected to Different Carbon Dioxide Levels After Flowering	102
<ol> <li>Effects of Carbon Dioxide Levels on Number of Seeds Per Plant, Seed Weight Per Plant and Weight of One Seed in Two Dry Bean Cultivars Grown Under Two Seasons .</li> </ol>	105
8. Correlation Coefficients Between Percentage of Pod Abscission and Other Reproductive Characters in Two Dry Bean Cultivars Subjected to Different Carbon Dioxide Levels After Flowering	106
CHAPTER 6	
<ol> <li>The Effects of Developmental Stage and Carbon Dioxide Levels on Free Sugar Concentration in Various Plant Parts of the Two Bean Cultivars</li></ol>	115
2. Effects of Carbon Dioxide and Time of Sampling on Free Sugar Concentration of Bean Tissues	118
3. Effects of Developmental Stages and Carbon Dioxide Levels on Starch Concentration in Various Plant Parts of the Two Dry Bean Cultivars	119
• Effects of Carbon Dioxide and Cultivar on	121

4.

5. Effective Sar

6. Corre of dr Tw of

7. Corr Pe Fr Be Di

8. Corr Pe Fr Dr Ca

CHAPTER 7 l. Lev

APPENDIX A A-1. Eff N A

A-2. Eff

A-3. Eff

		Pag
5.	Effects of Carbon Dioxide and Time of Sampling on Starch Concentration of Bean Tissues	124
6.	Correlation Coefficients Between Percentage of Pod Abscission and Available Carbohy- drate Fractions in Vegetative Tissues in Two Dry Bean Cultivars Under Two Levels of Carbon Dioxide	126
7.	Correlation Coefficients Between Seed Weight Per Plant and Available Carbohydrate Fractions in Vegetative Tissues in Two Dry Bean Cultivars Under Two Levels of Carbon Dioxide	127
8.	Correlation Coefficients Between Stem Weight Per Plant and Available Carbohydrate Fractions in Vegetative Tissues in Two Dry Bean Cultivars Under Two Levels of Carbon Dioxide	128
CHAPTE	ER 7	
1.	Levels of ABA, PA and DPA in Relation to Per- centage of Abscission of the Two Bean Cultivars	140
APPEND	DIX A	
A-1.	Effects of Daylengths on Number of Flowers, Number of Pods and Percentage of Pod Abscission at 3 Stages of Development of Two Dry Bean Cultivars Grown Under Greenhouse Conditions in 1975	147
A-2.	Effects of Daylengths on Number of Flowers, Number of Pods and Percentage of Pod Abscission at 3 Stages of Development of Two Dry Bean Cultivars Grown Under Greenhouse Conditions in 1976	148
A-3.	Effects of Daylengths on Leaf Area Per Plant, Number of Leaf Per Plant and Leaf Weight Per Plant at 3 Stages of Development of Two Dry Bean Cultivars Grown Under Green- house Conditions in 1975.	149

A-4. Effective Number 1997 Per 1997 Per

A-5. Effe Pe Pl Tw ho

A-6. Effe Pl Pl of Gr

APPENDIX B

B-1. Effe

B-2. Eff

B-3. Eff O

B-4. Eff o T

B-5. Eff

B-6. Dev

		Page
A-4.	Effects of Daylengths on Leaf Area Per Plant, Number of Leaf Per Plant and Leaf Weight Per Plant at 3 Stages of Development of Two Dry Bean Cultivars Grown Under Green- house Conditions in 1976	150
A-5.	Effects of Daylengths Effects on Root Weight Per Plant, Stem Weight Per Plant and Total Plant Weight at 3 Stages of Development of Two Dry Bean Cultivars Grown Under Green- house Conditions in 1975	151
A-6.	Effects of Daylengths on Root Weight Per Plant, Stem Weight Per Plant and Total Plant Weight at 3 Stages of Development of Two Dry Bean Cultivars Grown Under Greenhouse Conditions in 1976	152
APPEND	DIX B	
В-1.	Effects of Developmental Stages and Daylength on Free Sugar Levels in Vegetative Tissues of Two Phaseolus vulgaris L. Cultivars	153
В-2.	Effects of Developmental Stages and Daylength on Free Sugar Levels in Reproductive Tissues of Two <u>Phaseolus</u> <u>vulgaris</u> L	154
B-3.	Effects of Developmental Stages and Daylength on Starch Levels in Vegetative Tissues of Two $\underline{Phaseolus}$ $\underline{vulgaris}$ L. Cultivars	155
B-4.	Effects of Developmental Stages and Daylength on Starch Levels in Reproductive Tissues of Two Phaseolus vulgaris L. Cultivars	156
В-5.	Effects of Developmental Stages and Daylength on Total Nonstructural Carbohydrate Levels in Vegetative Tissues of Two <u>Phaseolus</u> vul- garis L. Cultivars	157
В-6.	Developmental Stages and Daylength Effects on Total Nonstructural Carbohydrate Levels in Reproductive Tissues of Two Phaseolus vul-	150

APPENDIX C C-1. Effect Flo Pool of

C-2. Effective Floring Poor of

C-3. Effe We an De Gr

C-4. Effe We an De Gr C-5. Effe Pe Le De G:

C-6. Effe Po L D G APPENDIX D

D-1. Eff

D-2. Eff

166

D-3. Effection Care tra

D-4. Effe Ca tr Dr

D-5. Effe Ca st in Be

		-	Page
D-3.	Effects of Stages of Pod Development and Carbon Dioxide Levels on Starch Concen- tration in Root, Stem and Leaf of the Two Dry Bean Cultivars	•	167
D-4.	Effects of Stages of Pod Development and Carbon Dioxide Levels on Starch Concen- tration in Seed and Pod Wall of the Two Dry Bean Cultivars	. j	168
D-5.	Effects of Stages of Pod Development and Carbon Dioxide Levels on Total Non- structural Carbohydrate Concentration in Root, Stem and Leaf of the Two Dry Bean Cultivars.		169
D-6.	Effects of Stages of Pod Development and Carbon Dioxide Levels on Total Non- structural Carbohydrate Concentration in Seed and Pod Wall of the Two Dry Bean		7.0
	Cultivars	1	L70

CHAPTER 2

1. Flow pe un

2. Pods ci ha

3. Pods fl 4. Flow po

5. Pod

6. Poo

CHAPTER 3

CHAPTER

2.

## LIST OF FIGURES

CHAPTE	R 2	Page
1.	Flower production in a specific day as a percentage of total flowers produced under greenhouse conditions	37
2.	Pods retained from flowers borne on a spe- cific day, as a percentage of total pods at harvest grown under greenhouse conditions .	38
3.	Pods retained per day as a percentage of flowers produced on that day	4]
4.	Flower production in a specific day as a percentage of total flowers produced under field conditions	42
5.	Pod retained from flowers borne on a specific day, as a percentage of total pods at harvest grown under field conditions	43
6.	Pods retained per day as a percentage of flowers produced on that day	46
CHAPTE	R 3	
1.	<pre>Interaction between developmental stages and   daylengths on percentage of pod abscission   (A), and total plant weight (B)</pre>	55
CHAPTE	R 6	
1.	Levels of starch in A. root; and B. stem; of two Phaseolus vulgaris L. cultivars during their reproductive phase, averaged from the two carbon dioxide levels	116
2.	Levels of starch in seed of two Phaseolus vulgaris L. cultivars during their reproductive phase, averaged from the two carbon dioxide levels.	123

CHAPTER 7

1. Grow

CHAPTER	7	Pag

1. Growth curves of abscising and persisting pods A. Black Turtle Soup; B. Seafarer . . 138

Abs:

excessive n

limiting yi pods the fa understood.

by the leve

The hypotheses is control:

is control

### INTRODUCTION

Abscission of flowers and immature fruits is characteristic of many economic plants, which produce excessive numbers of flowers, including dry beans (<a href="Phaseolus vulgaris">Phaseolus vulgaris</a> L.), and may be an important factor limiting yield. To prevent abscission of flowers and pods the factors controlling their abscission must be understood. Abscission may be regulated by hormones or by the level of nutrients available, chiefly carbohydrates and/or proteins.

The objectives of this study were to test two hypotheses concerning abscission, namely (a) abscission is controlled by carbohydrate level vs. (b) abscission is controlled by abscisic acid level.

plant. In identified

stalk which cells. An

this zone,

the organs.

traumatic o

and it may some chemi

usually as abscission

searched f

in this ph

synthesize

The induct

#### CHAPTER 1

#### LITERATURE REVIEW

## Introduction

Abscission is the process by which an organ, or part of an organ, separates from the main body of the plant. In fruits and leaves an abscission zone can be identified at the base of the organ or its supporting stalk which is characterized by smaller, less lignified cells. An abscission layer usually develops through this zone, which eventually brings about separation of the organs. In nature abscission occurs at defined stages in the life of a deciduous organ or following traumatic or accidental interruptions of its development, and it may be influenced by environmental changes. Also, some chemical message, sent to the abscission zone, is usually assumed to be responsible for the regulation of abscission layer formation. Physiologists have long searched for evidence of involvement of plant hormones in this phenomenon. These messengers are presumed to be synthesized outside of the abscission zone itself (82). The induction of these hormonal changes is believed to

be controlle internal one plant parts The that of lea: nechanisms first based (55), the s mones are p blade, and sion, analo Ana growth regu for the con This should of the fru angia whic Also, they correlativ fruit abso For exampl With persi In some ir occurs on! develop a

De

be controlled either by environmental factors or by internal ones, such as competition between different plant parts (82).

The physiology of fruit abscission parallels that of leaf abscission, suggesting similar regulatory mechanisms (55). Two arguments support such a hypothesis, first based on the presumed homology of the two organs (55), the second on analogies in their behavior. Hormones are produced in both the fruit and in the leaf blade, and injury to a fruit may result in rapid abscission, analogous to that observed on deblading.

Analyses of hormone levels and responses to growth regulators suggest that a simple, direct mechanism for the control of fruit abscission is unlikely (85). This should not be surprising in view of the complexity of the fruit. Unlike leaves, fruits enclose megasporangia which have a complex development of their own.

Also, they are heterotrophic and therefore more prone to correlative control.

Despite the proposed analogy between leaf and fruit abscission (55) several differences are noteworthy. For example, immaturity and active growth are associated with persistence in leaves, but with abscission in fruit. In some instances (e.g., avocado, tomato), abscission occurs only in young fruit; the mature fruit does not develop a complete abscission layer.

Evi grouped int

sion (23, 1 are often d such as ear

This is evi

Ev

time of flo

vars, comp

late flowe generation in snap be

flowering the case.

(early flo flowering)

(24). In

Since many reviews have dealt with leaf abscission (5, 6, 18, 55, 56, 90), this review will be limited to abscission of reproductive structures.

# Genetics Control of Reproductive Structure Abscission

Evidence for genes control of abscission can be grouped into two parts flowering time and rate of abscission (23, 107). The blossoms of early flowering lines are often damaged by adverse environmental conditions such as early spring frost, and abscise prematurely. The same thing happens in the very late flowering varieties. This is evident in some fruit trees, and certain introduced varieties of field crops.

Evidence for the existence of genes controlling time of flowering was obtained for Phaseolus vulgaris L. (23). In a cross between early and late flowering cultivars, complete dominance for early flowering was observed in  $\mathbf{F}_1$ , and a good fit to 3:1 and 1:1 ratios of early to late flowering plants was observed in  $\mathbf{F}_2$  and in  $\mathbf{F}_2$  x  $\mathbf{F}_1$  generations respectively. Similar results were reported in snap beans, and the dominant gene controlled the early flowering character (28). However, this is not always the case. In the cross of White Seeded Tendergreen (early flowering) x Bush Blue Lake OSC 949-1864 (late flowering), the late flowering character was dominant (24). In the cross of G.N. 1140 (early flowering) x

Nebraska #1 flowering i being domin of more tha flowering. tropical or (79) found backcross : time was qu most of the time exhib Thus genes among vari trolling e habitats ( daylength, of floweri

Gee reproduction reproduction and reproduction in the period apple variation same orches as a same orches determined determined percentage garden be

Nebraska #1 selection (late flowering), the time of flowering is controlled by two major genes, with earliness being dominant to lateness. This indicates involvement of more than one locus for the gene expressing time of flowering. In Goiano and Mexico 450, bean cultivars of tropical origin which are daylength insensitive, Ortega (79) found continuous variation of flowering time in  $F_2$ , backcross 1 and backcross 2, suggesting that flowering time was quantitatively inherited. He concluded that most of the genes involved in the control of flowering time exhibited additive effects and lack of dominance. Thus genes controlling time of flowering appear to differ among varieties. Daylength plays a major role in controlling earliness and in adaptation of cultivars to new habitats (36). Therefore, environment, and especially daylength, may interact with genes in the control of time of flowering.

Genes may control the rate of abscission of reproductive structures, as well as time of flowering. Early work of Heinicke (50) showed marked differences in the percentage of immature fruit abscission in three apple varieties. Since the trees were growing in the same orchard, the rate of fruit drop must have been determined genetically. Cowpea varieties differ in percentage of aborted ovules (75). Soybean (107) and garden bean (13) varieties exhibit different rates of

correlation

vs. yield p

blossom and

Unductive st

ing events

(1) La

is (2) Fa

> ma b:

(3) I<sub>1</sub>

C

ā

S

I

1

blossom and pod drop. In the latter species negative correlations between total number of blossoms per plant vs. yield per plant, and percentage blossom and pod drop vs. yield per plant were reported (13).

## Types of Abscission

Under natural conditions, the abscission of reproductive structures ensues when one or more of the following events occurs.

- Lack of pollination due to unfavorable environmental conditions or absence of insects, which is followed by abscission of the flower;
- (2) Failure of the pollen tube to reach the ovule usually results in the flower abscission. This may be due to genetically controlled incompatibility between the pollen tube and the style.
- (3) In some species, fertilization occurs and the ovule is formed; however, during fruit development the ovule ceases growth and abscission of the young fruit follows. In soybeans, pod abscission was found to occur during the early stages of embryo development, 3 to 7 days after anthesis (59). Competition for carbohydrates, mineral nutrients, and growth regulators may be responsible for this type of abscission.

(4) Fru mat

oco

Environmen The

responsive temperatur

and photop
The carboh

Li

utilized d

overall gr is less li

photosynth abscission increased

are reduce

deciduous

greater or

leaves (1

(4) Fruit maturation. Abscission of fruit at maturity is common in some fruits, however, this type of abscission does not commonly occur in field crops.

# Factors Controlling Reproductive Structures Abscission

### Environmental Factors

The abscission of reproductive structures is very responsive to environmental factors, including light, temperature,  ${\rm CO}_2$ , relative humidity, and moisture stress.

Light intensity. Both photosynthetic effects and photoperiodic effects of light influence abscission. The carbohydrates resulting from photosynthesis are utilized directly for deposition of cell walls. When environmental factors favor photosynthesis but restrict overall growth, cell walls are thicker and abscission is less likely to occur (9). The opposite is true when photosynthesis is limited. Cell walls are thinner and abscission rates are higher. Cotton boll drop is increased from 46 to 86% by shading (30). Pea yields are reduced 50% under 50% shade (71). Shading of soybean plants increases the abscission of pods (69). In deciduous fruit trees abscission of young fruit is greater on twigs deficient in carbohydrate or with few leaves (19).

or low (78)

and in subs

high level: (9, 127, 12

Va

with decres initiation regard to

leads to a buds (9).

strong sin buds alter

zones of t vernal lea

high temp soybean (

Photoperiod. Ojehohon et al. (77) reported that long days induce abscission of flower buds on the main axis of three South American cultivars of Phaseolus vulgaris L., and in subsequent experiments with one of these cultivars they showed that the buds began to fail to develop in long days at about the stage when meiosis occurred. This happened regardless of whether light intensity were high or low (78). Long photoperiods also increase the percentage of flower and pod abscission in soybeans (60, 106). Information on the physiology of the response is only fragmentary (52).

Various reports indicate that long days favor high levels of auxins, gibberellins and abscisic acid (9, 127, 128). Hormonal changes in leaves associated with decreasing day length could be one factor in the initiation of leaf abscission in the fall (9). With regard to vernal leaf abscission, increasing daylength leads to an upsurge of hormone synthesis, especially in buds (9). Consequently, buds and developing shoots became strong sinks, and the auxin moving downward from the buds altered the auxin gradient across the abscission zones of the old leaves. This would serve to promote vernal leaf abscission.

Temperature. Under natural growing conditions, high temperature increases flower and pod abscission in soybean (106), temperatures above 40°C being especially

is due to 1 photosynthe ature (40) abscission temperature In young co period of buds, flow with perio processes Such adver high tempe abscission duction ha of greenho increased enriched a concentrat storage ca tillers in whereas i increased grain fil dioxide e

Ca

effective (

effective (69). Cessation of pod growth at 40°C possibly is due to lowered rates of photosynthesis, since net photosynthesis of soybeans is very low at this temperature (40). In lima bean, the daily rate of blossom abscission was positively correlated with the maximum temperature and possibly the attendant low humidity (22). In young cotton fruits, abscission increased during the period of warm nights (30°C) (43). Abscission of flower buds, flowers and young fruits is frequently correlated with periods of hot weather (118). Various metabolic processes are adversely affected by excessive heat (104). Such adverse metabolic effects probably mediate the high temperature induction of flower and young fruit abscission.

Carbon dioxide. Marked increases in plant production have been obtained by carbon dioxide enrichment of greenhouse atmosphere (119). Soybean yields have been increased when plants were grown in a carbon dioxide enriched atmosphere (20). Increasing carbon dioxide concentration during the reproductive stages may increase storage capacity and yield. The number of ear-bearing tillers in barley was increased by such treatment (41), whereas in rice both grain number and grain size were increased (126). Carbon dioxide enrichment only during grain filling can also increase yield (41, 126). Carbon dioxide enrichment at flowering increased the number of

pods in soy was reduced increased h concentrati 350 to 1,00 fruits by also incre as well as indicate t ment are b abscission abscission appreciabl of ethylen

Al

Re together w lima bean

alone incr when combi

high relat retention

S flowering

flowers a

pods in soybeans under field conditions, but seed size was reduced, whereas enrichment during pod filling increased both seed size and yield (45). Increasing the concentration of carbon dioxide in the atmosphere from 350 to 1,000 ppm decreased abscission of young cotton fruits by nearly 50% (43). Carbon dioxide enrichment also increased the yield of sugar beet and kale (39), as well as many greenhouse crops (119). These results indicate that plants grown under carbon dioxide enrichment are better able to retain fruits.

Although there are few reports on its effects on abscission in intact plants, carbon dioxide retards abscission in explants; it reduces the rate of abscission appreciably (4), and counteracts the promotive action of ethylene (3).

Relative humidity. Low relative humidity, together with high temperature, promotes abscission in lima bean flowers and fruits (22, 63). High humidity alone increases pod set and retention, but is detrimental when combined with low temperature (38). In peanuts, high relative humidity favors the development and retention of fruits (64).

Soil moisture stress. Moisture stress during the flowering and pod filling period increases abscission of flowers and fruits in garden beans (13), and cotton (37).

Soil moistu Among the o auxin (49) ethylene re 127, 128). rainfall ( abscission Internal F flower abs tally unti flowers th on the bas being char the factor in which flower, w that the from the tended to grow more they wou!

Fa

In

on the ba Thus, the Soil moisture stress alters hormone levels in plants. Among the changes reported are: decreases in diffusible auxin (49) and cytokinin activity (54); increases in ethylene release (68) and abscisic acid levels (122, 127, 128). All these changes would favor abscission.

Factors such as nutrient deficiency (69), excess rainfall (14) and severe defoliation (67) also favor abscission of reproductive structures.

### Internal Factors

In lima beans, position on the raceme influences flower abscission (22). Fruit setting occurred acropetally until a "capacity set" was attained; the remaining flowers then abscised. Therefore, pods occurred largely on the basal portion of the raceme, the terminal part being characteristically barren. Ojehomon (76) examined the factors controlling pod retention in Vigna unquiculata, in which only the pods of the basal racemes, the first to flower, were normally retained. Autoradiographs showed that the upper flowers did receive some labelled assimilate from the subtending leaf. Ovaries which failed to develop tended to be smaller at the time of flower opening, to grow more slowly, and eventually to lose weight. However, they would abscise even if comparable in size to those on the basal racemes, unless the latter were removed. Thus, the basal racemes appeared to promote abscission

in the mon the assimi through th

A: reported

like horm

better de Powell an probably

V

pods form on the cl and anti-

pod, naph benzoic a cluded th

in lupine which de of flowe

clusive.

(flower

Posed th

in the more distal ones, not simply by sequestering all the assimilates from the subtending leaf, but probably through the action of diffusible hormones.

Although both Luckwill (66) and Wright (120) reported an inverse relationship between levels of auxinlike hormones and fruit abscission, the nature of the hormones analyzed remains uncertain. Their nature was better defined in subsequent studies (73, 85, 87), but Powell and Pratt (85) concluded that extracted auxins probably do not directly control abscission.

Van Steveninck showed that in lupine the first pods formed promoted abscission of the remaining flowers on the cluster (108, 109, 110, 111, 112). Both auxin and anti-auxin could substitute for the effect of the pod, naphthaleneacetic acid (NAA) and 2, 3, 5 triiodobenzoic acid (TIBA) being the most effective. He concluded that the controlling factor may be an "antiauxin" (110). A search for abscission accelerating factors in lupine revealed the presence of an auxin-like material which delayed abscission of pods, but promoted abscission of flowers (112).

Studies of auxin applications to fruit are inconclusive. Synthetic auxins may increase abscission (flower and young fruit) (12), or increase persistence (mature fruit and seedless fruit) (33). Hartmann proposed that if auxin increase persistence, antiauxin

auxins to 1 confirming "antiauxin Auxins alo abscission lators in with the i fruit (10, Ca energy sto accumulate when synth reserve ca exceeds s The level numerous 94, 97).

intensity
balance i
exceeds s
carbohydr

should pron

should promote abscission (46). Application of antiauxins to mature olives promoted abscission, apparently confirming this suggestion. However, both auxin and "antiauxin" promote abscission in lupine flowers (110). Auxins alone may be insufficient to account for fruit abscission (55). Evidence associating abscission stimulators in fruit has accumulated (111, 112), culminating with the identification of abscisic acid (ABA) in cotton fruit (10, 74).

#### Carbohydrate Availability

Carbohydrates are the primary source of reserve energy stored in the vegetative organs of plants. They accumulate during periods of high photosynthetic activity when synthesis exceeds utilization. Conversely, these reserve carbohydrates are degraded when utilization exceeds synthesis, and are essential for survival (99). The level of carbohydrate reserves may be influenced by numerous cultural and environmental factors (16, 58, 88, 94, 97).

Under conditions of low leaf area, low light intensity or supra-optimal temperature, the energy balance is likely to be negative, since the demand exceeds supply (16). Under such conditions, the reserve carbohydrates are likely to be utilized. Decreases in reserve carbohydrates are usually observed just after

defoliation
when suffi

Re

synonymous mentioned drates wh

building : erally, i

carbohydr starch (1 only as s

(TAC) had

term "tot replaceme

confusion obtain en drates.

Extracti

been rep

(16, 88

defoliation (88). Surplus carbohydrates can be restored when sufficient leaf area has developed to exceed the energy requirement of the crop.

Reserve carbohydrates mentioned in this sense are synonymous with the term "total available carbohydrates" mentioned in many papers (116), and include all carbohydrates which can be used as a source of energy or as building material, either directly or indirectly. Generally, in higher green plants, the bulk of available carbohydrates consists of sugars, fructosans, dextrin and starch (116), whereas hemicellulose and cellulose serve only as structural materials.

Although the term "total available carbohydrates" (TAC) had been widely used, Smith (96) introduced the term "total nonstructural carbohydrate (TNC)" as a replacement for "total available carbohydrate" to avoid confusion with animal physiology, ruminants being able to obtain energy from at least part of the structural carbohydrates.

## Carbohydrate Determination

### Extraction

Various extraction and analysis techniques have been reported for plant carbohydrates. Almost all the available literature deals with grasses and forage crops (16, 88, 99, 113); however, these methods have been

adapted fo (84, 114). In

by a giver others, th

judgment o to evaluat T

legumes a and subtre from these

the form

(116) or native to

fructosan has been with wate

in water. very smal

cated tha of gluco

80% etha hours wa in barle

extracte

adapted for horticultural (11, 121) and forest crops (84, 114).

In certain cases the form of carbohydrate stored by a given species dictates the techniques (99). In others, the choice may depend on the convenience and judgment of the investigator. This has made it difficult to evaluate critically data from different laboratories.

To develop an extraction technique, one must know the form of carbohydrate stored in the plant. Many legumes accumulate sucrose and starch (97), as do tropical and subtropical grasses (95). TNC has been extracted from these species with takadiastase enzyme preparations (116) or with acid (88). On the other hand, grasses native to temperate latitudes accumulate sucrose and fructosan (101, 102). Therefore, TNC from these species has been extracted with dilute acid solution (88), or with water (113), since fructosan is readily soluble in water. Fructosan-storing grasses may also accumulate very small amounts of water-insoluble carbohydrates (95).

The extraction of reducing sugars is less complicated than that of TNC. Reducing sugars consist mainly of glucose and fructose, which can be extracted by using 80% ethanol. Extraction in boiling 80% ethanol for 4 hours was found to be sufficient to remove all the sugar in barley leaves (124). Later, Yemm and Willis (125) extracted soluble sugars at 37°C with absolute alcohol,

the plant results us

followed by Extraction for two ho

Analysis Fo

niques are

Micro copp

a high de

inability Vegetativ

(1, 103).
complete
as eviden

The envir

Cess is 1

energy in

abscissio

followed by two further extractions with 70% ethanol. Extraction of white clover tissue with hot 80% ethanol for two hours removed all the glucose and fructose from the plant (62). Recent studies also report satisfactory results using 80% ethanol for soluble sugars (11, 121).

#### Analysis

Four commonly used carbohydrate analysis techniques are: (a) anthrone, (b) phenol-sulfuric,
(c) Nelson's arsenomolybdate and (d) Shaeffer-Somogyi
Micro copper (53).

#### Role of Carbohydrates in Abscission

In woody plants, rapid vegetative growth imposes a high demand for carbohydrates and mineral nutrients, and abscission of immature fruits may result from the inability of the fruits to compete successfully with vegetative tissues and with each other for carbohydrate (1, 103). If competition is increased by partial or complete leaf removal, the rate of abscission increases, as evident in cotton (32, 70), and soybean fruits (67). The environment can affect abscission partly by the effect on photosynthetic rate, as discussed previously.

The role of carbohydrate in the abscission process is not clear. Carbohydrate is used as a source of energy in many metabolic processes which may affect abscission. However, there is growing evidence for the

role of pla role of cas

Th

contained summer. A fruits or

studies.

noted absorber pl

abscissio sometimes

D relation

fruits.

reveals t

peach fru

Finally,

or no ef

role of plant hormones in abscission (6, 18), and the role of carbohydrates may be related to that of hormones (6).

# Role of Abscisic Acid (ABA) in Abscission

The recent discovery of ABA (7) aroused the interest of many physiologists conducting abscission studies. Bornman (15) found that Streptocarpus leaves contained five times as much ABA in autumn as in midsummer. Application of cotton extracts to intact young fruits or to defruited pedicels of cotton strongly promoted abscission (7). Application of ABA to leaves or entire plants in the field usually accelerated leaf abscission (98). However, repeated applications were sometimes required to obtain a response (98).

Davis and Addicott (26) reported a strong correlation between ABA content and abscission in cotton fruits. However, their data are for <u>cumulative</u> abscission, rather than <u>rate</u> of abscission. Critical analysis reveals that ABA content was better correlated with fruit growth rate than with abscission. Further, in immature peach fruit the concentration of an ABA-like inhibitor is greater in persisting than in abscising fruit (129). Finally, application of synthetic ABA to fruit has little or no effect on abscission (47, 48, 130).

The understood fruit absc:

were obser levels wer ABA is mos

intact pla

or to repe

is still to solve

process, therefore The physiology of the ABA effect is not well understood, and convincing evidence for its role in fruit abscission is still lacking. High levels of ABA were observed in young nonabscising leaves, whereas the levels were lower in old and abscising leaves (81, 105). ABA is most effective on explants (2, 7, 8, 25, 83, 97); intact plants respond only to very high concentrations or to repeated applications (21, 31, 35, 47).

Although abscission of flowers and immature fruits is becoming a problem in many economic crops, its cause is still unknown. This creates difficulty in an effort to solve the problem of abscission. Due to its complex process, abscission can be influenced by many factors, therefore care should be taken in any study on abscission.

l. Abbott th 48

2. Abeles 3. Abeles Al e

4. Abele T

5. Addic E 1 B

6. Addic 7. Addic

8. Addio

9. Addio

#### REFERENCES

- Abbott, D. L. 1960. The bourse shoot as a factor in the growth of apple fruits. Ann. Appl. Biol. 48: 434-438.
- Abeles, F. B. 1967. Mechanism of action of abscission accelerators. Physiol. Plant. 20: 442-454.
- Abeles, F. B., L. E. Craker and G. R. Leather. 1971. Abscission: The phytogerontological effects of ethylene. Plant Physiol. 47: 7-9.
- Abeles, F. B. and H. E. Gahagan. 1968. Abscission: The role of ethylene, ethylene analogues, carbon dioxide and oxygen. Plant Physiol. 43: 1255-1258.
- Addicott, F. T. 1965. Physiology of abscission. In Encyclopedia of Plant Physiology, XV (Part 2), 1094-1126. Ed. W. Ruhland. Springer-Verlag, Berlin.
- Addicott, F. T. 1970. Plant hormones in the control of abscission. Biol. Rev. 45: 485-524.
- Addicott, F. T., H. R. Carns, J. L. Lyon, O. E. Smith and J. S. McMeans. 1964. On the physiology of abscisins. In Régulateurs naturels de la croissance végétale, 687-703. Ed. J. P. Nitsch, CNRS, Paris.
- Addicott, F. T. and J. T. Lyon. 1969. Physiology of abscisic acid and related substances. Ann. Rev. Plant Physiol. 20: 139-164.
- Addicott, F. T. and J. L. Lyon. 1973. Physiological ecology of abscission. In Shedding of Plant Parts, 85-124. Ed. T. T. Kozlowski, Academic Press.

10. Addic Th fo Wa

ll. Aung, Ch du

12. Batje 13. Bink

14. Black

l5. Born

l6. Brow b

17. Buko

18. Carr

19. Char

20. Coo

21. Coo

- Addicott, F. T., J. L. Lyon, K. Ohkuma, W. E.
  Thiessen, H. R. Carns, O. E. Smith, J. W. Cornforth, B. V. Milborrow, G. Ryback and P. F.
  Wareing. 1968. Abscisic acid: A new name for
  abscisin II (Dormin). Science 159: 1943.
- Aung, L. H., F. Tognoni and A. A. De Hertogh. 1973. Changes in the carbohydrates of tulip bulbs during development. Hort. Sci. 8: 207-208.
- Batjer, L. P. 1965. Fruit thinning with chemicals. Agr. Information Bull. No. 289. U.S. Dept. Agr., Agr. Res. Serv.
- Binkley, A. M. 1932. The amount of blossom and pod drop on six varieties of garden beans. Proc. Amer. Soc. Hort. Sci. 29: 489-492.
- 14. Blackwell, F. L. C. 1969. Effects of weather, irrigation, and pod removal on the setting of pods and the marketable yield of runner beans Phaseolus multiflorus. J. Hort. Sci. 44: 371-384.
- Bornman, C. H. 1969. Laminal abscission in <u>Streptocarpus</u>. Abstracts. XI<sup>th</sup> Int. Bot. Congress. Seattle U.S.A., p. 18.
- 16. Brown, R. H. and R. E. Blaser. 1965. Relationships between reserve carbohydrate accumulation and growth rate in Orchardgrass and Tall Fescue. Crop Sci. 5: 577-582.
- Bukovac, M. J., F. Zucconi, R. P. Larsen and C. D. Kesner. 1969. Chemical promotion of fruit abscission in cherries and plums with special reference to 2-Chloroethanephosphonic acid. J. Amer. Soc. Hort. Sci. 94: 226-230.
- Carns, H. R. 1966. Abscission and its control. Ann. Rev. Plant Physiol. 17: 295-314.
- Chandler, W. H. 1951. Deciduous Orchards. Lea and Febiger Publ., Philadelphia.
- Cooper, R. L. and W. A. Brun. 1967. Response of soybeans to a carbon dioxide enriched atmosphere. Crop Sci. 7: 455-457.
- Cooper, W. C. and W. H. Henry. 1967. The acceleration of abscission and colouring of Citrus fruit. Proc. Fla. St. Hort. Soc. 80: 7-14.

22. Cordi

23. Coyne

24. Coyn o i S

25. Crac 26. Davi

27. Dela

28. Dic

29. Drey 30. Dun

31. Eag

32. Eat

33. Edg

- Cordner, H. B. 1933. External and internal factors affecting blossom drop and set of pods in lima beans. Proc. Amer. Soc. Hort. Sci. 30: 571-576.
- Coyne, D. P. 1970. Genetic control of a photoperiodtemperature response for time of flowering in beans (Phaseolus vulgaris L.). Crop Sci. 10: 246-248.
- Coyne, D. P. and R. H. Mattson. 1964. Inheritance of time of flowering and length of blooming period in <u>Phaseolus vulgaris</u> L. Proc. Amer. Soc. Hort. Sci. 85: 366-373.
- Cracker, L. E. and F. B. Abeles. 1969. Abscission: Role of abscisic acid. Plant Physiol. 44: 1144-1149.
- Davis, L. A. and F. T. Addicott. 1972. Abscisic acid: Correlations with abscission and with development in the cotton fruit. Plant Physiol. 49: 644-668.
- Dela Fuente, R. K. and A. C. Leopold. 1968.
   Senescence processes in leaf abscission. Plant Physiol. 43: 1496-1502.
- Dickson, M. H. 1967. Diallel analysis of seven economic characters in snap beans. Crop Sci. 7: 121-124.
- 29. Dreywood, R. 1946. Industr. Engng. Chem. Anal. ed. 18: 499.
- Dunlap, A. A. 1943. Low light intensity and cotton boll shedding. Science 98: 568-569.
- Eagles, C. F. and P. F. Wareing. 1964. The role of growth substances in the regulation of bud dormancy. Physiol. Plant. 17: 697-709.
- Eaton, F. M. and N. E. Rigler. 1945. Effects of light intensity, nitrogen supply and fruiting on carbohydrate utilization by the cotton plant. Plant Physiol. 20: 380-411.
- Edgerton, L. J., C. G. Forshey and G. D. Blanpied. 1967. Effect of summer applications of Alar (B 995) in reducing drop and improving color and firmness of apples. Proc. New York State Hort. Soc. 112: 204-206.

34. Edger la wi So

35. El-An 19 (d

36. Evans fl TA De

37. Ewing fa fi Te

38. Fishe

39. Ford c k

40. Fuku

41. Giff

42. Grig

43. Guir

- Edgerton, L. T. and W. J. Greenhalgh. 1969. Regulation of growth, flowering and fruit abscission with 2-Chloroethanephosphonic acid. J. Amer. Soc. Hort. Sci. 94: 11-13.
- El-Antably, H. M. M., P. F. Wareing and J. Willman. 1967. Some physiological responses to d,1-abscisin (dormin). Planta. 73: 74-90.
- 36. Evans, L. T. and R. W. King. 1974. Factors affecting flowering and reproduction in grain legume. TAC/FAO. Working Party on Grain Legumes, New Delhi, 1974.
- Ewing, E. C. 1918. A study of certain environmental factors and varietal differences influencing the fruiting of cotton. Mississippi Agr. Exp. Sta. Tech. Bull. 8: 1-95.
- 38. Fisher, V. J. and C. K. Weaver. 1974. Flowering, pod set, and pod retention of lima bean in response to night temperature, humidity and soil moisture. J. Amer. Soc. Hort. Sci. 99: 448-450.
- Ford, M. A. and G. N. Thorne. 1967. Effect of CO<sub>2</sub> concentration on growth of sugar beet, barley, kale and maize. Ann. Bot. N.S. 31: 629-644.
- Fukui, J., O. Mutsuo and I. Watanabe. 1965. Studies on the seed production of soybean. I. Effect of temperature on photosynthesis of soybean. Proc. Crop Sci. Soc. Japan 33: 432-436.
- 41. Gifford, R. M., P. M. Bremner and D. B. Jones. 1973. Assessing photosynthetic limitation to grain yield in a field crop. Aust. J. agric. Res. 24: 297-307.
- 42. Griggs, W. H., B. T. Iwakiri, R. B. Fridley and J. Mehlaschau. 1970. Effect of 2-Chloroethylphosphonic acid and cycloheximide on abscission and ripening of "Bartlett" pears. Hort. Science 5: 264-266.
- Guinn, G. 1974. Abscission of cotton floral buds and bolls as influenced by factors affecting photosynthesis and respiration. Crop Sci. 14: 291-293.

44. Hall, 19 of ab

45. Hardr at fe

46. Hartı 0

47. Hart

48. Hart

49. Hart

50. Hei

51. Heir

52. Hen

53. Hod

- 44. Hall, W. C., F. A. Herrero and F. R. H. Katterman. 1961. Leaf abscission in cotton. IV Effects of a natural promoter and amino acids on abscission in cotyledonary node explants. Botan Gaz. 123: 29-34.
- 45. Hardman, L. L. and W. A. Brun. 1971. Effect of atmospheric carbon dioxide enrichment at different developmental stages on growth and yield components of soybeans. Crop Sci. 11: 886-888.
- Hartmann, H. T. 1955. Induction of abscission of olive fruits by maleic hydrazide. Bot. Gaz. 123: 29-34.
- Hartmann, H. T., A. J. Heslop and J. Whisler. 1968. Chemical induction of fruit abscission in olives. Calif. Agric. 22: 14-16.
- Hartmann, H. T., A. Tombesi and J. Whisler. 1970.
  Promotion of ethylene evolution and fruit abscission in the olive by 2-chloroethanephosphonic acid and cycloheximide. J. Amer. Soc. Hort. Sci. 95: 635-640.
- 49. Hartung, W. and J. Witt. 1968. Uber der Einfluss der Bodenfeuchtigkeit auf den Wuchsstoffgehatt von Anastatica hierochuntica und Helianthus annuus. Flora (Jena). Abt. B 157: 603-614.
- Heinicke, A. J. 1917. Factors influencing the abscission of flowers and partially developed fruits of the apple (Pyrus malus). Cornell Univ. Agr. Exp. Sta. Bull. 393: 44-112.
- Heinze, P. H. and A. E. Murneek. 1940. Comparative accuracy and efficiency in determination of carbohydrates in plant material. Missouri Agri. Exp. Sta. Res. Bull. 314.
- 52. Hendricks, S. B. and H. A. Borthwick. 1965. The physiological functions of phytochrome. In "Chemistry and Biochemistry of Plant Pigments" 405-436. Ed. T. W. Goodwin, Academic Press, New York.
- 53. Hodge, J. E. and B. T. Hofrieter. 1962. Determination of reducing sugars and carbohydrates. In "Methods in Carbohydrate Chemistry," Vol. I, pp. 380-394. Ed. R. L. Whistler and M. L. Wolfrom, Academic Press, New York.

54. Itai, in 87

55. Jacob In Re

56. Jacob ab 57. Jacob 19

58. Jung e: c: a 3

59. Kato A a K

61. Koch

62. Laid

63. Lami

64. Lee

- 54. Itai, C. and Y. Vaadia. 1971. Cytokinin activity in water-stressed shoots. Plant Physiol. 47: 87-90.
- Jacobs, W. P. 1962. Longevity of plant organs: Internal factors controlling abscission. Ann. Rev. Plant Physiol. 13: 403-436.
- 56. Jacobs, W. P. 1968. Hormonal regulation of leaf abscission. Plant Physiol. 43: 1480-1495.
- Jacobs, W. P., J. A. Shield, Jr., D. J. Osborne. 1962. Senescence factor and abscission of Coleus leaves. Plant Physiol. 37: 104-106.
- Jung, G. A. and D. Smith. 1960. Influence of extended storage at constant low temperature on cold resistance and carbohydrate reserves of alfalfa and medium red clover. Plant Physiol. 35: 123-125.
- 59. Kato, I., S. Sakaguchi and Y. Naito. 1955. Anatomical observations on fallen buds, flowers and pods of soy-bean, <u>Glycine Max. M. Tokai-</u> Kinki Nat. Agr. Exp. Sta. Bull. 2: 159-168.
- Kiyosawa, S. and K. Kiyosawa. 1962. Effect of daylength on fruiting in soybean plant. Proc. Crop Sci. Soc. Japan 30: 342-345.
- 61. Koch, R. B., W. F. Geddes and F. Smith. 1951.

  The carbohydrates of Gramineae. I. The sugars of the flour of wheat (<u>Triticum vulgare</u>) Cereal Chem. 28: 424-430.
- 62. Laidlaw, R. A. and S. G. Reid. 1952. Analytical studies on the carbohydrates of grass and clovers. I. Development of methods for the estimation of free sugar and fructosan contents. J. Sci. Fd. Agric. 3: 19-25.
- Lambeth, V. N. 1950. Some factors influencing pod set and yield of the lima bean. Univ. of Missouri Agr. Exp. Sta. Res. Bull. 466.
- 64. Lee, T. A., D. L. Ketring and R. D. Powell. 1972. Flowering and growth response of peanut plants (Arachis hypogara. L.) at two levels of relative humidities. Plant Physiol. 49: 190-193.

65. Lipe, re Ph

66. Luckw in du to

67. McAli

68. McMi. 1' p P. 69. Mann

70. Maso

71. Mead

72. Nels

73. Nits

74. Ohk

<sup>75.</sup> Oje

<sup>76.</sup> Oje

- Lipe, J. A. and P. W. Morgan. 1973. Ethylene, a regulator of young fruit abscission. Plant Physiol. 51: 949-953.
- 66. Luckwill, L. C. 1953. Studies of fruit development in relation to plant hormones. I. Hormone production by the developing apple seed in relation to fruit drop. J. Hort. Sci. 28: 14-24.
- McAlister, D. F. and O. A. Krober. 1958. Response of soybeans to leaf and pod removal. Agron. J. 50: 674-677.
- 68. McMichael, B. L., W. R. Jordan and R. D. Powell. 1972. An effect of water stress on ethylene production by intact cotton petioles. Plant Physiol. 49: 658-660.
- Mann, J. D. and E. G. Jaworski. 1970. Comparison of stresses which may limit soybean yields. Crop Sci. 10: 620-624.
- Mason, T. G. 1922. Growth and abscission in Sea Island cotton. Ann. Bot. 36: 457-483.
- Meadley, J. T. and G. M. Milbourn. 1971. The growth of vining peas. III. The effect of shading on abscission of flowers and pods. J. Agric. Sci. 77: 103-108.
- Nelson, N. 1944. A photometric adaptation of the Somogyi method for the determination of glucose. J. Biol. Chem. 153: 375-379.
- Nitsch, J. P., C. Pratt, C. Nitsch, and N. J. Shaulis. 1960. Natural growth substances in concord and concord seedless grapes in relation to berry development. Amer. J. Bot. 47: 566-576.
- Ohkuma, K., F. T. Addicott, O. E. Smith and W. E. Thiessen. 1965. The structure of abscisin. II. Tetrahedron letters 29: 2529-2535.
- 75. Ojehomon, O. O. 1970. A comparison of the vegetative growth, development and seed yield of three varieties of cowpea, Vigna unguiculata (L.). Walp. J. Agric. Sci. 74: 363-374.
- Ojehomon, O. O. 1972. Fruit abscission in cowpea, Vigna unguicuiata. L. Walp. J. Expt. Bot. 23: 751-761.

77. Ojeho 19 ar Pł

79. Orted

80. Osbo b 81. Osbo B

82. Osbo a E

83. Palm

84. Park

85. Power 86. Rasm

87. Rati

88. Rey

- Ojehomon, O. O., A. S. Rathjen and D. G. Morgan. 1968. Effects of daylength on the morphology and flowering of five determinate varieties of Phaseolus vulgaris. L. J. Agric. Sci. 71: 209-214.
- Ojehomon, O. O., M. S. Zehni and D. G. Morgan. 1973.
   The effects of photoperiod on flower-bud development in Phaseolus vulgaris. Ann. Bot. 37: 871-884.
- Ortega, Y. S. 1971. Inheritance of flowering time in French bean (<u>Phaseolus</u> <u>vulgaris</u>). Agronomic Tropical 21: 319-328.
- Osborne, D. J. 1955. Acceleration of abscission by a factor produced in senescent leaves. Nature 176: 1161-1163.
- Osborne, D. J. 1968. Defoliation and defoliants. Botan. Gaz. 123: 29-34.
- Osborne, D. J. 1973. Internal factors regulating abscission. In Shedding of Plant Parts, 125-148, Ed. T. T. Kozlowski, Academic Press.
- Palmer, R. L., H. Z. Hield and L. N. Lewis. 1969.
   Citrus leaf and fruit abscission. Proc. 1st. Int. Citrus Symp., 3: 1135-1143.
- 84. Parkerson, R. H. and F. W. Whitmore. 1972. A correlation of stem sugars, starch and liquid with wood formation in Eastern White pine. Forest Sci. 18: 178-183.
- Powell, L. E. and C. Pratt. 1966. Growth promoting substances in the developing fruit of peach (Prunus persica L.). J. Hort. Sci. 41: 331-348.
- Rasmussen, H. P. and M. J. Bukovac. 1969. A histochemical study of abscission layer formation in the bean. Amer. J. Bot. 56: 69-76.
- 87. Ratner, A., R. Goren and S. P. Monselise. 1969. Activity of pectin esterase and cellulase in the abscission zone of <u>Citrus</u> leaf explants. Plant Physiol. 44: 1717-1723.
- Reynolds, J. H. and D. Smith. 1962. Trend of carbohydrate reserves in alfalfa, smooth bromegrass and timothy grown under various cutting schedules. Crop Sci. 2: 333-336.

89. Rubi

90. Rubi 0 3 91. Shaf i

92. Shaf

93. Sha:

95. Smi

96. Smi

98. Smi

99. sm:

- Rubinstein, B. and A. C. Leopold. 1962. Effects of amino acids on bean leaf abscission. Plant Physiol. 37: 398-401.
- Rubinstein, B. and A. C. Leopold. 1964. The nature of leaf abscission. Quarterly Rev. Biol. 39: 356-372.
- Shaffer, P. A. and A. F. Hartmann. 1921. The iodimetric determination of copper and its use in sugar analysis I. J. Biol. Chem. 45: 349-364.
- Shaffer, P. A. and A. F. Hartmann. 1921. The iodimetric determination of copper and its use in sugar analysis II. J. Biol. Chem. 45: 365-390.
- Shaffer, P. A. and M. Somogyi. 1933. Copper-Iodometric reagents for sugar determination. J. Biol. Chem. 100: 695-713.
- Smith, D. 1950. Seasonal fluctuations of root reserves in red colver <u>Trifolium pratense</u> L. Plant Physiol. 25: 702-710.
- Smith, D. 1968. Classification of several native North American grasses as starch or fructosan accumulators in relation to taxonomy. Jour. Brit. Grassl. Soc., 23: 306-309.
- 96. Smith, D. 1969. Removing and analyzing total nonstructural carbohydrates from plant tissue. Research Report #41. University of Wisconsin, 12 pp.
- Smith, D. and L. F. Graber. 1948. The influence of top growth removal on the root and vegetative development of biennial sweet clover. J. Am. Soc. Agron. 40: 818-831.
- 98. Smith, O. E., J. L. Lyon, F. T. Addicott and R. E. Johnson. 1968. Abscission physiology of abscisic acid. In Biochemistry and Physiology of Plant Growth Substances, 1547-1560. Ed. F. Wightman and G. Setterfield, Runge Press, Ottawa.
- 99. Smith, D., G. M. Paulsen and C. A. Raguse. 1964. Extraction of total available carbohydrates from grass and legume tissue: Plant Physiol. 39: 960-962.

100. Somodicio

103. Swee

105. Ture

106. Van

107. Van

109. Van

110. Var

- 100. Somogyi, M. 1937. A reagent for the copperiodometric determination of very small amounts of sugar. J. Biol. Chem. 117: 771-776.
- Sprague, V. G. and J. T. Sullivan. 1950. Reserve carbohydrates in orchardgrass clipped periodically. Plant Physiol. 25: 92-102.
- 102. Sprague, V. G. and J. T. Sullivan. 1952. Reserve carbohydrates in orchard grass cut for hay. Plant Physiol. 28: 304-313.
- 103. Sweet, G. B. 1973. Shedding of reproductive structures in forest trees. In "Shedding of Plant Parts," 341-382. Ed. T. T. Kozlowski, Academic Press, New York.
- 104. Treshow, M. 1970. Environment and plant response. McGraw-Hill, Publ., New York.
- 105. Tureskaya, R., V. Kefeli, M. Kutacek, K. Vackova, N. Tschumarovski and T. Krupnikova. 1968. Isolation and some physiological properties of natural plant growth inhibitors. Biol. Plantarum, 10: 205-221.
- 106. Van Schaik, P. H. and A. H. Probst. 1958. Effects of some environmental factors on flower production and reproductive efficiency in soybeans. Agron. Jour. 50: 192-197.
- 107. Van Schaik, P. H. and A. H. Probst. 1958. The inheritance of inflorescence type, peduncle length, flowers per node, and percent flower shedding in soybeans. Agron. J. 50: 98-102.
- 108. Van Steveninck, R. F. M. 1957. Influence of peamosaic virus on the reproductive capacity of vellow lupins. Bot. Gaz. 119: 63-70.
- 109. Van Steveninck, R. F. M. 1957. Factors affecting the abscission of reproductive organs in yellow lupins (<u>Lupinus luteus L.</u>) I. The effect of different patterns of flower removal. J. Expt. Bot. 8: 373-381
- 110. Van Steveninck, R. F. M. 1958. Factors affecting the abscission of reproductive organs in yellow lupins (Lupinus luteus L.) II. The effects of growth substances, defoliation, and removal of lateral growth. J. Expt. Bot. 9: 372-383.

111. Van S ac Ne
112. Van S th
113. Wait
113. Wait
114. Warg
1 115. Webs

ll6. Wein

117. Wit

118. Wit

120. Wr

121. Wr

- 111. Van Steveninck, R. F. M. 1959. Abscission accelerators in lupins (Lupinus luteus L.)
- 112. Van Steveninck, R. F. M. 1959. Factors affecting the abscission of reproductive organs in yellow lupins (<u>Lupinus luteus L.</u>) III. Endogenous growth substances in virus-infected and healthy plants and their effect on abscission. J. Expt. Bot. 10: 367-375.
- 113. Waite, R. and J. Boyd. 1953. The water soluble carbohydrates of grasses. I. Changes occurring during the normal life cycle. J. Sci. Fd. Agric. 4: 197-204.
- 114. Wargo, P. M. 1971. Seasonal changes in carbohydrate levels in roots of sugar maple. U. S. Forest Ser. Res. Pao. N.E. 213: 1-8.
- 115. Webster, B. D. 1973. Anatomical and histochemical changes in leaf abscission. In Shedding of Plant Parts 45-84. Ed. T. T. Kozlowski, Academic Press. New York.
- 116. Weinmann, H. 1947. Determination of total available carbohydrates in plants. Plant Physiol. 22: 279-290.
- 117. Wittenbach, V. A. and M. J. Bukovac. 1974. Cherry fruit abscission. Evidence for time of initiation and involvement of ethylene. Plant Physiol. 54: 494-498.
- 118. Wittwer, S. H. 1954. Control of flowering and fruit setting by plant regulators. In "Plant Regulators in Agriculture," 62-80. Ed. H. B. Tukey, Wiley, Publ., New York.
- 119. Wittwer, S. H. and W. M. Robb. 1964. Carbon dioxide enrichment of greenhouse atmospheres for food crop production. Econ. Bot. 18: 34-56.
- 120. Wright, S. T. C. 1956. Studies of fruit development on relation to plant hormones. III. Auxins in relation to fruit morphogenesis and fruit drop in the black currant <u>Ribes</u> <u>nigrum</u>. J. Hort. Sci. 31: 196-211.
- 121. Wright, R. D. and L. H. Aung. 1975. Carbohydrates in two rhododendron cultivars. J. Amer. Soc. Hort. Sci. 100: 527-529.

122. Wright Street Stre

124. Yemi

125. Yem

127. Zee

128. Zee

129. Zuo

130. Zu

- 122. Wright, S. T. C. and R. W. P. Hiron. 1970. The accumulation of ABA in plants during wilting and under other stress conditions. In "Plant Growth Substances," 291-298. Ed. D. J. Carr, Springer Verlag, Publ., New York.
- 123. Yamaguchi, S. 1954. Some interrelations of oxygen, carbon dioxide, sucrose and ethylene in abscission. Ph.D. Dissertation, University of California, Los Angeles.
- 124. Yemm, E. W. 1935. The respiration of barley plants. I. Methods for the determination of carbohydrates in leaves. Proc. Roy. Soc. B, 117: 483-504.
- 125. Yemm, E. W. and A. J. Willis. 1954. Stomatal movements and changes of carbohydrate in leaves of Chrysanthemum maximum. New Phytol. 53: 373-396.
- 126. Yoshida, S., J. H. Cock and F. T. Parao. 1972. Physiological aspects of high yields. In "Rice Breeding," IRRI, Los Banos, pp. 455-468.
- 127. Zeevaart, J. A. D. 1971. Effects of photoperiod on growth rate and endogenous gibberellins in the long-day rosette plant spinach. Plant Physiol. 47: 821-827.
- 128. Zeevaart, J. A. D. 1971. (+)-Abscisic acid content of spinach in relation to photoperiod and water stress. Plant Physiol. 48: 86-90.
- 129. Zucconi, F. 1975. Reassessment of the relationship between hormonal and developmental changes during abscission with particular reference to peach (Prunus persica L. Batsch) fruit. Ph.D. Dissertation, Michigan State University.
- Zucconi, F., R. Stosser and M. J. Bukovac. 1969. Promotion of fruit abscission with abscisic acid. BioScience, 19: 815-817.

ABSCISSI

]

tive stru

dry beans productio

and perce

from a co to a long Black Tun

had the h

#### CHAPTER 2

## ABSCISSION OF FLOWERS AND FRUITS IN PHASEOLUS VULGARIS L.

# I. VARIETAL DIFFERENCE IN FLOWERING PATTERN AND ABSCISSION

# Abstract

Developmental patterns and abscission of reproductive structures (flowers and pods) were studied in both greenhouse and field experiments with three cultivars of dry beans (<a href="Phaseolus vulgaris">Phaseolus vulgaris</a> L.). The pattern of flower production, pod retention, number of flowers produced and percentage of reproductive structures abscised varied with cultivar. The pattern of flower production varied from a concentrated skewed pattern for cultivar Seafarer to a longer more normally distributed pattern for cultivar Black Turtle Soup. In general, the first-formed flowers had the highest probability of setting pods and producing mature seed.

fruit.

obtained immature

of the re was lost loss var

8), and

needed o

and the paper the growth ;

ditions contras

hopeful

the gen

#### Introduction

The seed yield is determined primarily by the number of flowers per plant, the percentage fruit set, the number of seeds per fruit, and seed size. If abscission could be prevented or decreased, yield of dry beans might be increased, provided other physiological limits were not encountered.

Dry bean, in common with many other crop plants grown for seed, produce many more flowers than mature fruit. One of the reasons that maximum yield is not obtained in this crop is abscission of flowers and immature pods. Swen (9) reported that from 78 to 87% of the reproductive potential in 5 varieties of soybeans was lost by abscission. In field beans the estimated loss varies from a low of 75 to a high of 97% (3, 4, 7, 8), and a figure of 80% was reported in lupins (2).

In order to increase yields, information is needed on flowering patterns, patterns of abscission, and the basic cause or causes of abscission. In this paper the patterns of flowering, abscission, and fruit growth and development under field and greenhouse conditions are described in three dry bean cultivars with contrasting growth habits. The information obtained hopefully will serve as a base for further studies on the genetic and physiological causes of abscission.

I Turtle Sc for their

seeds of the green at 26 ± 5

the seed At flowe uniformi

and tagge each day

At matur and the Variance

appropri Field Ex

at East

double : Were spa

block d

per rep

#### Materials and Methods

## Greenhouse Experiment

Three dry bean cultivars, Seafarer (bush), Black
Turtle Soup (semi-vine) and Michelite (vine) were selected
for their contrasting growth and flowering habits. Four
seeds of each cultivar were planted in 20 35-cm pots in
the greenhouse in early April with temperature maintained
at 26 ± 5°C at East Lansing. Ten days after emergence
the seedlings were thinned to one uniform plant per pot.
At flowering time, 15 of each cultivar were selected for
uniformity and flowers that opened each day were counted
and tagged with color-coded wire, using a different color
each day. These flowers were observed until maturity.
At maturity the number of filled pods, plant dry weight
and the yield components were recorded. Analysis of
variance and correlation analysis were carried out where
appropriate.

# Field Experiment

Ten dry bean cultivars were grown in field plots at East Lansing in the summer of 1974. Each plot was a double row, 15 m long with rows 1.5 m apart. Plants were spaced 50 cm apart in the row. A randomized complete block design was used with 4 replications and 60 plants per replication per cultivar. Flowers from 10 plants

per culti coded wir followed

analyzed

Greenhou

more flo higher r differen

(Table ]

weight,

nonsigni nifican

obtained ponents

but not

is demo

number

but had

number

also ha


per cultivar in each replication were tagged with colorcoded wire each day, and the flowering pattern was followed until maturity.

At maturity, the plants were harvested and analyzed as described above.

#### Results


### Greenhouse Experiment

Seafarer and Michelite produced significantly more flowers than Black Turtle Soup; however, their higher rates of abscission resulted in a nonsignificant difference between cultivars in number of pods retained (Table 1). No cultivar differences occurred in stem weight, pod weight, or total plant weight, but the differences were seen in root weight (Table 1). With the nonsignificant difference in vegetative weight, no significant difference in seed yield among cultivars was obtained (Table 1). Cultivar differences in vield components were found in seeds per pod and weight per seed but not in number of pods or yield (Table 1). In general, the phenomenon of yield component compensation is demonstrated by these data. Michelite had a high number of pods, low seeds per pod and low seed weight but had the same yield as Black Turtle Soup (Table 1). On the other hand, Black Turtle Soup, with a low pod number, high seeds per pod and high average seed weight. also had a yield similar to the others. This suggests



Character		Cultivar	
	Seafarer	Michelite	Black Turtle Soup
No. Flowers/plant	40.9 a	44.9 a	19.0 b
No. Pods/plant	15.2 m	19.5 m	13.7 m
% Abscission	62.8 y	56.6 y	27.9 x
Dry wt./plant (g)			
Stems	5.4 a	4.7 a	3.5 a
Roots	5.1 mn	6.6 n	3.0 m
Pods	16.8 x	16.7 x	20.4 x
Total	27.3 p	31.0 p	27.0 p
#seeds/pod	4.1 a	4.2 a	5.5 b
wt./seed (g)	0.21 y	.16 x	.22 y
seed wt./plant (g)	12.7 m	14.1 m	15.7 m

 $<sup>^{\</sup>rm a}{\rm Means}$  within a given character followed by similar letters are not significantly different at the 1% level by Tukey's Test.




that there is a yield limit set by the plant but that there are several ways to get to that end point via variations in level of yield components.

Seafarer had a short and highly concentrated flowering period (6 days) reaching its peak of flower production in 3 days from beginning of the first flower. Michelite and Black Turtle Soup, on the other hand, had a longer and less concentrated flowering period (15-18 days) with their peak of flower production at 8 days from the opening of the first flower (Fig. 1). To determine which flowers were most likely to set pods and produce seed, the data were expressed as pods produced on a given day that are retained to maturity, as a percentage of the total number of pods harvested (Fig. 2). The peak of pod retention for Seafarer was at 2 and 3 days while its peak of flower production was at 3 days (Figs. 1 and 2). The peak of pod retention for Michelite and Black Turtle Soup was at 4 days while their peaks of flower production were at 8 days.

## Field Experiment

The 10 cultivars studied can be classified into 3 groups, according to the number of flowers produced; those which produced the highest number of flowers were Michelite, 0639, and 0686, those with the next highest were Pinto 114, Seafarer, Black Turtle Soup and Jamapa, and those which produced the lowest number of flowers







Fig. 1. Flower production in a specific day as a percentage of total flowers produced under greenhouse conditions.







Fig. 2. Pods retained from flowers borne on a specific day, as a percentage of total pods at harvest grown under greenhouse conditions.



were Cran-028, Montcalm and Swedish Brown (Table 2). A similar trend was seen among these cultivars in the number of pods retained to maturity, that is, cultivars which produced more flowers had a tendency to retain higher number of pods (Table 2). However, the abscission rate did not follow the trend of flowers and pod production; plants with fewer flowers and fewer pods had the abscission rate as high as those with higher number of flowers and pods (Table 2). Cultivar differences were seen in total plant weight which resulted in the difference in seed weight per plant (Table 2).

The flowering pattern of the three cultivars selected for detailed study indicated Seafarer to have a slight earlier peak of flower production in 7 days from beginning of the first flower than Michelite and Black Turtle Soup (11-12 days) (Fig. 4). The peak of pod retention in Seafarer occurred at 5 days after the first flower which was ahead of that in Michelite and Black Turtle Soup (Fig. 5). In all three cultivars, the flowers that were borne earlier had a higher chance of setting and developing to maturity (Fig. 6).

## Discussion


Abscission data from the field and greenhouse grown plants indicates that there is more abscission in the field (Table 1 and 2). This higher abscission rate under the field conditions in all cultivars may be

Table 2

Mean Values per Space-planted Plants of Flowers Produced, Pods Retained at Harvest and Percentage of Flower and Pod Abscission

Cultivars	Number of Flowers	Number of Pods	% Abscission	Seed wt. (g) Per Plant	Total Plant Wt. (g)
Michelite	281 a	82 ab	71 ab	51 b	94 bed
6290	259 ab	84 a	q 89	44 bc	79 cdef
9890	222 b	73 abc	q 99	55 b	110 ab
Pinto 114	178 c	58 cd	q 99	87 a	131 a
Seafarer	168 c	53 d	q 89	33 c	54 £
Black Turtle Soup	147 cd	68 bcd	52 c	57 b	102 bc
Jamapa	147 cd	64 cd	56 c	51 b	88 bcde
Cran-028	107 de	35 e	q 89	48 bc	72 def
Montcalm	100 e	23 e	76 a	20 p	87 bcde
Swedish Brown	97 e	33 е	q 99	41 bc	63 ef

Letters not in common indicate significant differences by the Duncan's Multiple Range Test (5% level). Note:



10 Percent 9

a pero




Fig. 4. Flower production in a specific day as a percentage of total flowers produced under field conditions.




Fig. 5. Pod retained from flowers borne on a specific day, as a percentage of total pods at harvest grown under field conditions.

attribut condition diseases Also it much gree to that high ra associa on fiel between and pod

> absciss vars u

Apart f cultiva growing the lea Michel:

due to were g time (

(10). Vars v

farer of Mi

attributed to greater variations in the environmental conditions such as moisture stress, heat, wind, pest and diseases to which the plants were occasionally subjected. Also it was noticed that under the field conditions a much greater number of flowers were produced as compared to that of greenhouse grown plants (Table 1 and 2). A high rate of abscission in the field probably can be associated with the greater number of flowers produced on field-grown plants. A high negative correlation between the yield per plant and the percentage of flower and pod abscission was reported in garden beans (1). Apart from the higher abscission rate under the field. cultivar differences in abscission existed under both growing conditions. Black Turtle Soup, which produced the least number of flowers, had the lowest while Michelite and Seafarer had the highest percentage of abscission (Table 1 and 2).

The pattern of flowering was longer in all cultivars under the field-grown plants as compared to that of the greenhouse-grown plants (Fig. 1 and 4). This may be due to the longer daylength in summer when the plants were grown in the field, and better growth rates at that time (Table 1 and 2), the same as was shown in soybean (10). The patterns of pod retention for the three cultivars were similar under both greenhouse and field. Seafarer had an earlier peak for pod retention than those of Michelite and Black Turtle Soup. This pattern of pod

retenti duced f ripe po phenome

6). The flowers develop

or by

mately These

growth

in eit

and ph are no also )

to it

retention peaks indicates that the flowers that are produced first are most likely to be retained to produce ripe pods containing good seeds (Fig. 3 and 6). This phenomenon was also shown in lupin (11) and cowpea (5, 6). This may be explained either by assuming that early flowers have the first share of nutrients for their development, thus having a greater chance of survival, or by the release of an inhibitory substance to inhibit flowers set at a later time.

Although Seafarer and Michelite differ widely in growth habit and flowering patterns, they share approximately 92% of their germplasm in common (Adams, per com.). These two cultivars have similar percentage of abscission in either greenhouse or field-grown conditions (Table 1 and 2). This indicates that there may be separate genetic and physiological factors controlling abscission which are not affected by growth habit. These factors must also be considered in breeding to better fit the plant to its environment especially when gross changes in plant architecture are proposed.

Black Turtle Soup Seafarer Michelite

1.00




Fig. 6. Pods retained per day as a percentage of flowers produced on that day (data from field-grown plants).

l. Bin

2. Gre

3. Ho

. Ka

5. Oj

6. Oj

7. Ro

8. Sc

#### REFERENCES

- Binkley, A. M. 1932. The amount of blossom and pod drop on six varieties of garden beans. Proc. Amer. Soc. Hort. Sci. 29: 489-492.
- Greenwood, E. A. N., P. Farrington and J. D. Beresford. 1975. Characteristics of the canopy, root system and grain yield of a crop of <u>Lupinus</u> angustifolius cv. Unicrop. Aust. J. Agric. Res. 26: 497-510.
- Hodgson, G. L. and H. E. Blackman. 1956. An analysis
  of the influence of plant density on the growth
  of <u>Vicia faba</u>. I. The influence of density on
  the pattern of development. J. Expt. Bot. 7:
  147-165.
- Kambal, A. E. 1968. A study of the agronomic characters of some varieties of <u>Vicia faba</u> Sudan Agric. J. 3: 1-10.
- Ojehomon, O. O. 1970. Effect of continuous removal of open flowers on the seed yield of two varieties of cowpea, Vigna unguiculata (L) Walp. J. Agric. Sci. 74: 375-381.
- 6. Ojehomon, O. O. 1972. Fruit abscission in cowpea, Vigna unguiculata L. Walp. 1. Distribution of 14C-assimilates in the inflorescence, and comparative growth of ovaries from persisting and abscising open flowers. J. Expt. Bot. 23: 751-761.
- Rowlands, D. G. 1961. Fertility studies in the field bean (Vicia faba L.) II. Inbreeding Heredity 16: 497-508.
- Soper, M. H. R. 1952. A study of the principal factors affecting the establishment and development of the field bean (<u>Vicia</u> <u>faba</u>) J. Agric. Sci., Camb. 42: 335-346.

9. Sw

10. Va

ll. Va

- Swen, M. A. 1933. Factors affecting flower shedding in soybean. Ph.D. dissertation, University of Illinois, Urbana.
- Van Schaik, P. H. and A. H. Probst. 1958. Effects of some environmental factors on flower production and reproductive efficiency in soybeans. Agron. J. 50: 192-197.
- Van Steveninck, R. F. M. 1957. Factors affecting the abscission of reproductive organs in yellow lupins (<u>Lupinus luteus L.</u>). I. The effect of different patterns of flower removal. J. Expt. Bot. 8: 373-381.

ABSCI

bean (

The nu

(10 da

Weight

greate

under

#### CHAPTER 3

ABSCISSION OF FLOWERS AND FRUITS IN PHASEOLUS VULGARIS L.

II. GROWTH AND POD ABSCISSION AS AFFECTED BY

DIFFERENT DAYLENGTHS

#### Abstract

Greenhouse-grown plants of two cultivars of dry bean (Phaseolus vulgaris L.) were subjected to 8, 13, and 18-hour photoperiods after the flower buds became visible. The numbers of flowers that opened on each plant were counted daily. Plants were harvested at peak flowering (10 days after initiating photoperiodic treatment), at pod filling (20 days), or at maturation (40 days). Weights of roots, stems, leaves, fruits and seeds were recorded, and leaf areas determined. Pod abscission was greater under short photoperiods and differed for the two cultivars. Plants were larger and yielded more seed under longer daylengths.

as the damage that soil most plant no mechani photosy in cott Factors plant damage that da

ontoge that p plants to mea

ation can be

fruit area

photo:

## Introduction

A number of environmental factors have been cited as the cause of abscission of reproductive structures.

Among these factors are extremes of temperature, humidity, soil moisture, light duration and intensity, inadequate plant nutrition, disease, insect infestation, and mechanical forces such as wind and rain.

Abscission may be promoted by a deficiency of photosynthate. For example, shading increases abscission in cotton (5, 6, 7), soybean (3, 11) and vining peas (13). Factors which reduce the amount of light reaching the plant canopy, such as cloudy weather (9, 12) and close spacing (2) increase abscission in young cotton fruit (6, 12), and soybean (10). Partial or complete defoliation also increases abscission. Most of these results can be attributed to a deficiency of photosynthate.

Abscission can occur at several stages in ontogeny. However, Meadley and Milbourn (13) showed that pod abscission in pea was markedly increased when plants were shaded from flowering onwards. In order to meet the increased demand for photosynthate during fruit development either photosynthetic rates or leaf area must be increased. In soybean the rates of leaf photosynthesis are much higher during the pod filling stages (4, 8).

effect daylen

Turtle
in flo

greenh days a plant

of eac

was me

plant

divid under

house Were

cons

With incal

plan

The purpose of this study was to measure the effects on growth and pod abscission of differential daylengths.

## Materials and Methods

Two dry bean cultivars, "Seafarer" and "Black
Turtle Soup" were chosen for study because they differed
in flowering habit and rate of pod abscission (Chapter 2).
Three seeds were planted in each of 150 35-cm pots in the
greenhouse in early April, 1975 at East Lansing. Ten
days after emergence the seedlings were reduced to one
plant per pot. At flower initiation 10 uniform plants
of each cultivar were harvested, and separated into root,
stem (including petioles), and leaf. Leaf area per plant
was measured and, after freeze drying of all plant parts,
dry weights were determined.

At the time of flower initiation 90 uniform plants of each cultivar were selected, and randomly divided into three groups of 30 plants, which were placed under 8, 13, or 18-hour photoperiods in the same greenhouse. For the 8-hour day, black polyethylene curtains were opened at 8 a.m. and closed at 4 p.m. each day throughout the experiment. The 13- and 18-hour treatments consisted of natural light plus supplemental lighting with a mixture of high intensity from fluorescent and incandescent lights. The intensity of the light at the plant level was 70,000-75,000 lux, ample for maximum

photosy to avoi was mai

night.

opened flowers ing (10

experion harves

at pod

ments,

(

(

the s

Howe

photosynthesis. A black screen separated the treatments to avoid any interference. The greenhouse temperature was maintained at 25-29°C in the daytime and 18-21°C at night.

All plants were numbered, and the flowers that opened each day were recorded to give the number of flowers produced. Plants were harvested at peak flowering (10 days after the beginning photoperiodic treatment), at pod filling (20 days), and at maturity (40 days). The experiment was analyzed factorially with three factors, harvesting time, cultivar, and daylength. In each harvest, 10 plants per cultivar were used for all treatments, and the parameters measured in each harvest were:

- Number of pods and number of seeds per plant (second and third harvest only);
- (2) Number of trifoliate leaves per plant (first and second harvest only, since leaves had senesced on mature plants);
- (3) Leaf area per plant (first and second harvest only);
- (4) Root, stem (including petiole), leaf, seed and pod dry weight.

The above experiment was repeated in 1976, using the same greenhouse, cultivars and lights as before. However, 25 cm pots were used instead of 35 cm pots.

Pod Abs

more population percent

show the hours ever,

ever,

days a

of pod the 8stage

In ger from 8 culti

decrea Absci

inclusion

Well

flowe

### Results and Discussion

#### Pod Abscission

Both cultivars produced more flowers and retained more pods as daylength increased (Table 1). In 1975, percent pod abscission was markedly reduced in Black Turtle Soup under 13-hour day, while Seafarer did not show this reduction until daylength was extended to 18 hours (cultivar x daylength significant at 1%). However, no interaction was evident in 1976.

Pod abscission was not recorded at peak flowering since the pods were still attached to the plants. However, at pod filling many pods had abscised. More than one-half of the pods lost having fallen in the first 20 days after anthesis in both cultivars. In Seafarer a greater portion of the pods abscised at the early stages of pod development (Table A-1, A-2, Appendix A). Under the 8-hour day many pods abscised at the pod filling stage in both cultivars (Table A-1, A-2, Appendix A). In general, considering both years increasing daylength from 8 to 13 to 18 hours reduced pod abscission in both cultivars (Table 1). At the mature stage pod abscission decreased progressively as daylength increased (Fig. 1A). Abscission data recorded at the pod filling stages included both late flowers and young pods. The abscission of these structures may involve hormonal levels as well as nutrients. Similarly, shading of peas after flowering increased pod abscission (12).

ole 1

Table 1

Table 1

Effects of Daylengths and Harvesting Time on Number of Plowers, Number of Pods, and Percentage of Pod Absolssion of Two Dry Bean Cultivars Grown Inder Greeninouse Conditions in 1975 and 1976 Absolssion of Two Dry Bean Cultivars Grown Inder Greeninous Conditions in 1975 and 1976

		Number of Flowers	Flowers	Number of Pods	of Pods	% Abscission	ission	
		1975	1976	1975	1976	1975	1976	
i	Harvesting Time							
	Peak flowering	17.2 a	11.5 a	16.4 c	10.4 €	1	1	
	Pod filling	17.4 a	13.6 b	9.9 b	7.3 b	36.2 a	42.1 a	
	Maturity	18.7 a	12.8 ab	7.6 a	5.5 a	53.7 b	54.1 b	
II.	Cultivar							
	Black Turtle Soup	10.8 m	8.3 m	8.6 m	6.3 m	30.5 m	38.2 m	
	Seafarer	24.7 n	17.0 n	14.0 n	9.2 n	59.4 n	62.0 n	
III.	Daylengths							
	8 Hours	10.1 ×	10.4 x	x 6.5		53.7 z	57.9 z	
	13 Hours	18.3 y	12.4 y	10.2 y	7.7 Y	47.4 y	46.8 y	
	18 Hours	24.8 z	15.2 z	17.8 z		33.7 x	39.7 x	
Inte	Interactions							
	I × II	n.s.	n.s.	*	*	n.s.	n.s.	
	I x III	n.s.	n.s.	*	*	*	n.s.	
	II × III	*	n.s.	*	n.s.	*	n.s.	
	IX II X III	n.s.	n.s.	*	n.s.	n.s.	n.s.	

 $^{4}_{\text{Means}}$  within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.




Fig. 1. Interaction between developmental stages and daylengths on percentage of pod abscission (A), and total plant weight (B). (Data taken from 1975, averaged over two cultivars.)

pods a hour d abscis

in Bla

under correl abscis

with h photop negati

negat: under

veget.

the co

between istic:

signi of bo

for b

titio

Competition for nutrients between flowers, young pods and growing apices was probably severe under the 8hour day. The correlation between percentage of pod abscission and number of flowers produced was significant in Black Turtle Soup under 8- and 13-hour days but not under the 18-hour photoperiod (Table 2). The positive correlation agrees with data for garden bean in which abscission of flowers and pods is greater in varieties with high flower number (1). However, under the 18-hour photoperiod the number of flowers borne per plant was negatively correlated with abscission (Table 2). The negative correlation was insignificant in both cultivars under the 18-hour day. Because Seafarer is determinate. vegetative growth competed less with fruit set than was the case in the indeterminate Black Turtle Soup. Thus the correlation between abscission and flower number was significant only at the 13-hour photoperiod (Table 2).

The preponderance of negative relationships between percentage of pod abscission and the characteristics representing vegetative growth at the 18-hour photoperiod, though individually not always statistically significant (Table 2) nevertheless implies that the plants of both cultivars are producing photosynthate sufficient for both an increase in growth and a removal of competition between developing pods.

Table 2

Correlation Coefficients Between Percentage of Pod Abscission or Leaf Area and Other Parameters in Two Dry Bean Cultivars Subjected to Different Daylength Treatments After Flowering (date taken at the pod filling stage from 1975 grown plants)

40	Bla	Black Turtle Soup	dnos		Seafarer	
Faraniecers	8 Hours	13 Hours	18 Hours	8 Hours	13 Hours	18 Hours
% Pod Abscission vs:						
Flowering period	.22	.28	.28	20	.28	.25
Number of flowers	.71ª	.74b	24	.34	.61a	04
Number of pods	60.	.29	47	57	26	730
Number of seeds	d77.	.58	-,31	.23	.31	78 <sup>D</sup>
Seed weight	.64a	. 59	.18	. 47	. 49	20
Root weight	.53	.27	55	.13	.13	33
Stem weight	.44	.61a	24	03	.21	42
Leaf weight	.61ª	.40	43	13	.30	38
Leaf number	.69ª	.46	19	26	90.	04
Leaf area	.51	.57	51	18	.17	40
Leaf Area vs:						
Flowering period	.73b	.13	.13,	.60ª	.33	.43
Number of flowers	.740	.69	.74.	.755	.48	200
Number of pods	·67ª	.51	۵64.	. 780	.51	≈/8.
Number of seeds	.67a	g88.	.61a	.64ª	g68.	.49
Seed weight	.69ª	.65ª	90.	.55	.72ª	.50

<sup>a</sup>Significant at 5% level.

bsignificant at 1% level.

cantly

nifica both o

photop nifica days

nific for s

Plant

dayle

A-5,

### Leaf Development

Leaf number, area, and weight increased significantly with daylength (Table 3). Leaf area and leaf weight per plant were significantly greater in Black Turtle Soup than in Seafarer in 1975; only the latter difference was significant in 1976. However, the number of leaves per plant was not affected by cultivar, documenting the larger leaf size in the Black Turtle Soup (Table 3). Leaf number and area per plant did not change significantly with time in 1975 (Table 3) indicating that no leaf growth occurred after the pods were set.

Leaf area was positively correlated with yield components in all daylengths (Table 2). This is consistent with many reports that greater photosynthetic capacity results in higher yield. Leaf area was significantly correlated with length of flowering time in both cultivars under 8-hour day, but not under longer photoperiods (Table 2). Yield (seed weight) was significantly correlated with leaf area under 8 and 13-hour days in Black Turtle Soup (Table 2). However, no significant correlation existed between leaf area and yield for Seafarer plants under 8 and 18-hour days.

# Plant Development

Weight of all vegetative parts increased with daylength in both cultivars and in both years (Table 4, A-5, and A-6). This was associated with a greater number

Table 3

Effects of Daylengths, Harvesting Times and Cultivars on Leaf Weight Per Plant, Leaf Area Per Plant and Number of Trifoliate Leaves Per Plant Grown Under Creanhouse Conditions in the Year 1975 and 1976 (data represents means for treatment) a

		Leaf wt/	Leaf wt/Plant (g)	Leaf Area/	Leaf Area/Plant (cm2)	Number of Trifo	Number of Trifoliate Leaf/Plant
		1975	1976	1975	1976	1975	1976
i.	Harvesting Time						
	Peak flowering Pod filling	3.7 a	2.3 a 2.5 a	1394 a 1472 a	991 a 1131 b	17 a 16 a	12 a 13 b
II.	Cultivar						
	Black Turtle Soup Seafarer	4.7 n 3.6 m	2.5 m 2.3 m	1624 n 1242 m	1149 n 974 m	16 a 17 a	12 m 13n
III.	Daylengths						
	8 Hours 13 Hours 18 Hours	2.9 x 3.8 Y 5.7 z	1.6 x 2.1 y 3.5 z	1172 x 1460 y 1667 z	839 x 1026 y 1320 z	14 x 16 y 19 z	10 x 12 y 15 z
Inte	Interactions						
	IxI	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
	IXI	n.s.	*	n.s.	n.s.	n.s.	n.s.
	IX IX III	n.s.		n.s.	n.s.	n.s.	n.s.

Amens within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.

with :

plant ing; while

plant filli weigh

> after sink

and s

organ

orga: phot

Seed

plan leng

plar (Tak

Turt

and percentage of pods retained (Table 1). Greater vegetative growth (Table 3 and 4) was also associated with lower abscission (Table 1) when 1975 values for Black Turtle Soup are compared with those for Seafarer.

The weight of the vegetative portions of the plants in this experiment did not increase after flowering; root and stem weights tended to decrease (Table 4), while leaf weight remain unchanged (Table 3). Total plant weight increased from peak flowering to the pod filling stages (Table 4), as a result of increasing pod weight. It could be assumed from these results that, after flowering, pod development comprises such a strong sink for assimilate (or nutrients) that pods are effective and successful competition with vegetative organs. A more likely explanation is simply that the vegetative organs have reached their final size, as determined by genetic potential in this environment, and that these organs are, in fact, no longer competing for the available photosynthate (or nutrients) with the pods.

# Seed Yield

The number of normal-sized seeds produced per plant was greater for both cultivars under longer day-length (Table 5), resulting in greater seed weight per plant. Weight per seed was not affected in 1976 (Table 5); effects varied with cultivar in 1975. Black Turtle Soup seeds were larger under 13 and 18 hour than

Table 4

Effects of Daylengths, Harvesting Times and Cultivars on the Weight per Plant of Root, Stem and Total Plant Grown Under Greenhouse Conditions in the Year 1975 and 1976 (data represents means for treatment) a

		Root wt/	Root wt/Plant (g)	Stem wt/	Stem wt/Plant (g)	Total Pla	Total Plant wt (g)
		1975	1976	1975	1976	1975	1976
ij	Harvesting Time						
	Peak flowering			2.0 b		7.2 a	4.8 a
	Pod filling Maturity	1.1 b 0.9 a	1.0 a 0.9 a	2.0 b	1.5 b	12.5 c 11.3 b	8.3 b
ii.	Cultivar						
	Black Turtle Soup Seafarer	1.4 n 1.1 m	0.9 m	1.9 n 1.7 m	1.2 m	11.0 n 9.6 m	6.4 m
III.	Daylengths						
	8 Hours						3.9 x
	18 Hours	1.0 y 2.0 z	0.8 y 1.5 z	1.6 y 2.7 z	1.1 y 1.9 z	9.2 y 16.1 z	6.4 y 10.3 z
Inte	Interactions						
	I × II I × III	n *	n.*	* *	. s.		 
	II × III I × II × III	n.s.	* u	* u	n.s.	. s.	n.s.

 $^{3}$ Means within the same parameter followed by similar letters are not significantly different at the 18 level by Duncan's Multiple Range Test.

Table 5

Effects of Daylength on Number of Seeds per Plant, Total Seed Weight per Plant and Weight per Seed in Two Dry Bean Cultivars Grown in 1975 and 1976 Seasons<sup>a</sup>

		No. See	No. Seed/Plant	Total S	Total Seed Wt. (9)	Wt/Se	Wt/Seed (g)
		1975	1976	1975	1976	1975	1976
i.	Cultivar						
	Black Turtle Soup Seafarer	38.8 a 35.0 a	25.9 a 23.7 a	5.4 4.8 a	5.1 a 5.7 a	0.22 a	0.14 a
II.	Daylength (hrs.)						
	8 13 18	15.3 x 28.9 y 66.4 z	11.5 x 26.6 y	2.4 4.9 x	2.7 x x z z x x	0.23 x 0.24 x	0.16 x 0.17 x
Inte	Interaction			2	7 7.0	× 77.0	X 97.0
	I x II	n.s.	n.s.	n.s.	n.s.	*	n.s.

under for S

> growt absci

ditio

under 8-hour daylengths, whereas the reverse was true for Seafarer.

Within the same variety, plants having greater growth before flowering usually associate with lower abscission level. Also favorable environmental conditions after flowering are required for seed development and fruit retention.

7.

8.

9.

10.

#### REFERENCES

- Binkley, A. M. 1932. The amount of blossom and pod drop on six varieties of garden beans. Proc. Amer. Soc. Hort. Sci. 29: 489-492.
- Brown, K. J. 1971. Plant density and yield of cotton in northern Nigeria. Cotton Growing Rev. 48: 255-266.
- Cartter, J. L. and E. E. Hartwig. 1962. The management of soybeans. Adv. in Agron. 14: 360-412.
- Dornhoff, G. M. and R. M. Shibles. 1970. Varietal differences in net photosynthesis of soybean leaves. Crop Sci. 10: 42-45.
- Dunlap, A. A. 1943. Low light intensity and cotton boll shedding. Science 98: 568-569.
- Eaton, F. M. and D. R. Ergle. 1954. Effects of shade and partial defoliation on carbohydrate levels and the growth, fruiting and fiber properties of cotton plants. Plant Physiol. 29: 39-49.
- Eaton, F. M. and N. E. Riglez. 1945. Effect of light intensity, nitrogen supply, and fruiting on carbohydrate utilization by the cotton plant. Plant Physiol. 20: 380-411.
- Ghorashy, S. R., J. W. Pendleton, R. L. Bernard and M. E. Bauer. 1971. Effect of leaf pubescence on transpiration, photosynthetic rate, and seed yield of three near-isogenic lines of soybeans. Crop Sci. 11: 426-427.
- Goodman, A. 1955. Correlation between cloud shade and shedding in cotton. Nature 176: 39.
- McAlister, D. F. and O. A. Krober. 1958. Response of soybeans to leaf and pod removal. Agron. J. 50: 674-677.

11.

12.

13.

- Mann, J. D. and E. G. Jaworski. 1970. Comparison of stresses which may limit soybean yields. Crop Sci. 10: 620-624.
- Mason, T. G. 1922. Growth and abscission in Sea Island cotton. Ann. Bot. 36: 457-483.
- Meadley, J. T. and G. M. Milbourn. 1971. The growth of vining peas. III. The effect of shading on abscission of flowers and pods. J. Agric. Sci. 77: 103-108.

ABSC:

(Phas

repro

seed during with

tion

the Turt

free

stem

#### CHAPTER 4

ABSCISSION OF FLOWERS AND FRUITS IN PHASEOLUS VULGARIS L.


III. RELATIONSHIP OF POD ABSCISSION TO DIFFERENTIAL

LEVELS OF CARBOHYDRATES PRODUCED BY


VARIATION IN DAYLENGTH

## Abstract

"Black Turtle Soup" and "Seafarer" dry bean (Phaseolus vulgaris L.) cultivars were subjected to 8, 13 and 18-hour photoperiods in the greenhouse during their reproductive phase. Free sugar, starch, and total nonstructural carbohydrate concentration of root, stem, leaf, seed, and pod wall were determined at different stages during pod development. Free sugar concentration increased with photoperiod in all tissues. Higher starch concentration was found in all tissues except in the root where there was no difference in starch concentration between the short (8-hour) and the long (18-hour) days. Black Turtle Soup, an indeterminate cultivar, contained more free sugar in all tissues than Seafarer, a determinate cultivar. Black Turtle Soup contained more starch in stem and leaf, but not in root and seed. Free sugars in root and leaf rose from peak flowering to pod filling, then



decreased. However, free sugars in stem, seed, and pod wall declined as the pod matured. In general, starch concentration of vegetative organs (root, stem, and leaf) decreased as the plants matured, while that of the seeds increased. The general pattern of higher available carbohydrate under long photoperiods was associated with greater vigor, higher yield, and reduced pod abscission. Pod abscission data indicated an inverse relationship to available carbohydrate concentration in vegetative organs.



### Introduction

Seed yield is determined by the number of flowers produced, the percentage of fruit set, the number of seeds per fruit, and seed size. Flower and young fruit abscission is extensive in many economic crops (4, 6, 14, 15), and is often related to environmental conditions, especially those that influence photosynthesis (3).


Abscission of young fruits may be controlled by levels of nitrogenous or carbohydrate reserves (5, 16). Spraying cotton plants with 20% sucrose and 1% nitrogen as urea actually increased abscission (1). However, the endogenous carbohydrate level was not influenced. Girdling increased carbohydrate level, and a 2% urea spray increased the nitrogen level (1). However, combining the two treatments considerably increased boll abscission. These results raise a question on the role of carbohydrate in fruit abscission (1).

In previous work, pod abscission was reduced and seed yield increased in dry beans by exposure to long photoperiods during the reproductive phase (see Chapter 3). The present study was designed to determine the effect of photoperiod on levels of available carbohydrates.

### Materials and Methods

### Sample Preparation

Two dry bean cultivars, "Seafarer" and "Black
Turtle Soup," were chosen because of differences in plant



type, flowering habits, and pod abscission rates. Seeds were planted in 35-cm pots in the greenhouse at East Lansing in early April 1975. Growing conditions have been described previously (see Chapter 3).

Plants were harvested at (a) flower initiation, just before beginning photoperiod treatments, (b) peak flowering, 10 days after flower initiation, (c) pod filling, 20 days after flower initiation, and (d) maturity, 40 days after flower initiation. At each harvest 10 uniform plants of each cultivar were separated into root, stem (including petioles), leaf, pod, and seed. The samples were freeze-dried, weighed, ground in a Wiley mill having a 40-mesh sieve, and stored in scintillating vials at room temperature prior to extraction.

# Determination of Free Sugars


The method used was a modification from Laidlaw and Reid (7). Free sugars were extracted by refluxing 100 mg samples in 80% ethanol in a water bath at 60°C for two hours, with occasional shaking. Samples were filtered through Whatman #1 paper and the filtrate was evaporated under vacuum at 50°C to near dryness. Volume was adjusted to 25 ml with distilled water. The solution was clarified with 10% lead acetate and then deleaded with potassium oxalate (8). Volume was adjusted, to 50 ml in the case of root and pod wall and 100 ml for

stem, leaf and seed tissues, with distilled water prior to analysis.

For analysis of free sugars, 5 ml of 70% anthrone reagent (19) was added to each 1.0 ml of extract. Before and after the addition of anthrone reagent the tube was immersed in cold water and shaken to obtain thorough mixing. The solution was heated in boiling water for 10 minutes, cooled to room temperature and absorbance was determined at 620 nm. Sugar content was expressed as glucose equivalents per gram dry weight.

#### Determination of Total Nonstructural Carbohydrates (TNC)

The extraction employed was modified from Smith (11). One hundred mg of tissue was placed in a 45 by 400 mm test tube, 15 ml distilled water was added, and the solution was boiled for 5 minutes to gelatinize the starch. After cooling to room temperature, 10 ml of buffer, pH 4.45, and 10 ml 0.5% takadiastase (Clarase 900) enzyme solution were added. The tubes were shaken in a water bath at 45°C for 44 hours (17). The solution was filtered through Whatman #1 paper into a 125 ml Erlenmeyer flask, and 2 ml of 10% neutral lead acetate was added to precipitate the proteins. After centrifugation at 12,000 xg for 5 minutes, the supernatant fluid was decanted into a second 125 ml flask containing 100 mg powdered potassium oxalate. These mixtures were kept



overnight in the refrigerator. The solution was stored overnight at 4°C, then centrifuged at 12,000 xg for 10 minutes, filtered through Whatman #52 paper, and made up to 200 ml with distilled water. TNC was determined colorimetrically at 620 nm using the anthrone-reagent technique as described earlier (19), and expressed as mg glucose per gram dry weight.

Since TNC in legumes consists mainly of free sugars and starches (11, 12), starch concentration was estimated as the difference between TNC and soluble sugars.


The data were analyzed factorially using 10 plants  $\label{eq:plant} \text{per treatment (one plant per replication).}$ 

# Results and Discussion

# Free Sugar

In general, free sugar levels increased with photoperiod in root, stem, leaf, seed and pod wall (Table 1). However, in the leaf highest levels occurred in plants subjected to 13-hour days (Table 1). Under the 18-hour daylength the movement of assimilate out of the leaf might have been more rapid, resulting in less free sugar in the leaf.

Free sugars were higher in all tissues of Black
Turtle Soup than in Seafarer (Table 1), indicating either
a difference in rate of assimilation of carbon dioxide
or a difference in rate of translocation of sugar, or




Developmental Stages, Cultivars and Daylength Effects on Free Sugar Level in Various Plant Parts^a (mg. glucose per g. dry weight) Table 1

I. Time Peak Pod f	Time Peak flowering Pod filling	Root	4	0+0			١	6000		200	
I. Tim Pea Pod Mat	le lk flowering   filling		,	2 - 61		Leaf		מפני		Pod	
Pea Pod Mat	k flowering										
Pod	filling	12.4		74.6	υ	55.5	ø	1		176 c <sup>D</sup>	
Mat		17.1	υ	62.7	q	66.1	р	9.99	В	141 b	
	Maturity	13.7		34.9	ಡ	ı		71.5	В	26.1 a	ď
II. Cul	Cultivar										
	ningt Purtle Soun	16.2	E		E	68.8	ш	70.2		122.4 m	E
Sea	Seafarer	12.7 n	ď	47.0	ц	52.8	п	67.8	E	106.8 n	¤
III. Day	Daylength										
		12 1	>	45.4	×	52.8	×	26.7	×		×
8	8 hours	10	4 >	61.2	۱ >	67.2	2	6.99	×		×
18	13 hours 18 hours	16.3	n N	65.7	, N	62.4	>	83.3	N	126.4 y	⊳
Interactions	suo					4		*		*	
× +	II.	*		* -		. +		*		*	
: > + ⊢	1 7 111	*		k k		. +		9		*	
II	II X II	*		* *		*		n.s.		*	

 $a_{\rm Means}$  within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.

b<sub>Seed</sub> not separate from pod.



both. Differences in rates of translocation of  $^{14}\mathrm{C}$  assimilates among bean cultivars have been reported (18).

Differences in response of the two cultivars to increasing daylength was evident in free sugar accumulation (Table 2). Free sugar concentration of root, stem, and leaf was proportional to photoperiod in Black Turtle Soup, but not in Seafarer (Table 3). In the latter, the effects of photoperiod were nonsignificant in root, stem. and leaf sugars which were not increased by extending the photoperiod from 13 to 18 hours. This could be due to the differences in growth habit, for Black Turtle Soup plants continue growth after the reproductive stage is attained. Seafarer, on the other hand, is determinate and makes only slight additional growth after the reproductive phase has been reached. This may have reduced free sugar content under longer photoperiods (Table 3). Cultivar did not affect free sugar levels in seed and pod wall except under the 18-hour photoperiod (Table 3). This may be due to the greater ability of Black Turtle Soup to accumulate free sugars in these organs under longer daylength.

In roots of Black Turtle Soup free sugar rose from flower initiation to pod filling, then decreased at maturity (Table 2). However, in Seafarer, free sugar levels did not change between pod filling and maturity. The accumulation of free sugar in roots at the later





Table 2

Effects of Cultivar and Time of Sampling on Free Sugar Concentrations (mg glucose/g dry weight) in Bean Tissues^a

Time After				Tissue		
Flower Initi- ation (days)	Cultivar	Root	Stem	Leaf	Seed	Pod Wall
0	Black Turtle Soup	11.5 a	117.1 9		1 1	
	Seafarer	13.2 b	55.1 cd	44.1 a	ı	ı
	Mean	12.3 p	86.1 s	73.8 r	1	
•	מויסף פון+זיים לפינת	14.9 b	88.4 £	61.7 c	1	187 d
07	Seafarer	9.9 a	60.9 d	49.3 a	1	166 c
	Mean	12.4 p	74.6 r	55.5 p	1	176 r
		000	9 8 97	75.9 d	73.3 b	158 c
20	Black Turtle Soup	14.0 p	48.6 C	56.3 b	59.9 a	124 b
	Mean	17.1 q	62.7 q	66.1 q	d 9.99	141 q
		12 2 4	38.6 h	1	67.1 ab	22.7
40	Black Turtle Soup	14.1 b	31.3 a	1	75.8 b	
	Seararer	13.7 r	34.9 p	1	71.5 p	26.1 p


 $^{\rm A}_{\rm Means}$  within columns and sets (abc, pqr) with similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.



Effects of Photoperiod and Cultivar on Free Sugar Concentration (mg/g dry wt) in Bean Tissues  $^{\rm A}$ Table 3


Black Turtle Soup 12.3 a 53.6 b Seafarer 12.0 a 32.7 a Mean 12.1 x 45.4 x Black Turtle Soup 16.4 b 71.5 c Seafarer 13.4 a 50.8 b Mean 14.9 y 61.2 y Black Turtle Soup 19.8 c 78.7 d Seafarer 12.7 a 52.8 b Mean 16.3 z 65.4 z	Photoperiod	+[]			Tissue		
Black Turtle Soup 12.3 a 53.6 b Seafarer  Mean  12.1 x 45.4 x  Black Turtle Soup 16.4 b 71.5 c Seafarer  Mean  14.9 y 61.2 y  Black Turtle Soup 19.8 c 78.7 d Seafarer  13.4 a 52.8 b  Mean  16.3 z 65.4 z	(hours)	cartivar	Root	Stem	Leaf	Seed	Pod Wall
Mean 12.1 x 45.4 x Black Turtle Soup 16.4 b 71.5 c Seafarer 13.4 a 50.8 b Mean 14.9 y 61.2 y Black Turtle Soup 19.8 c 78.7 d Seafarer 12.7 a 52.8 b Mean 16.3 z 65.4 z	8	Black Turtle Soup Seafarer	12.3 a 12.0 a		58.3 b 47.4 a	94.5 ab 87.7 a	115 a 106 a
Black Turtle Soup 16.4 b 71.5 c Seafarer  Mean  14.9 y 61.2 y Black Turtle Soup 19.8 c 78.7 d Seafarer  Mean		Mean	12.1 x	45.4 x	52.8 x	91.9 x	110.5 x
Mean 14.9 y 61.2 y Black Turtle Soup 19.8 c 78.7 d Seafarer 12.7 a 52.8 b Mean 16.3 z 65.4 z	13	Black Turtle Soup Seafarer	16.4 b		76.9 d 57.4 b	103.8 bc 101.6 b	110 a 103 a
Black Turtle Soup 19.8 c 78.7 d Seafarer 12.7 a 52.8 b Mean 16.3 z 65.4 z		Mean	14.9 Y		67.2 z	102.7 Y	106.5 x
16.3 z 65.4 z	18	Black Turtle Soup Seafarer	19.8 c 12.7 a	78.7	71.2 c 53.5 b	128.8 d 112.7 c	142 b 111 a
		Mean	16.3 z	65.4 z	62.4 Y	120.8 z	126.5 y

 $^{\rm a}{\rm Means}$  within columns and sets (abc, xyz) with similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.



stages of pod development in Seafarer may imply that there is a genetically programmed characteristic of this variety, or that root sink strength exceeds stem or pod strength. The level of free sugar in stem decreased as the pods matured in both cultivars (Table 2). Free sugar in leaves increased from peak flowering to pod filling (Table 2). The marked reduction in leaf free sugar from flower initiation to peak flowering stages in Black Turtle Soup may be due to the movement of sugar from the leaves to the flowers. The decreased abscission of flowers and young pods at peak flowering of Black Turtle Soup as compared to Seafarer (Chapter 3) may be related to the drop in leaf sugar level at this stage. Levels of free sugar in seeds were not affected by cultivar, but fell rapidly between peak flowering and pod filling (Table 2 and 3). In both cultivars the sugar level in pod walls rapidly decreased as the seed matured (Table 2). This may reflect redistribution from the pod walls to the growing seeds, although a decreased rate of photosynthesis in aging pod walls may also be held accountable.

Interactions between photoperiod and developmental stages were noted for sugar concentration of root, stem, leaf, seed and pod wall (Table 4). In the root, at peak flowering and at pod filling, free sugar level increased as photoperiod was extended to 13-hour, but no further increase occurred at 18-hour (Table 4). Levels were




Effects of Photoperiod and Time of Sampling on Free Sugar Concentration (mg/g dry weight) of Bean Tissues  $^{\rm A}$ Table 4

Photoperiod	Time After			Tissue		
(hour)	riower initi- ation (days)	Root	Stem	Leaf	Seed	Pod Wall
œ	O			73.8 c	,	۱
	10					159 cd <sup>2</sup>
	20		47.3 C	61.5 b		145 c
	40	11.1 ab	21.3 a	1	64.3 b	26.8 a
	Mean	12.2 x	55.6 x	29.8 x	26.7 x	110.3 ×
	c	4c 5 CT	2 1 38		1	1
7	,	12.0 00		200	•	174 d
	TO	13.0 cd		0 10	4 0 1	122 b
	20	18.5 e		13.5 C	0.70	
	40	12.5 abc	35.3 b	1	72.9 bc	
	Mean	14.2 y	67.4 y	69.3 z	⊼ 6.99	106.7 ×
81	c			73.8 c	1	1 9
)	,	200	100		1	195 e
	TO			2.5	89.5 d	126 c
	20	18.0 e		63.50	0 .	27.2 a
	40	17.5 e	48.2 c	1	11.1 C	
	Mode	15.2 2	70.8 z	66.2 y	83.3 z	126.1 Y

 $^2 M_{\rm cen}$  with similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.

bSeed not separate from pod.



proportional to photoperiod at maturity. In stems, the levels of free sugars at the peak flowering stage were similar under all photoperiod (Table 4), with the 18-hour treatment slightly lower than the 13-hour. At pod filling and at maturity, level was proportional to photoperiod. Under all photoperiods stem free sugar decreased as the pods reached maturity (Table 4) possibly due to redistribution from the stem to the growing pods. Free sugar in leaves was significantly lower under 8-hour than under longer photoperiods at the peak flowering stage (Table 4), possibly because of a light period insufficient for maximal photosynthesis. However, the level increased at the pod filling stages under both the 8 and 13-hour daylengths, but not under the 18-hour photoperiod (Table 4). This may be due to the stimulating effect of longer daylength on sugar movement out of the leaf. Free sugar concentration in seeds was proportional to photoperiod at all sampling dates, with maximum differences at pod filling, when the 18-hour treatment remained high relative to the 8- and 13-hour treatments (Table 4). The concentration decreased in all daylength treatments from peak flowering to pod filling. At maturity, however, the concentration increased slightly under 8- and 13-hour daylengths, while decreasing slightly under the 18-hour treatment. The increase in free sugar levels in seeds under the 8- and 13-hour daylengths was


3.5

;

accompanied by a decrease in stems and roots (Table 4). Free sugar levels in pod wall decreased independently of photoperiod as maturation advanced (Table 4), possibly due to re-allocation to the growing seeds. In pea, the pod exports all of its assimilate to the enclosed seed (2).

#### Starch

In vegetative tissues starch levels did not consistently increase with photoperiod (Table 5), levels in roots being lower under the 18-hour day in both cultivars (Table 6). No cultivar difference in starch level was evident under any of the daylength regimes. However, differences were seen in stems of Black Turtle Soup under the 13- and 18-hour daylengths (Table 6) since it stored more starch in stems than Seafarer. Also, Black Turtle Soup, which showed a greater growth response to longer daylength (13, and see Chapter 3), had a higher level of starch in the stems under the longer daylengths. This response did not occur in Seafarer. The starch level was found significantly higher only in leaves of Black Turtle Soup under the 18-hour day (Table 6). The level of starch in seed and pod wall was not affected by increasing daylengths (Table 6). No cultivar differences for starch were found in these organs under the longer daylengths (13 and 18 hours).




Developmental Stages, Cultivars and Daylength Effects on Starch Level in  $\ensuremath{\mathsf{Various}}$  Plant Parts^a Table 5

				Mg Gl	rcose ber	g D	Mg Glucose per g Dry Weight	
		Root	п	Stem	Leaf	J.	Seed	Pod
1	Time							
	peak flowering	20.4				д	1	56.5 b
	Pod filling Mathrity	16.4	фД	117.4 1 25.3	b 28.7	ಡ	389.0 c 343.3 b	357.8 c 38.4 a
11	Cultivar							
	alle Solla	19.9	E	108.3	n 77.2	q	363 m	142.4 m
	Seafarer	19.1	E	63.6	m 41.3	E	369 ш	159.4 n
TII.	Daylength							
		18.8	×		x 46.3	×	332 x	126 x
	8 nours	23.9	>	79.0	y 46.9		167 y	402 z
	13 nours 18 hours	15.9	×		84.6	×		159 Y
Inter	Interactions							
-		*		*	*		n.s.	*
	IXI	*		*	*		*	*
	IX III	5		*	*		*	*
	II × III	*		*	*		n.s.	n.s.

 $^{\rm a}_{\rm Means}$  followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.

b<sub>Seeds</sub> not separate from pod.




Effects of Photoperiod and Cultivar on Starch Concentration (mg/g dry weight) in Bean Tissues  $^{\rm A}$ 


Table 6

Photoperiod	1000			Tissue		
(hours)	Cultivar	Root	Stem	Leaf	Seed	Pod Wall
ω	Black Turtle Soup Seafarer	17.8 ab 19.7 bc	62.6 a 47.2 a	49.3 a	227 a 250 b	107 a 145 b
	Mean	18.8 x	54.9 x	46.3 x	239 x	126 x
13	Black Turtle Soup Seafarer	24.4 d 23.5 cd	102.8 b 55.3 a	58.5 a 35.3 a	286 cd 287 d	167 c 168 c
	Mean	23.9 Y	Y 0.67	46.9 x	287 z	167 Y
18	Black Turtle Soup Seafarer	17.6 ab 14.2 a	159.5 c 88.3 b	123.9 b 45.4 a	268 bc 258 b	153 bc 165 c
	Mean	15.9 z	123.9 z	84.6 y	263 y	159 Y

 $^{\rm a}_{\rm Means}$  within columns and sets (abc, xyz) with similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.



The pattern of starch accumulation in roots during pod development of the two cultivars showed that at the onset of flowering Black Turtle Soup stored more starch in roots than Seafarer (Table 7). This difference remained until the peak flowering stage. However, when seeds started to develop, the starch level in the roots of Black Turtle Soup dropped to the same level as Seafarer and subsequently there were no cultivar differences (Table 7). Black Turtle Soup stems contained higher starch levels in the flower initiation and pod filling stages than Seafarer: cultivar differences in starch level in the stems disappeared at maturity (Table 7). The high starch level in the stem of Black Turtle Soup may have been due to the indeterminate character and prolonged flowering behavior of this cultivar. This ability to continue vegetative growth after flower initiation can also result in higher starch level in leaves of Black Turtle Soup (Table 7). No differences were evident between cultivars for the starch level in seeds during pod development (Table 7). The starch level in immature pods of both cultivars was lower at the peak flowering stage. At the pod filling stage when seeds could be separated from the pod wall, starch levels in green pod wall of both cultivars rose considerably (Table 7). At maturity the seeds still contained a high




Effects of Cultivar and Time of Sampling on Starch Concentration (mg glucose/g dry weight) in Bean Tissues  $\!\!\!^{\rm a}$ Table 7

Time After	+			Tissue		
ation (days)	Cultival	Root	Stem	Leaf	Seed	Pod Wall
0	Black Turtle Soup Seafarer	21.2 cde 7.1 a		62.4 b 37.1 a 51.1 b 33.3 a	1 1	11
	Mean	14.1 p	56.7 q	56.7 q 35.2 p		
10	Black Turtle Soup Seafarer	23.0 e 17.9 bcd	142.9 d 87.5 c	142.9 d 121.4 c 87.5 c 58.3 b	1.1	54.9 b <sup>b</sup> 58.2 b
	Mean	20.4 q		115.2 r 89.9 q	ı	56.6 q
20	Black Turtle Soup Seafarer	15.2 b 17.5 bc	156.7 d 78.0 c	33.0 a 24.3 a	379.8 b 398.3 c	338.0 c 377.5 d
	Mean	16.4 p	117.4 r	28.7 p	389.0 q	357.8 r
40	Black Turtle Soup Seafarer	21.6 cde 22.0 de	25.4 a 25.2 a	1 1	347.0 a 339.5 a	34.3 a 42.5 ab
	Mean	21.8 q	25.3 p	1	343.3 p	38.4 p


<sup>a</sup>Means within columns and sets (abc, pgr) with similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.

<sup>b</sup>Seed not separate from pod.



starch content (Table 7), but starch in the matured, yellow pod wall dropped to a very low level (Table 7).

Under the 8- and 13-hour daylength, the starch level in roots followed a similar pattern for both cultivar (Table 8). It was high at peak flowering, dropped markedly at the pod filling stage where carbohydrate reserve was in great demand for embryo development, and increased again at maturity where the final size of various organs had been achieved (Table 8). Under the 18-hour daylength the plants retained more flowers and pods (see Chapter 3, Table 1), thus more carbohydrates may have been available to support these reproductive organs and less was left for the root. This trend was especially evident at peak flowering (Table 8) where the starch level in root decreased. From peak flowering to pod filling where more than one-half of pod abscission had occurred, the plants had adjusted the number of pods to be retained. This may have caused a rise in starch level in roots at the pod filling stage under the 18-hour daylength. The interaction between daylength and stage of pod development on starch level in stems was shown in Table 8. The longer the daylength, the more starch was found in the stems at peak flowering and pod filling (10 and 20 days after flower initiation). At maturity, no significant differences were found for starch levels in stems between the 13- and 18-hour daylengths, but a significantly




Effects of Photoperiod and Time of Sampling on Starch Concentration (mg glucose/g dry weight) of Bean Tissues  $^{\rm A}$ 

Photoperiod	Time After	Ď		Tissue		
(hour)	riower initiatation (days)	Root	Stem	Leaf	Seed	Pod Wall
α	0	14.2 ab	56.7 b	35.2 b	1	ı
•	1.0		83.1 bc	75.5 c	1	52.4 b
	20	12.5 a	69.3 b	17.0 a	367 b	309.2 c
	40		12.4 a	1	297 a	16.4 a
	Mean	17.6 x	55.4 x	42.6 x	332 x	126 x
	c	14.2 ab	56.7		•	1
13	0 -	28.1 d	95.5	47.3 b	1	56.3 b
	200	17.5 b		46.6 b	442 c	418.1 e
	40	26.2 d	33.5	1	361 b	28.2 a
	Mean	21.5 Y	73.4 ⅓	43.0 x	401 z	167 y
	c		56.7	35.2 b	1	,
18	5			146.8 d	1	61.0 b
	20 0	19.0 p	Н	22.4 ab	357 b	346.0 d
	40			1	371 b	70.5 b
	Mean	15.5 x	107.1 z	68.1 Y	364 Y	159 Y

 $^{\rm a}_{\rm Means}$  within columns and sets (abc, xyz) with similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.


bseed not separate from pod.



lower starch level was found under the 8-hour regime (Table 8). The starch level in leaves reached the maximum at peak flowering and then declined rapidly. This was seen in both 8- and 18-hour daylength treatment (Table 8), while in the 13-hour day the starch level did not drop at the pod filling stage. No explanation for this is apparent. The starch level in seed and pod wall reached a maximum at pod filling and declined at maturity (Table 8). The rate of decline was more rapid in the pod wall than in the seed (Table 8) possibly due to the export of starch from pod wall to seed as was shown in field pea (2). This pattern was seen in all three daylength regimes, except no decrease was seen in starch level in the seed at maturity in the 18-hour day (Table 8). Perhaps the great decrease in starch in stems from pod filling to maturity may be related to the changes in the level of starch in seeds under this 18-hour day (Table 8).

#### Relationship of Pod Abscission to Levels of Available Carbohydrates

Increasing the daylength during the reproductive phase increased plant weight, seed yield, and pod retention in these two dry bean cultivars. In this study increasing the daylength after flower initiation also caused an increase in free sugar and starch levels in roots, stems and leaves. A relationship may exist between available carbohydrate in the vegetative organs and percentage of



pod abscission as was suggested in the nutritional hypothesis first put forward by Mason (9). In this study no precise relationship could be established between percentage of pod abscission and available carbohydrate levels in the vegetative organs of these two cultivars (Table 9). However there was a general trend of negative correlation between percentage of pod abscission and available carbohydrates in most organs. This indicated that the higher the carbohydrate levels, the lower the abscission rate. The data presented here, though consistent with Mason's explanation, do not prove the level of available carbohydrates to be the cause of pod abscission in beans. Since abscission is a complex process, other factors than carbohydrate levels, such as nitrogenous levels or even endogenous hormonal levels, may play a part in determining pod abscission. Future studies toward establishing the balance of these factors in relation to pod abscission are needed.


Table 9

Correlation Coefficients Between Percentage of Pod Abscission and Available Carbohydrate in the Vegetative Tissues of the Two Drybean (Phaseolus Vulgaris L.) Cultivars Grown Under 3 Daylengths

		Bla	Black Turtle Soup	dnos		Seafarer	
	Characters	8 Hours	1 1	13 Hours 18 Hours	8 Hours	13 Hours	18 Hours
i.	At pod filling stage						
	Free sugar in root	.05	41	.11	48 70a	.31	.41
	Free sugar in stem	26	39	115	60	.12	333
	Starch in root	.43	-22 -75b	.15	98 - 88p	10	25
	Starch in stem Starch in leaf	01	57	.45	.14	67b	.07
II.	At maturity						
	Free sugar in root	.10 .63b	21 61a	12	1.27	23 66ª	.32
	Starch in root Starch in stem	01	57	38	- 35	01	29


asignificant at 5% level.

bsignificant at 1% level.




#### REFERENCES

- Eaton, F. M. and D. R. Ergle. 1953. Relationship of seasonal trends in carbohydrate and nitrogen levels and effects of girdling and spraying with sucrose and urea to the nutritional interpretation of boll shedding in cotton. Plant Physiol. 28: 503-520.
- Flinn, A. M. and J. S. Pate. 1970. A quantitative study of carbon transfer from pod and subtending leaf to the ripening seeds of the field pea (Pisum arvense L.) J. Expt. Bot. 21: 71-82.
- Guinn, G. 1974. Abscission of cotton floral buds and bolls as influenced by factors affecting photosynthesis and respiration. Crop Sci. 14: 291-293.
- Greenwood, E. A. N., P. Farrington and J. D. Beresford. 1975. Characteristics of the canopy, root system and grain yield of a crop of <u>Lupinus angustifolius</u> cv. Unicrop. Aust J. Agric. Res. 26: 497-510.
- Hawkins, R. S., R. L. Matlock and C. Hobart. 1933.
   Physiological factors affecting the fruiting of cotton with special reference to boll shedding.
   Arizona, Agri. Exp. Sta. Tech. Bull. 46: 361-407.
- Hicks, D. R. and J. W. Pendleton. 1969. Effect of floral bud removal on performance of soybeans. Crop Sci. 9: 435-437.
- Laidlaw, R. A. and S. G. Reid. 1952. Analytical studies on the carbohydrates of grassland clovers. I. Development of methods for the estimation of free sugar and fructosan contents. J. Sci. Fd. Agric. 3: 19-25.
- Loomis, W. E. 1926. A study of the clearing of alcoholic plant extracts. Plant Physiol. 1: 179-189.



- Mason, T. G. 1922. Growth and abscission in Sea Island Cotton. Ann. Bot. 36: 457-483.
- Ojehomon, O. O. 1972. Fruit abscission in cowpea, Vigna unguicuiata L. Walp. J. Expt. Bot. 23: 751-761.
- Smith, D. 1969. Removing and analyzing total nonstructural carbohydrates from plant tissue. Research Report #41. University of Wisconsin, 12 pp.
- Smith, D. and L. F. Graber. 1948. The influence of top growth removal on the root and vegetative development of biennial sweet clover. J. Amer. Soc. Agron. 40: 818-831.
- Subhadrabandhu, S., M. W. Adams and D. Penner. 1976.
   Abscission of reproductive structures in dry bean. II. Effects of daylengths on growth and pod abscission in Phaseolus vulgaris L. Agronomy Abstracts, p. 77.
- Van Schaik, P. H. and A. H. Probst. 1958. Effects of some environmental factors on flower production and reproductive efficiency in soybeans. Agron. J. 50: 192-197.
- Van Steveninck, R. F. M. 1957. Factors affecting the abscission of reproductive organs in yellow lupins (Lupinus Luteus L.) I. The effect of different patterns of flower removal. J. Expt. Bot. 8: 373-381.
- Wadleigh, C. H. 1944. Growth status of cotton plant as influenced by the supply of nitrogen. Arkansas Agr. Exp. Sta. Bull. 446: 1-138.
- Weinmann, H. 1947. Determination of total available carbohydrates in plants. Plant Physiol. 22: 279-290.
- 18. Wien, H. C., S. L. Altschuler, J. L. Ozbun and D. H. Wallace. 1976. 14C-assimilate distribution in Phaseolus vulgaris L. during the reproductive period. J. Amer. Soc. Hort. Sci. 101: 510-513.
- Yemm, E. W. and A. J. Willis. 1954. Stomatal movements and changes of carbohydrate in leaves of Chrysanthemum maximum. New Phytol. 53: 373-396.



#### CHAPTER 5


ABSCISSION OF FLOWERS AND FRUITS IN <u>PHASEOLUS</u> <u>VULGARIS</u> L.

IV. GROWTH AND POD ABSCISSION AS AFFECTED BY

CARBON DIOXIDE LEVELS

# Abstract

Abscission of young pods was studied in the green-house in two Phaseolus vulgaris L. cultivars over two seasons. Plants were grown under 300 (ambient air) and 800 ppm carbon dioxide after their flower buds became visible. High carbon dioxide level reduced pod abscission, increased plant size and seed yield. The relationship between pod abscission and growth is discussed.




## Introduction

Since the report of Wittwer (5) on increasing crop yield by carbon dioxide enrichment, numerous reports have confirmed his observations. Yields of field-grown soybean have been increased nearly 30% by carbon dioxide enrichment (1,200 ppm) (4), as a result of both an increase in number of pods filled and a decrease in pod abortion. In cotton an increase in carbon dioxide concentration in the atmosphere decreased abscission of floral buds and bolls (3). In barley (2), rice (6), and soybean (4) increasing the carbon dioxide concentration during the reproductive phase increased yield.

Growing dry bean plants under long photoperiods after flowering increased yield (see Chapter 3). This was associated with less pod abscission and larger plants. The study reported here was designed to determine the effects of increasing carbon dioxide levels during the pod development period on growth and pod abscission of dry beans.

## Materials and Methods

Two dry bean cultivars, "Seafarer" and "Black
Turtle Soup," were grown in the greenhouse at East
Lansing in the spring of 1975 and 1976 (for procedures, see
Chapter 3). When the plants reached the flower initiation stage, 30 of each cultivar were placed in each
of two clear polyethylene chambers (7.2 by 3 by 3 meters).



In each chamber air drawn in from one end was released at the other end. Carbon dioxide was fed into the mixing chamber of carbon dioxide and air. The gas mixture was introduced into one chamber with a fan at the rate of 148 cubic meters per hour. The concentration at the plant level was maintained in the range of 750 to 850 ppm, as maintained with an infra-red gas analyzer. This was continued for 40 days.

Ambient air was blown into the second chamber at the same rate. The carbon dioxide concentration was in the range of 300 to 350 ppm. Three harvests were made, pod abscission and growth were measured as before (Chapter 3).

## Results and Discussion

## Pod Abscission

Plant subjected to a high carbon dioxide level produced more flowers and retained more pods than the control (Table 1). Response was similar in both cultivars (Table 2). Data for both 1975 and 1976 showed the same trends (Table 1). Pod abscission was reduced more by carbon dioxide level in Black Turtle Soup than in Seafarer (Table 2). This was associated with larger plant (Table 3), more leaves, greater leaf area and leaf weight per plant in Black Turtle Soup (Table 4).

Pod abscission was not recorded at peak flowering (10 days after flower initiation). At pod filling stage




Table 1

Stages of Pod Development and Carbon Dioxide Levels Effect on Number of Flowers, Number of Pods and Percentage of Abscission in Two Dry Bean Cultivars Grown Under the Two Seasons\*

		No. F	No. Flowers	No. Pods	Pods	% Abscission	ission
		1975	1976	1975	1976	1975	1976
i.	Developmental Stages						
	Peak flowering	16.8 a	12.3 a		10.9 b	1	. 1
	Pod filling Maturity	15.8 a 15.7 a	12.9 a 15.5 b	11.5 b 8.8 a	8.2 a 7.3 a	25.1 a 41.1 b	28.4 a 47.3 b
II.	Cultivar				-		
	Black Turtle Soup Seafarer	10.5 m	8.3	8.8 E 6	6.5 m	21.9 m	20.5 m
III.	CO <sub>2</sub> Level				:	;	
	300 ppm 800 ppm	14.6 x 17.6 y	10.4 x 16.7 y	9.3 x 13.8 y	6.0 × 11.5 y	44.2 y 22.0 x	46.5 y 22.2 x
Inte	Interactions			•			
	Ix I	*	s d	*	* *	n.s.	n.s.
	IX III	n.s.	n.s.	*	n.s.	n.s.	n.s.
	11 X 11I	n.s.	n.s.	n.s.	n.s.	n.s.	*
	III x II x I	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.

 $^{\rm a}{\rm Me}_{\rm ans}$  within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test,





Table 2

Effects of Carbon Dioxide Levels on Number of Flowers, Number of Pods Retained, and Percentage of Pod Abscission in Two Dry Bean Cultivars Grown under the Two Seasons (data are means at all harvesting stages)

THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER.							The second secon
Carbon Dioxide	+	No. Flo	No. Flowers/Plant No. Pods/Plant % Pod Abscission	No. Pods	s/Plant	% Pod Ab	scission
Levels (ppm)	כמוכוימו	1975	1975 1976	1975	1976	1975 1976 1975 1976	1976
300	Black Turtle Soup 9.2 a 5.4 p	9.2 a	5.4 p	5.5 a	4.1 p	5.5 a 4.1 p 34.48 b 26.09 q	26.09 q
	Seafarer	20.0 €	20.0 c 15.3 r	8.1 b	7.9 g	8.1 b 7.9 q 53.88 c 66.88 s	88.99
800	Black Turtle Soup 11.9 b 11.3 g	11.9 b	11.3 q	11.4 c	9.0 g	11.4 c 9.0 g 9.30 a 14.88 P	14.88 p
	Seafarer	23.3 d	23.3 d 22.0 s	13.7 d	14.1 r	13.7 d 14.1 r 34.75 b 43.57 r	43.57 r

 $^{\rm a}_{\rm Means}$  within a given parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.



H

Table 3

Effects of Carbon Dioxide Levels on Dry Weight Per Plant of Root, Stem and Total Plant Weight in Two Dry Bean Cultivars Grown Under the Two Seasons (data are means at all harvesting stages)<sup>a</sup>

ant Wt.	1976	7.0 a 5.9 q	6.8 a 3.9 p	13.4 c 10.1 s	11.6 b 7.6 r
Total Plant Wt. (g)	1975 1976	7.0 a	6.8 a	13.4 c	11.6 b
Wt./Plant (g)	1975 1976	1.3 a 1.0 p	1.4 a 0.8 p	2.3 c 2.2 r	1.9 b 1.9 q
Stem Wt	1975	1.3 a	1.4 a	2.3 0	1.9 b
Root Wt./Plant Stem Wt./Plant (g) (g)	1975 1976	1.0 a 1.2 q	1.0 a 0.7 p	1.6 c 2.1 s	1.4 b 1.4 r
Root Wt./	1975	1.0 a	1.0 a	1.6 c	1.4 b
7 c77 i + [,1]		Black Turtle Soup	Seafarer	Black Turtle Soup	Seafarer
Carbon Dioxide	(1.24)	300		800	

 $^{\rm a}_{\rm Means}$  within a given parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.





Table 4

Effects of Carbon Dioxide Levels on Leaf Area Per Plant, Number of Leaf Per Plant and Leaf Weight Per Plant in Two Dry Bean Cultivars Grown Under the Two Seasons (data are means at all harvesting stages)

Carbon Dioxide	Cultivar	Leaf A	rea/Plant m <sup>2</sup> )	<pre>Leaf Area/Plant Number Trifoliate Leaf Weight/ (cm2)</pre> Leaf/Plant (9)	ifoliate lant	Leaf V Plar	eaf Weight/ Plant (g)	
mevers (ppm)		1975	1975 1976	1975 1976	9261	1975	1975 1976	
300	Black Turtle Soup	668 a	420 p	11 a 11 p	11 p	2.4 b	2.4 b 2.5 q	
	Seafarer	659 a	342 p	13 a 10 p	d 01	2.0 a	2.0 a 1.6 p	
800	Black Turtle Soup	930 b	930 b 1043 r	19 b 14 r	L4 r	5.0 d	5.0 d 6.3 s	
	Seafarer	739 a	739 a 723 g	13 a 12 g	L2 q	3.4 c	3.4 c 3.2 r	

 $^{\rm a}{\rm Means}$  within a given parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.



and at maturation, high carbon dioxide reduced pod abscission in both cultivars (Table 1, Appendix C tables C-1 and C-2). About half of the pod abscised at pod filling stage in Black Turtle Soup under the control treatment, but only one-third were lost at this same stage when this cultivar was grown under high carbon dioxide (Table C-1). However, for Seafarer, a cultivar with a high abscission level, about two-thirds of pod loss occurred at the pod filling stage when they were grown under ambient carbon dioxide. This ratio was decreased to one-half when high carbon dioxide level was supplied to the plants (Table C-1). Similar results were obtained in 1976 (Table C-2). Two interpretations are offered for these results. First, high competition for carbohydrates occurred during the early stages of pod development, and this competition will determine the amount of pods to be retained. This was supported by the high portion of pods abscised at pod filling in relation to that at maturity when plants were grown under a low carbon dioxide level. The second explanation was shown in the drop of this ratio of pod abscision at pod filling and at maturity with carbon dioxide enrichment. This indicated that more carbohydrate was made available under a high carbon dioxide regime for feeding young pods at this stage. Percentage of pod abscission obtained by carbon dioxide enrichment was similar to that obtained by increasing daylength, as reported earlier (Chapter 3).

## Leaf Development

With high carbon dioxide levels more leaves per plant were obtained. This resulted in greater leaf area and higher leaf weight produced by the plant (Table 5). With a low carbon dioxide level there was no significant difference in number of leaves and leaf area per plant of the two cultivars, but when the plants were subjected to the higher carbon dioxide level cultivar differences in number of leaves and leaf area were obtained. Black Turtle Soup exhibited greater response to increasing carbon dioxide than Seafarer (Table 4). This response occurred in both the 1975 and 1976 experiments (Table 4). This high response to increasing carbon dioxide level in Black Turtle Soup can be related to its growth habit. Black Turtle Soup is a semi-vine bean and the plant can still make some vegetative growth after flowering. Therefore, when more carbohydrate was synthesized under with a high carbon dioxide level, vegetative growth may also be promoted and this resulted in more leaves being produced. For Seafarer, a determinate type, vegetative growth does not continue much after the plant reaches the reproductive phase, therefore, more carbohydrate produced by carbon dioxide enrichment did not result in more leaves being produced (Table 4). However, the high carbon dioxide treatment did result in lower percentage of pod abscission. A negative correlation between percentage





Table 5

Stages of Pod Development and Carbon Dioxide Levels Effect
on Number of Leaf and Leaf Area per Plant in Two Dry
Bean Cultivars Grown Under the Two Seasons<sup>a</sup>

		No 1	Leaf	Leaf Plant	Area/ (cm <sup>2</sup> )
		1975	1976	1975	1976
ı.	Developmental Stage	s			
	Peak flowering Pod filling Maturity	15.8 b 12.6 a	10.7 a 12.9 b	742 a 756 a	549 a 716 b
II.	Cultivar				
	Black Turtle Soup Seafarer	15.2 m 13.3 m	12.3 n 11.3 m		732 n 533 m
III.	CO <sub>2</sub> Level				
	300 ppm 800 ppm	12.2 x 16.2 y			381 x 883 y
Inter	actions				
	I x III I x III	* n.s. **	** ** *	n.s. **	n.s. **
	I x II x III	*	n.s.	n.s.	n.s.

 $<sup>^{\</sup>rm a}{\rm Means}$  within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.



of pod abscission and leaf area per plant was also noticed in both cultivars under both levels of carbon dioxide (Table 6). However, at 300 ppm carbon dioxide, at which leaf area may be more important in supplying carbohydrate for pod retention, a highly significant correlation occurred. This significant correlation was seen in the indeterminate cultivar, Black Turtle Soup. In Seafarer, the determinate type bean, no significant correlation between leaf area and percentage of pod abscission was found. At the high carbon dioxide level (800 ppm), more ideal for photosynthesis, leaf area per se may not be as critical for supporting the young pods, therefore, no significant correlation between leaf area and pod abscission was found (Table 6).

The number of leaves per plant was significantly lower at the pod filling stage as compared to peak flowering (Table 5). This was due to the loss of some old leaves at pod filling. However, leaf area and leaf weight per plant were not significantly different at these two stages. This indicated that very little or no leaf growth occurred after peak flowering in 1975. This was in contrast to the 1976 data where there were indications of leaf growth after this stage (Table 5).

# Plant Development

Greater growth and larger plants were observed with the carbon dioxide enrichment treatment. Increase

enter de la companya de la companya

4


Table 6

Correlation Coefficients Between Percentage of Pod Abscission and Other Growth Characters in Two Dry Bean Cultivars Subjected to Different Carbon Dioxide Levels After Flowering (data taken at pod filling stage)

	Black Tu	Black Turtle Soup	Seaf	Seafarer
Characters	300 ppm CO <sub>2</sub>	800 ppm CO <sub>2</sub>	300 ppm CO <sub>2</sub>	800 ppm CO <sub>2</sub>
Root weight	.62ª	33	24	.42
Stem weight	03	68ª	23	11
rosf weight	-,80 <sup>b</sup>	36	32	80.
rest norgh-	39	02	60.	.03
Leaf area	76 <sup>b</sup>	47	22	14

asignificant at 5% level.

bsignificant at 1% level.



in root, stem, leaf and total plant weight occurred in both cultivars (Table 3) in both years of study. The increase in vegetative growth with the high carbon dioxide level in both cultivars was associated with a higher number of pods retained (Table 2).

Black Turtle Soup produced more dry weight in the vegetative organs than the Seafarer (Table 3). The greater growth response with high carbon dioxide enrichment was also evident by reduced pod abscission in Black Turtle Soup. Whether the higher vegetative weight determined pod abscission was not known, but the data show a relation between these characters (Table 6).

Total plant weight was greater with the carbon dioxide enriched treatment after flowering, while under the ambient carbon dioxide level this weight was decreased (Table 2). This may be due to the loss of senescing leaves at maturation. The slightly higher total plant weight at maturity under high carbon dioxide enrichment was due to the increased weight of reproductive organs (seeds and pods). This was also seen in decreased weight of root and stem (Table C-4) from peak flowering to maturation. This loss in vegetative weight compensated for an increase in reproductive weight. This dynamic situation in weight allocation affected the yield of a plant. Competitive ability of various organs at any developmental stage of plants grown under a particular environment would undoubtedly influence pod abscission.

· . 11:0

### Seed Yield

Plants growing under the high carbon dioxide level (800 ppm) produced more mature seeds. This was seen in both cultivars (Table 7). The more seed produced, the greater the seed weight per plant. The high number of seed produced under the high carbon dioxide level resulted in slightly smaller seeds produced in 1975 (Table 7). This may have been due to the competition between the more numerous seeds in the pods. However, this seed size difference was not observed in 1976 (Table 7).

Cultivar differences in seed yield existed in response to increasing carbon dioxide concentration.

Under the high carbon dioxide level, Black Turtle Soup produced more seed and had higher yield than Seafarer in both years (Table 7). This greater response by Black Turtle Soup was also seen in vegetative growth and pod abscission.

The correlation between percentage of pod abscission and other reproductive structures at pod filling was given in Table 8. At the ambient carbon dioxide level, the number of flowers produced in both cultivars was positively correlated to percentage of pod abscission. This suggested that under this carbon dioxide level at the pod filling stage, competition for carbohydrate occurred between the flowers and young pods. The more reproductive organs, the more competitive force and the




Table 7

Effects of Carbon Dioxide Levels on Number of Seeds Per Plant, Seed Weight Per Plant and Weight of One Seed in Two Dry Bean Cultivars Grown Under Two Seasons (data are means at all harvesting stages)<sup>9</sup>

Carbon Dioxide Levels (ppm)	Cultivar	No. See	No. Seeds/Plant	Seed Wt./Plant (g)	lant	Weight/Seed (9)	eq
		1975	1975 1976	1975 1976	92	1975 1	1976
300	Black Turtle Soup	23 a	23 a 22 p	4.1 b 4.0 p	d 0	0.17 b 0.16 p	16 p
	Seafarer	16 a	16 a 21 p	2.5 a 3.1 p	J p	0.15 ab 0.15 p	15 p
800	Black Turtle Soup	53 c	53 c 54 r	8.1 c 7.1 q	1 4	0.13 a 0.13 p	13 p
	Seafarer	39 b	39 b 37 q	6.9 c 5.7 q	J 4	0.13 a 0.18 q	18 g

 $^3 M_{\rm beans}$  within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.





Table 8

Correlation Coefficients Between Percentage of Pod Abscission and Other Reproductive Characters in Two Dry Bean Cultivars Subjected to Different Carbon Dioxide Levels After Flowering

	400	Black Tur	Black Turtle Soup	Seafarer	arer
	Characters	300 ppm CO <sub>2</sub>	800 ppm CO <sub>2</sub>	300 ppm CO <sub>2</sub>	800 ppm CO <sub>2</sub>
i.	At Pod Filling Stage				
	Flowering period	.30 .78b	12	18 .63a	.36 .79ª
	Number of Pods	. 58	57	37	46
	Seed weight	458.	.52	. 42	. 78 <sup>b</sup>
II.	At Maturity				
	Flowering period	.59ª	55	.33	25
	Number of flowers	g88.	.11	.44	90°-
	Number of pods	.46	41	- 86~	2.5
	Number of seeds	.70ª	57	47	2.4
	Seed weight	927.	36	37	T#


<sup>a</sup>Significant at 5% level.

bsignificant at 1% level.




more they would abscise. However, when the plants were grown under the high carbon dioxide level, a negative correlation was seen in Black Turtle Soup while the positive correlation was still observed in Seafarer (Table 8). The negative correlation in Black Turtle Soup may have resulted from a greater vegetative growth response by this cultivar to the higher carbon dioxide level. Therefore, flowers and young pods had a greater share of the assimilate produced, thus fewer pods would have to sacrifice themselves by abscissing. However, for Seafarer which is a determinate plant, vegetative growth did not occur after flowering, the plants exhibited less growth response to the increased carbon dioxide level. Also this cultivar produced many flowers in a short flowering time as compared to Black Turtle Soup (see Chapter 2). Thus the demand for food from these flowers and young pods was made in a short time period; even if plants had grown larger under the high carbon dioxide, the increase in growth may not have been sufficient to meet the demand from those flowers and pods. Therefore, a high percentage of pod abscission still occurred as more flowers were produced.

Negative correlation between percentage of pod abscission and yield (seed weight) at maturity indicated that as percentage of abscission went up lower yield resulted. This was seen in Black Turtle Soup grown at



300 ppm carbon dioxide and in both cultivars at the 800 ppm carbon dioxide level (Table 8). Although pod abscission can affect seed vield, in dry beans there are many components affecting yield (1). Thus, nonsignificant negative correlations under these conditions may be explained. It is possible that percentage of pod abscission might be significantly correlated to any one of the vield components instead of total vield in this case. The significantly positive correlation of 0.77 between percentage of pod abscission and seed vield in Black Turtle Soup grown under ambient air (300 ppm carbon dioxide) may be explained as follows. The limited photosynthesis may have been insufficient for retaining many pods. Thus the competition among pods for growth could result in a condition that the more pods abscised the greater the chance for the remaining pods to grow.

The previous study of cultivar response to increasing daylength showed positive response to daylength in both cultivars, but Black Turtle Soup seemed to exhibit a greater response than Seafarer (Chapter 3). A similar result was also found when these two cultivars were grown under high carbon dioxide concentration. These two experiments pointed out two things. First, Black Turtle Soup had greater response when grown under conditions favoring photosynthesis after flowering. The second was that since both extending daylength and




carbon dioxide enrichment gave similar results in pod abscission, seed yield, and growth rate, there must be similar physiological responses of plants that result from these two environmental factors. These physiological responses may be responsible for determining the abscission capacity of the plants. If that physiological factor is not carbohydrate availability, then it must be some factors influenced by or closely associated with the levels of carbohydrate.

codir:

1. 10.

#### REFERENCES

- Adams, M. W. 1967. Basic of yield component compensation in crop plants with special reference to the field bean. Phaseolus vulgaris L. Crop Sci. 7: 505-510.
- Gifford, R. M., P. M. Bremner and D. B. Jones. 1973. Assessing photosynthetic limitation to grain yield in a field crop. Aust. J. Agric. Res. 24: 297-307.
- Guinn, G. 1974. Abscission of cotton floral buds and bolls as influenced by factors affecting photosynthesis and respiration. Crop Sci. 14: 291-293.
- Hardman, L. L. and W. A. Brun. 1971. Effect of atmospheric carbon dioxide enrichment at different developmental stages on growth and yield components of soybeans. Crop Sci. 11: 886-888.
- Wittwer, S. H. and W. M. Robb. 1964. Carbon dioxide enrichment of greenhouse atmospheres for food grop production. Econ. Bot. 18: 34-56.
- Yoshida, S., J. H. Cock and F. T. Parao. 1972.
   Physiological aspects of high yields. In "Rice Breeding" IRRI, Los Banos, pp. 455-468.




#### CHAPTER 6

ABSCISSION OF FLOWERS AND FRUITS IN PHASEOLUS VULGARIS L.

V. RELATIONSHIP OF POD ABSCISSION TO DIFFERENTIAL LEVELS
OF CARBOHYDRATES PRODUCED BY VARIATION IN CARBON
DIOXIDE CONCENTRATION

## Abstract

Levels of free sugar, starch and total nonstructural carbohydrates were determined in two dry bean
(Phaseolus vulgaris L.) cultivars grown under ambient
(A300 ppm) and enriched (A800 ppm) level of carbon
dioxide. Free sugars were higher in both vegetative (root,
stem, and leaf) and reproductive (seed and pod wall) tissues in both cultivars grown under high carbon dioxide,
while starch concentration was higher only in the vegetative tissues. Black Turtle Soup (indeterminate) contained
significantly more starch in the vegetative tissues, but
not in the seed, than did Seafarer (determinate). Free
sugar and starch decreased at maturity in the vegetative
tissues, but not in seed, where most of the carbohydrate
was stored as starch. A shift of available carbohydrates
from the vegetative tissues and the pod wall to the seed



occurred with maturation. The role of available carbohydrates in pod abscission, seed yield, and vegetative growth is discussed.

er sates esable

#### Introduction

An extensive review article by Wittwer and Robb (5), emphasizing tremendous increases in crop production obtained by carbon dioxide enrichment of greenhouse atmospheres, aroused the attention of crop physiologists in the early sixties. Carbon dioxide enrichment has since been reported to increase yield in soybean (1, 3), wheat (4) and cotton (2).

Our previous studies with <u>Phaseolus vulgaris</u> L. showed that growth rate, seed yield, and pod retention could be increased by raising the ambient carbon dioxide concentration (see Chapter 5). The objective of the present study was to determine the effects of such treatment on carbohydrate concentration of various tissues.

## Materials and Methods

# Sample Preparation

Two dry bean cultivars, "Black Turtle Soup" (indeterminate) and "Seafarer" (determinate), were grown and treated with two carbon dioxide levels as previously described (see Chapter 5).

Harvesting times, handling of samples during harvesting, preparation of samples, and analysis of free sugars and total nonstructural carbohydrate has been described elsewhere (Chapter 4). Free sugar, starch, and total nonstructural carbohydrate concentrations are expressed as mg glucose equivalent per g of dry tissues.

(pla

Free

free cult free howe

> the Thes Seaf

and Soup

at p was acti

the carr

hypo Pod

. . . (=10

The data were analyzed factorially, using 10 replications (plants) per treatment.

### Results and Discussion

## Free Sugar

Increasing the carbon dioxide level increased the free sugar concentration in all tissues analyzed for both cultivars (Table 1). Black Turtle Soup retained more free sugars in the root, stem, and leaf than did Seafarer; however, the reverse occurred in seeds and pods, although the difference was not significant (1%) in pods (Table 1). These data indicate a stronger sink effect of seeds in Seafarer than in Black Turtle Soup.

No interaction was observed between cultivar and carbon dioxide level except in the leaf. Black Turtle Soup leaves contained more sugar at 300 ppm carbon dioxide, but no difference was evident at 800 ppm (data not shown).

The free sugar levels in all tissues were high at peak flowering and decreased as the pods developed, as was seen in a previous study (Chapter 4). Several interactions were noted between carbon dioxide level and growth stage. As the pods matured, the free sugar level in the stem and leaves decreased at a faster rate at the low carbon dioxide level than at the high carbon dioxide (Fig. 1A and 1B). These results are consistent with the hypothesis that available free sugars are limiting during pod formation and filling. The free sugar level in the


Table 1

The Effects of Developmental Stage and Carbon Dioxide Levels on Free Sugar Concentration in Various Plant Parts of the Two Bean Cultivars<sup>a</sup>

			Mg Gluc	Mg Glucose per g Dry Weight	/ Weight	
		Root	Stem	Leaf	Seed	Pod
i.	Time					
	Peak flowering	23.9 b		69.7 a	1	267.7 cb
	Pod filling	18.9 a	56.2 b	73.2 a	90.5 b	117.0 b
	Maturity	20.9 a	38.8 a	1	80.6 a	37.7 a
II.	Cultivars					
	Black Turtle Soup	24.2 n	62.3 n	76.4 n	84 m	137 m
	Seafarer	18.2 m	44.8 m	m 9.99	87 n	145 m
III.	CO2Level					
	300 maa 008	18.9 x	42.6 x	58.9 x	x 89	125 x
	mdd 008	23.5 Y	64.5 Y	84.1 Y	102 y	157 Y
Inter	Interactions					
	11 * 1	n.s.	n.s.	n.s.	n.s.	n.s.
	1111	n.s.	*	*	**	*
	1 A 111	n.s.	n.s.	*	n.s.	n.s.
	IX II X III	n.s.	*	n.s.	*	*

 $^{\rm d}_{\rm Means}$  within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.

b<sub>Seed</sub> not separate from pod.






Fig. 1. Levels of starch in A. root; and B. stem; of two Phaseolus vulgaris L. cultivars during their reproductive phase, averaged from the two carbon dioxide levels.

12.11

leaves of plants exposed to 800 ppm carbon dioxide increased between peak flowering and pod filling (Table 2).

The level of free sugar in flowers and newly set pods (seed could not be separated at this stage) was high under both the high and low carbon dioxide treatment at peak flowering but decreased sharply with maturation (Table 2). High carbon dioxide significantly increased seed free sugar concentration at all developmental stages. Free sugars in seeds dropped between pod filling and maturity under the high carbon dioxide treatment, but increased slightly during the same period at the ambient carbon dioxide level (Table 2). Greater pod retention in the high carbon dioxide treatment (Chapter 5) probably created greater competition for sugar among seeds, resulting in a decline in free sugar level from pod filling to maturity. Much pod abscission occurred under the ambient carbon dioxide level and fewer pods and seeds were retained (Chapter 5). This reduced competition and prevented a decline in concentration in the seeds. Free sugar content of pod walls decreased with maturation under both carbon dioxide levels (Table 2).

## Starch

High carbon dioxide increased starch concentrations in vegetative tissues (root, stem and leaf), but decreased them in the seed and pod wall (Table 3). Starch





Table 2

Effects of Carbon Dioxide and Time of Sampling on Free Sugar Concentration (mg glucose/g dry weight) of Bean Tissues<sup>A</sup>

	"	1	2		) c				ր 2	
	Pod			243	104.7 26.1			291.6	129.	49
	g				м 0 Ф				υ 0 Ι	2 2
	Seed		1	1 9	72.6		1	1 2	D C 077	88
Tissues	Leaf	, , ,	04.1.0 64.6.1.0	53.3 3	, i	82 1 3	74.97	0 0 0	1	1
	Stem	70 07	0.65	49.8 b	18.7 a	79.5 6	72.0 d	62.7 c	0 000	
	Root	70 % PK	22.7 bc	16.4 a	17.7 a	20.3 ab	25.1 c	21.4 b	24.1 bc	
Time After Flower Initiation	(days)	0	10	20	40	0	10	20	40	
Carbon Dioxide	revel (ppm)	300				800				

 $^{\rm a}{\rm Means}$  within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.

bSeed not separate from pod.




Effects of Developmental Stages and Carbon Dioxide Levels on Starch Concentration in Various Plant Parts of the Two Dry Bean Cultivars  $^{\rm A}$ Table 3

		2	lg Glucose Eq	nivalent per	Mg Glucose Equivalent per g Dry Weight	
		Root	Stem	Leaf	Seed	Pod
i	Time					
	Peak flowering	21.5 b	194 b	355 b	ı	150 bb
	Pod filling	38.0 ℃	178 b	206 a	348 b	406 c
	Maturity	4.7 a	67.2 a	1	269 a	35.0 a
II.	Cultivars					
	Black Turtle Soup	20.8 n	223 n	342 n	307 m	205 n
	Seafarer	15.3 m	70.5 m	218 m	309 ш	190 m
III.	CO <sub>2</sub> Level					
	300 ppm		123 x	139 x	340 y	210 y
	mdd 008	24.1 Y	170 Y	422 Y	277 ×	184 x
Inter	Interactions					
	IXI	*	*	n.s.	*	n.s.
	IX I	*	*	*	*	*
	II x III	n.s.	*	*	n.s.	*
	I x II x III	n.s.	*	*	n.s.	n.s.

 $^{\rm a}{\rm Means}$  within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.

bSeed not separate from pod.



accumulation was greater in the vegetative organs and pods in Black Turtle Soup than in Seafarer, however, no differences were evident in seed (Table 3). No interaction was seen between carbon dioxide levels and cultivars in root starch concentration (Table 3). Although starch concentration in the stems increased significantly with increasing carbon dioxide concentration in both cultivars (Table 3), the increase was not significant in Seafarer (Table 4). Starch concentration was consistently low in Seafarer stems even under field conditions; this may be characteristic of the determinate growth habit. Carbon dioxide enrichment markedly increased starch accumulation in leaves of both cultivars (Table 4). Interaction was again significant (Table 3), with a 4-fold increase in Seafarer as compared with a 3-fold increase in Black Turtle Soup (Table 4). Carbon dioxide-cultivar interaction in seed starch concentration was nonsignificant (Table 3).

The concentration of starch in the vegetative tissues decreased with maturity (Table 3). This decrease, especially in stems and leaves coincided with an increase in starch concentration in seeds and pod walls. Starch concentration in all tissues was lower at maturity than at pod filling (Table 3), possibly due to hydrolysis.

Significant interactions occurred between cultivars and starch concentration in roots, stems and seeds

. Fr 1,000

2011

- 1 A

. . .

.


. . . .

Table 4

Effects of  ${\rm CO}_2$  and Cultivar on Starch Concentration (mg glucose/g dry weight) in Bean Tissues<sup>a</sup>


Carbon Dioxide				Tissue		
Level (ppm)	Cultivar	Root	Stem	Leaf	Seed	Pod
300	Black Turtle Soup Seafarer		182.2 b 63.5 a	186.6 b 92.1 a	13.1 a 182.2 b 186.6 b 295.0 b 11.1 a 63.5 a 92.1 a 292.2 b	211.9 b 208.6 b
	Mean	12.1 x	12.1 x 122.8 x 139.3 x 293.6 y	139.3 x	293.6 Y	210.2 y
800	Black Turtle Soup Seafarer	28.5 c 19.6 b	263.0 c 77.4 a	498.4 d 344.9 c	28.5 c 263.0 c 498.4 d 221.0 a 19.6 b 77.4 a 344.9 c 216.2 a	197.6 b 171.1 a
	Mean	24.0 y	170.2 y	421.6 Y	24.0 y 170.2 y 421.6 y 218.6 x	184.3 x

 $^{\rm a}{\rm Means}$  within columns and sets (abc, xyz) with similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.



during the reproductive phase (Table 3). In roots, the starch concentration was higher in Black Turtle Soup at all developmental stages except maturity (Fig. 1A), but the difference was minimal at peak flowering. On the other hand, the difference was maximal at this time in the stem (Fig. 1B). In both cultivars, the starch levels in roots and stems decreased as the seeds matured (Table 3). In leaves there was no interaction between cultivar and growth stage during pod development (Table 3). As the pods developed, the starch concentration in leaves decreased (Table 3). Cultivar-developmental stage interaction on seed starch was shown in Figure 2, where starch concentration in seeds of both cultivars increased from peak flowering to pod filling, then decreased at maturity (Fig. 2).

Interaction between carbon dioxide levels and stage of pod development on starch concentration was noted (Table 3). In root, stem and leaf, carbon dioxide enrichment increased starch accumulation (Table 5), but response varied considerably with time. However, in seed, carbon dioxide enrichment decreased starch concentration regardless of the stage of pod development (Table 5). The greater number of seeds produced apparently reduced the starch available for storage in each seed. The rapid decline in starch concentration in the pod wall at maturity (Table 5) paralleled data obtained in a previous






Fig. 2. Levels of starch in seed of two <u>Phaseolus vulgaris</u> L. cultivars during their reproductive phase, averaged from the two carbon dioxide levels.





Table 5

Effects of Carbon Dioxide and Time of Sampling on Starch Concentration (mg glucose/g dry weight) of Bean Tissues^A

Carbon Dioxide	Time After			Tissues		
Level (ppm)	(days)	Root	Stem	Leaf	Seed	Pod
300	0	32.3 d		166.7 b		
	10	12.0 b	197.9 d	225.4 c	1	200.7 c
	20	22.2 c		53.3 a	405.1 c	400.6 d
	40	2.0 a		1		
800	0	32.3 d	97.4 b	166.7 b	1	
	10	31.1 d	189.7 d	7	1	100.5 b
	20	33.7 d	217.3 d	358.3 d	291.8 b	412.1 d
	40	7.4 ab	103.6 b		263.6 a	40.5 a

 $^{\rm a}{\rm Means}$  within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.



experiment (Chapter 4), and suggests rapid utilization of starch from the pod wall by the growing seeds.

### Relation of Pod Abscission to Available Carbohydrates

The data presented in Chapter 5 showed that carbon dioxide enrichment during the reproductive phase increased plant size, seed yield, and pod retention. In this study, using the same growing conditions, increasing the carbon dioxide concentration increased free sugar and starch in root, stem, and leaf. Available carbohydrate concentration paralleled vegetative growth, seed yield, and pod retention, suggesting a direct influence of carbohydrate in limiting abscission. However, correlation coefficients between percentage of pod abscission and free sugar and starch levels under the two carbon dioxide levels did not reveal any significant relationships (Table 6). Nonsignificant negative correlations were observed between abscission rates and levels of available carbohydrate. Starch concentration of stems was negatively correlated with percentage of abscission in both cultivars at least at pod filling (Table 6). This may indicate that the carbohydrates temporarily stored in stems are used to retain pods. No significant correlations were evident between available carbohydrate in vegetative tissues or stem weight and seed yield (Table 7 and 8). Abscission is a complex process whose mechanism is not fully




Table 6

Correlation Coefficients Between Percentage of Pod Abscission and Available Carbohy-drate Fractions in Vegetative Tissues in Two Dry Bean Cultivars Under Two Levels of Carbon Dioxide

	2,000	Black Tur	Black Turtle Soup	Seafarer	ırer
	כוומד מכרכד מ	300 ppm CO <sub>2</sub>	800 ppm CO <sub>2</sub>	300 ppm CO <sub>2</sub>	800 ppm CO <sub>2</sub>
ı.	At Pod Filling Stage				
	Free sugar in root	78b	.17	.29	.05
	Free sugar in leaf	.02	09	.41	12
	Starch in stem Starch in leaf	68 a	.19	21	75b
II.	At Maturity				
	Free sugar in root Free sugar in stem Starch in root Starch in stem	59	41 34 09	1.19	.2124

asignificant at 5% level.

bSignificant at 1% level.




Table 7

Correlation Coefficients Between Seed Weight Per Plant and Available Carbohydrate Fractions in Vegetative Tissues in Two Dry Bean Cultivars Under Two Levels of Carbon Dioxide

	200000000000000000000000000000000000000	Black Tur	Black Turtle Soup	Seafarer	arer
	Citataccers	300 ppm CO <sub>2</sub>	800 ppm CO <sub>2</sub>	300 ppm CO <sub>2</sub>	800 ppm CO <sub>2</sub>
i.	At Pod Filling Stage				
	Free sugar in root	66ª	20	10	45
	Free sugar in stem	.41	.67ª	.27	.16
	Free sugar in leaf	.12	.04	.48	.16
	Starch in root	90	21	45	43
	Starch in stem	59	70ª	.30	62ª
	Starch in leaf	57	.21	.30	14
II.	At Maturity				
	Free sugar in root	.38	.22	60.	20
	Free sugar in stem	90	.12	.15	02
	Starch in root	60.	51	.26	.18
	Starch in leaf	.52	25	.13	.41

<sup>a</sup>Significant at 5% level.

bsignificant at 1% level.





Table 8

Correlation Coefficients Between Stem Weight Per Plant and Available Carbohydrate Fractions in Vegetative Tissues in Two Dry Bean Cultivars Under Two Levels of Carbon Dioxide

	Characters	Black Tu	Black Turtle Soup	Seafarer	arer
		$300 \mathrm{\ ppm\ CO}_2$	800 ppm CO <sub>2</sub>	300 ppm co <sub>2</sub>	800 ppm co <sub>2</sub>
i.	At Pod Filling Stage				
	Free sugar in root	.14	16	47	0.4
	Free sugar in stem	49	53	44	500
	Free sugar in leaf	.08	07	. 41	21. 1
	Starch in root	.31	- 53	- 15	
	Starch in stem	.57	16	. 23	32
	Starch in leaf	15	19	90.	.31
II.	At Maturity				
	Free sugar in root	.32	.64ª	.25	- 26
	Free sugar in stem	.23	.64ª	15	17
	Starch in root	.35	.48	60	7.5
	Starch in stem	.33	.43	17	41

<sup>a</sup>Significant at 5% level.

bSignificant at 1% level.




understood. Available carbohydrate probably plays only an indirect role in the phenomenon.

6.19

÷

#### REFERENCES

- Cooper, R. L. and W. A. Brun. 1967. Response of soybeans to a carbon dioxide enriched atmosphere. Crop Sci. 7: 455-457.
- Guinn, G. 1974. Abscission of cotton floral buds and bolls as influenced by factors affecting photosynthesis and respiration. Crop Sci. 14: 291-293.
- Hardman, L. L. and W. A. Brun. 1971. Effect of atmospheric carbon dioxide enrichment at different developmental stages on growth and yield components of soybeans. Crop Sci. 11: 886-888.
- Krenzer, Jr., E. G. and D. N. Moss. 1975. Carbon dioxide enrichment effects upon yield and yield components in wheat. Crop Sci. 15: 71-74.
- Wittwer, S. H. and W. M. Robb. 1964. Carbon dioxide enrichment of greenhouse atmospheres for food crop production. Econ. Bot. 18: 34-56.



#### CHAPTER 7


ABSCISSION OF FLOWERS AND FRUITS IN PHASEOLUS VULGARIS L.

VI. THE RELATIONSHIP BETWEEN POD ABSCISSION AND

ENDOGENOUS ABSCISIC, PHASEIC, AND DIHYDRO
PHASEIC ACIDS IN THE PEDICELS AND PRUITS

# Abstract

In two cultivars of dry bean, populations of fruits having low vs. high abscission potential were established by removing early-opening flowers from half the plants. Fruits were harvested 4 to 5 days after anthesis and separated according to length, which was negatively correlated with abscission potential. Abscisic acid (ABA), phaseic acid (PA), and dihydrophaseic acid (DPA) contents were determined in methanol extracts of both pods and pedicels, using electron capture gas liquid chromatography (GLC). None of the treatments affected the content of these 3 compounds in pedicels. ABA content of pods was positively correlated with abscission potential of the 2 cultivars and with fruit load, but was unaffected by fruit size. PA content increased with fruit size and with fruit load, but was not affected by cultivar. DPA content was greater in the cultivar exhibiting the higher rate of



abscission, but was unaffected by pod size or fruit load. I conclude that levels of extractable ABA, PA, and DPA do not regulate fruit abscission in bean.

albads :

## Introduction

During fruit abscission, both persisting (actively growing) and abscising fruits occur in the same population (3, 7). In order to make valid comparisons, one must be able to recognize and separate persisting and abscising fruits. Earlier attempts to do so were based on criteria such as fruit size, color, or loosening (2, 3, 5), which are useful only in the terminal stages of abscission (7), well after induction has occurred.

Growth inhibitors have been frequently viewed as potential abscission-regulation hormones (2, 4). Abscisic acid has been identified and believed to have this abscission regulating property (7). In peach, greater levels of abscisic acid was found in abscising (i.e. loose) than in persisting (i.e. tight) fruits (5). A strong correlation between abscisic acid content and abscission was reported in cotton fruit (2). However, the abscisic acid data presented in that paper could also be interpreted to be correlated with fruit growth. The evidence of abscisic acid levels correlated with growth rates was presented in soybean pods (9). Studying abscission in peach fruit, Zucconi (12) found levels of abscisic acid to be negatively correlated with abscission.

My purpose was to test the hypothesis that ABA content controls fruit abscission in bean. Differential treatments, which had previously been shown to result in

growie

11 .11

614.6

4 .

1 - "

1 .....

101.

14

25

.

. .

-

. . .

1 + + 71 a

. Jaw?

CONDLICT.

different rates of abscission, were applied, and pods were sampled during the induction phase for subsequent analysis. PA and DPA, previously shown to be metabolites of ABA (6), were also measured.

### Materials and Methods

#### Plant Culture

Two Phaseolus vulgaris L. cultivars, "Black Turtle Soup" and "Seafarer," were selected for their contrasting growth habits and abscission potentials. Seeds were planted in soil mixture (1:1:1 ratio of sand: sandy loam and peat) in 25 cm. pots in a greenhouse maintained at 24 ± 3°C on February 16, 1976 for Black Turtle Soup and February 21, 1976 for Seafarer. The reason for sowing at different times is to get the two cultivars flowering at the same time. On emergence, the seedlings were thinned to one per pot. When the plants reached flowering stage on March 26, 1976, they were divided into two groups of 150 plants of each cultivar and differential treatments applied.

Flowers opening six days or later after the first flower on the same plant have a high abscission potential in comparison with those opening earlier (Chapter 2). In the control treatment, flowers borne on the first five days were left on the plant. In the second group of plants, all flowers opening during the first five days were removed. Flowers opening the sixth day and thereafter

- Hestin

2 M - 13 DW

1921

1

.

4

95

. .

Wer-

were tagged with color-coded wire, and the pods from these tagged flowers were harvested at 2-day intervals, immediately frozen on dry ice and stored at  $-10\,^{\circ}\text{C}$ .

Eighty plants of each cultivar and treatment were used to record the rate of abscission. For each cultivar and treatment, 40 flowers that were borne on day six were tagged and individually numbered. Pods from these tagged flowers were measured every day and abscission was recorded.

After freeze-drying, pods sampled on day 4 and 5 were divided into 3 groups: small (< 1.5 cm), medium ( 1.5-3.0 cm) and large (> 3.0 cm). An abscission potential was assigned to each subgroup based on the relationship between pod length and abscission probability determined experimentally from comparable pods monitored (in vivo) throughout their development.

# Extraction

Pedicels (20 mg) and fruits (100 mg) were homogenized in 10 ml methanol, following the addition of 100 ng t, t-ABA to correct for losses. No t, t-ABA was recovered from nonspiked samples, and the amount of c, t-ABA recovered from blanks containing 5'5-ABA alone was negligible. The homogenates were left for 6 to 12 hours at 2°C in the refrigerator, then filtered, and the tissue was rinsed with methanol. The filtrate was reduced to

dryness in a flash evaporator at 40°C and resuspended in 20 ml distilled water.

## Fractionation

The extract was adjusted to pH 8.0 with 0.1 N  ${\rm NH_4OH}$  and washed 3 times with 10 ml methylene chloride  $({\rm CH_2Cl_2})$ . The  ${\rm CH_2Cl_2}$  fraction was discarded. The water phase was adjusted to pH 3.0 with 0.1 N formic acid and partitioned 3 times with 15 ml ethyl acetate (EtAc) to obtain the acidic fraction, which was evaporated and stored at -10°C until analyzed.

# Analysis

The samples were resuspended in 0.5 ml methanol and methylated with diazomethane according to Schlenk and Gellerman (8). The methylated samples were evaporated under nitrogen, the residues resuspended in 2 and 1 ml ethyl acetate for the pod and pedicel samples, respectively, and 2  $\mu$ l of each sample was injected into a Packard 7300 Gas-Liquid Chromatograph equipped with a 63 Ni electron capture detector operated at 5v. The column was 2 mm i.d X 1.83 m, packed with 1% XE-60, 12,500 Centistrokes (CS or CSTK - the viscosity rating), on Gaschrome Q 80/100 mesh. Column, inlet and detector temperatures were 200°, 240°, and 240°C, respectively. The carrier gas was N<sub>2</sub> at a flow rate of 40 ml/min at 40 psi. Nitrogen scavenger gas was supplied to the detector at 90 ml/min.

dryne s. Sugar.

. 50, 0a

Data were quantified by measuring peak heights and comparing with a standard curve (semi-log) for each compound measured. Peak heights were linear over a range of 0.05 to 1.0 ng/µl ABA, 0.10 to 2.0 ng/µl PA and 0.05 to 2.0 ng/µl DPA. Data were corrected for losses during extraction and fractionation from the percentage of recovery of the standard t,t-ABA, assuming parallel losses, in c,t-ABA, PA and DPA.

#### Results

#### Pod Growth vs. Abscission

Persisting and abscising pods could not be distinguished on the basis of length during the first three days after pod setting. However, the growth rate of abscising pods began to decrease thereafter, and the differences in pod size increased until the pods abscised from the plants (Fig. 1A and B).

Control vs. deflowered plants differed markedly in percentage of pod abscission. Only 30 and 20% of the pods abscised in the latter vs. 55 and 75% in the former for "Black Turtle Soup" and "Seafarer" respectively. Within treatments and cultivars, the degree of abscission was strongly correlated with pod size on days 4 and 5, smaller pods having a much greater tendency to abscise (Table 1).

4 -4

to the bas

a ja disa

11.00

15.

17.0

. ...

\* 311

.

6 224




Fig. 1. Growth curves of abscising and persisting pods. A. Black Turtle Soup; B. Seafarer.

### Hormone Content vs. Abscission

ABA. ABA content of "Seafarer" pods was higher than that of "Black Turtle Soup" pods, particularly in the control treatment (Table 1). Deflowering reduced ABA content in all cases. However, the effect of pod size was inconsistent. Thus, ABA content paralleled abscission in the case of cultivar and treatment effects, but not in size effect. None of the treatments affected ABA content of the pedicels.

PA. Low levels of PA were detected in both pod and pedicel. PA was higher in pods from control plants than in those from deflowered plants in all but one case (Table 1). However, large pods contained more PA than medium or small pods, indicating no correlation with abscission. Cultivar did not affect PA content. PA levels in the pedicels were not affected by any of the treatments.

<u>DPA</u>. "Seafarer" pods contained significantly more DPA than did "Black Turtle Soup" pods (Table 1). However, neither deflowering nor pod size affected DPA content of the pedicels.

## Discussion

Although the work of both Martin and Nishijima
(5) and Davis and Addicott (2) suggests a positive
relationship between the content of ABA or ABA-like

.. ei


JEE.

- . . . (6)

Levels of ABA, PA and DPA in Relation to Percentage of Abscission of the Two Bean Cultivars $^{\rm A}$ Table 1

Small Small Medium Large Small Medium	Abscission							
Small Medium Large Large Seafarer Small Medium		Abscission	Æ,	ABA		PA		DPA
			Pod	Pedicel	Pod	Pedicel	Pod	Pedicel
	Long	46	10.0	19.3	0	2.3	4.2	7.8
	High	99	65.0	24.0	0	0.7	25.3	8.1
		c	10.4	17.5	0	0	8.0	5.0
	High	33	73.0	16.7	9.9	1.7	22.6	6.4
		c	8.0	18.0	0	0	1.0	3.9
	High	00	0.66	32.8	64.7	0	13.4	4.7
	,	1.1	101.0	23.9	0	2.5	52.9	5.6
Medium	High	88	159.0	19.7	11.8	1.3	36.0	5.6
Medium		•	0 03	22.9	0	0	41.6	3.5
	LOW	33 0	101.0	17.7	4.0	1.8	31.0	4.7
	наты				, ,,	0 0	48 6	6.9
Larde	Low	00	81.0	14.4	61.8	0.7	49.0	2.4
	High							
Means for:		22 a	44.2 a	21.4 a	11.9 a	0.8 a	11.4 a	6.0 a
RTS			d 8.66	20.4 a	15.4 a	1.4 a	43.2 b	4.3 a
1000			03 7 ×	21.7 x	2.9 x	1.7 x	29.8 x	v 0.9
Small		97 p			2.6 x	x 6.0	24.0 x	4.9 x
		16 a	63.1 x					u
Medium		0	69.2 x	22.3 x	35.7 Y	0.7 ×	X 0.87	× .
903.5					2.7 m	1.3 m	24.8 m	5.4 m
המדפת	Low	10 a 35 b	46.4 m 97.7 n	20.9 m	24.8 n	ш 6.0	29.7 m	4.8 m

Ageans within the same parameter followed by similar letters are not significantly different at the 5% level by Duncan's Multiple Range Test.




inhibitors in fruits and their abscission, Zucconi (12) has pointed out the fallacy of measuring ABA content of abscising fruits, viz., that one is measuring the result of abscission rather than its cause. ABA content should be measured during induction. Zucconi (12) actually demonstrated a negative correlation between content of an ABA-like inhibitor and the induction of abscission in peach fruits.

In the present study, ABA content of the fruits paralleled relative abscission potential as affected by cultivar and treatment (deflowering), but fruit size, which greatly affected abscission, had little effect on ABA content. I conclude from this that the cultivar and treatment effects on ABA content were fortuitous, rather than being causally related to abscission. If ABA content were in fact responsible for abscission, one would expect differences to be apparent in the pedicel, where abscission actually occurs. However, this was not the case.


PA and DPA have not been implicated directly in abscission, and their levels appear to bear little relationship to fruit retention. There was no correlation between relative levels of PA vs. DPA in the pods; PA levels were significantly affected by deflowering and by pod size, while DPA content varied only with cultivar.

If ABA regulates abscission, it must do so independently of concentration in the tissue as a whole. Concentrations might be very different in the cells of the abscission zone itself, for example. On the other hand, diffusible ABA could control abscission independently of the size of the pool of extractable ABA.



#### REFERENCES

- Addicott, F. T. and J. L. Lyon. 1969. Physiology of abscisic acid and related substances. Ann. Rev. Plant Physiol. 20: 139-164.
- Davis, L. A. and F. T. Addicott. 1972. Abscisic acid: correlations with abscission and with development in the cotton fruit. Plant Physiol. 49: 644-648.
- Leuty, S. J. and M. J. Bukovac. 1968. A comparison of the growth and anatomical development of naturally abscising, non-abscising and naphthaleneacetic acid-treated peach fruits. Phytomorphology 18: 372-379.
- Luckwill, L. C. 1953. Studies of fruit development in relation to plant hormones. I. Hormone production by the developing apple seed in relation to fruit drop. J. Hort. Sci. 28: 14-24.
- Martin, G. C. and C. Nishijima. 1972. Levels of endogenous growth regulators in abscising and persisting peach fruits. J. Amer. Soc. Hort. Sci. 97: 561-565.
- Milborrow, B. V. 1974. The chemistry and physiology of abscisic acid. Ann. Rev. Plant Physiol. 25: 259-307.
- Ohkuma, K., J. L. Lyon, F. T. Addicott, and O. E. Smith. 1963. Abscisin II, an abscissionaccelerating substance from young cotton fruit. Science 142: 1592-1593.
- Powell, L. E. and C. Pratt. 1966. Growth promoting substances in the developing fruit of peach (Prunus persica L.). J. Hort. Sci. 41: 331-348.
- Quebedeaux, B., P. B. Sweetser, and J. C. Rowell. 1976. Abscisic acid levels in soybean reproductive structures during development. Plant Physiol. 58: 363-366.



- Schlenk, H. and J. L. Gellerman. 1960. Esterification of fatty acids with diazomethane on a small scale. Anal. Chem. 32: 1412-1414.
- Swanson, B. T. Jr., H. F. Wilkins, C. F. Weiser, and I. Klein. 1975. Endogenous ethylene and abscisic acid relative to phytogerontology. Plant Physiol. 55: 370-376.
- Zucconi, F. 1975. Reassessment of the relationship between hormonal and developmental changes during abscission with particular reference to peach (<u>Prunus persica L. Batsch) fruit. Ph.D. dissertation, Michigan State University.</u>

#### CHAPTER 8


#### SUMMARY AND CONCLUSIONS

Elucidation of the causes of physiological processes is difficult, especially in an area as complex as abscission. However, an effort must be made to generate reasonable hypotheses, and to design appropriate experiments to test them, if progress is to be made.

In this work, differences were established in abscission potential of reproductive structures in dry bean cultivars.

Among the bean cultivars studied, Black Turtle
Soup and Seafarer were selected for their contrasting
growth habits and abscission rates. Black Turtle Soup, an
indeterminate type, had a lower pod abscission rate than
Seafarer, a determinate bush bean.


Exposure of either cultivar to long photoperiods or high carbon dioxide levels after the onset of flowering increased vegetative growth and leaf area, reduced pod abscission and consequently resulted in higher seed number and greater seed yield. Black Turtle Soup was more responsive to both factors than was Seafarer.




Levels of free sugar and starch in root, stem and leaf increased with photoperiod and carbon dioxide level. Available carbohydrate concentration in vegetative tissues was higher in the Black Turtle Soup cultivar, consistent with larger plants, greater leaf area and a lower abscission rate. However, the results provide only correlative evidence in support of the carbohydrate hypothesis.

Levels of abscisic acid and its related compounds in pod and pedicel were not related to the role of abscission in either cultivar. Young pods of Seafarer contained more abscisic acid and dihydrophaseic acid than did those of Black Turtle Soup. Levels in the pedicel were not affected by cultivar.

In conclusion, pod abscission in dry bean is not totally dependent on available carbohydrate levels. Many nutritional factors such as nitrogen, or carbohydrate/ nitrogen ratio could regulate abscission. Although endogenous abscisic acid content was not correlated with abscission, other naturally occurring growth regulators, such as ethylene, cytokinin and auxin could play a role. The interplay between the nutritional and hormonal levels at a particular stage of development may control pod abscission. Since abscission is a complex process influenced by many environmental factors, care must be taken to design critical experiments to test available hypotheses.







# APPENDIX A

GROWTH AND POD ABSCISSION AS AFFECTED BY DIFFERENT DAYLENGTHS




Table A-1

Effects of Daylengths on Number of Flowers, Number of Pods and Percentage of Pod Abscission at 3 Stages of Development of Two Dry Bean Cultivars Grown Under Greenhouse Conditions in 1975<sup>a</sup>

Dev. Stages	Cultivar		Daylengths	ngths			Dayl	Daylengths	w		Dayle	Daylengths	
		80	13	18	Mean	80	13	18	Mean	ω	13	18	Mean
Peak Flowering B	Black Turtle Soup	7	00	17	10	7	8	16	10	1	1	,	1
	Seafarer	12	59	31	24	12	27	30	23	1	1	,	1
×	Mean	10	18	24	17a	6	11	23	16c	,	ı	ı	1
Pod Filling	Black Turtle Soup	00	6	17	11	S	7	16	6	29.6	21.1	9.3	20.0
	Seafarer	12	27	32	24	2	6	19	11	48.5	66.5	42.1	52.3
zi	Mean	6	18	25	17a	S	8	11	10b	39.0	43.8	25.7	36.2a
Martin Ma	Black Turtle Soun	6	œ	17	11	6	2	12	7	63.0	30.1	29.7	40.9
	Seafarer	14	59	32	26	m	7	16	8	73.8	71.8	53.6	66.4
, 2,	Mean	::	18	56	19a	m	9	14	8a	68.4	50.9	41.7	53.7b
Mean	ď	10×	187	252		¥9	6× 10y	182		53.7z	53.7z 47.4y 33.7x	33.7x	
	BTS		11	E				E 6			30.5	2 m	
cultivar -	Seafarer		25	25 n				14 n			59.4	4 n	
Interactions	,												
1) Dow stade x cultivar	tivar		ď	n.s.				*			n.	n.s.	
	toneriod		2	0				*			*		
3) Cultivar x photoperiod	period		*	*				*			*		
4) Dev. stage x cultivar	tivar							;			2	0	
x photoperiod			ď	n.s.									

<sup>8</sup>Means within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.




Table A-2

Effects of Daylengths on Number of Flowers, Number of Pods and Percentage of Pod Abscission at 3 Stages of Development of Two Dry Bean Cultivars Grown Under Greenhouse Conditions in 1976<sup>a</sup>

		No	No. of Flowers	Flowe	rs		No.	No. of Pods	ds	de	Pod A	& Pod Abscission	u.
Dev. Stages	Cultivar		Daylengths	ngths			Dayl	Daylengths	s		Dayle	Daylengths	
		80	13	16	Mean	80	13	16	Mean	8	13	18	Mean
Peak Flowering	Black Turtle Soup Seafarer Mean	6 14 10	6 18 12	9 15	7 10 11a	13	6 16 11	13	14 10c				1.1
Pod Filling	Black Turtle Soup Seafarer Mean	7 11	118 113	12 22 17	9 14a	ខេត	7 8 5	212	7 8 4 d	37.6 67.3 67.3	27.3 56.5 41.9	14.0 50.0 32.0	26.3 57.9 42.1a
Maturity	Black Turtle Soup Seafarer Mean	13	9 15	11 22 17	9 17 13a	24 E	0 9 0	r 8 8	la e a	52.0 74.6 63.3	40.6 62.7 51.6	33.6 61.1 47.3	42.1 66.2 54.1b
	Mean	10x	12y	152		×9	8y	102		57.9z	46.8y	39.7x	
Mean	BTS			E 8				ш 9			34.	34.18 m	
cultivar	Seafarer		L	17 n				u 6			62.	62.02 n	
Interactions													
The same of the same	100		2	0				*			u	n.s.	
	Dev. stade x photoperiod		ď	n.s.				*			n	s.	
	photoperiod		ď	n.s.			-	n.s.			ď	n.s.	
<ol> <li>Dev. stage x cultivar x photoperiod</li> </ol>	k cultivar od		ď	n.s.			-	n.s.			ц	n.s.	

<sup>a</sup>Means within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.




Table A-3

Effects of Daylengths on Leaf Area Per Plant, Number of Leaf Per Plant and Leaf Weight Per Plant at 3 Stages of Development of Two Dry Bean Cultivars Grown Under Greenhouse Conditions in 1975<sup>a</sup>

			Leaf Area/Plant	ea/Plan	t.	No	No. Leaf/Plant	f/Pla	nt	Leaf	Weigl	Leaf Weight/Plant (g)	t (g)
Dev. Stage	Cultivar		Daylengths	ngths			Daylengths	ngths			Dayle	Daylengths	
		80	13	18	Mean	80	13	18	Mean	8	13	18	Mean
Peak Flowering	Black Turtle Soup Seafarer Mean	1282 1135 1208	1478 1271 1374	1903 1297 1600	1554 1234 1394a	14	16 16 16	21 17 19	17 17 17a	2.3	3.3	7.1	4.3 3.1
Pod Filling	Black Turtle Soup Seafarer Mean	1229 1041 1135	1742 1350 1546	2111 1357 1734	1694 1250 1472a	13 13	15 17 16	20 20	16 17 16a	3.5	5.0 4.1	6.5	5.0 4.6b
	Mean	1172×	1172× 1460y	1667z		14x	16y	19z		2.9x	3.84	5.7x	
Mean	BTS		1624 n	c			16 1	E			4	4.7 n	
cultivar	Seafarer		1242 m	E			17	17 m			3	3.6 ш	
Interactions													
1) Dev. stage x	stage x cultivar		n.s.				'n.	n.s.			п	.8.	
2) Dev. stage x	Dev. stage x photoperiod		n.s.				ď	n.s.			d	n.s.	
	photoperiod		*				*					*	
<ol> <li>bev. stage x cultivar x photoperiod</li> </ol>	cultivar od		n.s.				ü.	n.s.			п	n.s.	

<sup>8</sup>Means within the same parameter followed by similar letters are not significantly different at the 18 level by Duncan's Multiple Range Test.





Table A-4

Effects of Daylengths on Leaf Area Per Plant, Number of Leaf Per Plant and Leaf Weight Per Plant at 3 Stages of Development of Two Dry Bean Cultivars Grown Under Greenhouse Conditions in 1976<sup>a</sup>

			Leaf Area/Plant	ea/Plan	ıt	N	No. Leaf/Plant	f/Pla	nt	Leaf	f Weig	Leaf Weight/Plant (g)	t (g)
Dev. Stage	Cultivar		Daylengths	gths			Daylengths	ngths			Dayl	Daylengths	
		8	13	18	Mean	8	13	18	Mean	œ	13	18	Mean
Peak Flowering	Black Turtle Soup Seafarer Mean	903 681 792	1071 907 989	1279 1108 1194	1084 899 991a	999	===	13 15	12 12 12a	1.9	2.2	3.1	2.3 2.3
Pod Filling	Black Turtle Soup Seafarer Mean	868 903 886	1121 1003 1062	1651 1241 1446	1214 1049 1131b	6 11 10	12 14 13	15 16 15	12 14 13b	1.5	2.3	3.5	2.3 2.3
	Mean	839x	1026y	1026y 1320z		10×	12y	152		1.6x		2.1y 3.5z	
Mean	BTS		1149 n	п			12	12 m			2	2.5 m	
cultivar	Seafarer		974	E			13	13 m			2	2.3 ш	
Interactions													
1) Dev. stage x cultivar	cultivar		n.s.				'n.	n.s.			ď	n.s.	
	photoperiod		n.s.				n.	n.s.					
	hotoperiod		n.s.				n.	n.s.			ď	n.s.	
<ol> <li>Dev. stage x cultivar x photoperiod</li> </ol>	cultivar		n.s.				'n.	n.s.			ď	n.s.	

Ameans within the same parameter followed by similar letters are not significantly different at the 18 level by Duncan's Multiple Range Test.



able A-5

Effects of Daylengths Effects on Root Weight Per Plant, Stem Weight Per Plant and Total Plant Weight at 3 Stages of Development of Two Dry Bean Cultivars Grown Under Greenhouse Conditions in 1975<sup>a</sup>

		Root	Root Weight (g)/Plant	(d) :	lant	Stem	Weigh	Stem Weight (g)/Plant	Plant	To	tal Pla	Total Plant Wt. (g)	(6)
Dev. Stage	Cultivar		Daylengths	gths			Dayle	Daylengths			Dayle	Daylengths	
		8	13	18	Mean	ω	13	18	Mean	80	13	18	Mean
Peak Flowering	Black Turtle Soup Seafarer Mean	::::	1.2	3.4	2.1 1.7 1.9c	1:3	1.6	3.6	2.2 1.7 2.0b	5.2	5.3	13.1 8.2 10.7	8.2 6.1 7.2a
Pod Filling	Black Turtle Soup Seafarer Mean	0.7	0.8	2.0 1.6	1.2 1.0 1.1b	1.2	1.8	2.9	2.1 1.8 2.0b	5.4	12.6 12.8 12.7	18.2 19.0 18.6	12.7 12.4 12.5c
Maturity	Black Turtle Soup Seafarer Mean	0.0	0.0	1.0	1.0 0.9a	1.1	1.3	2.3	1.6 1.6	5.00	10.0 8.0 9.0	20.6 17.1 18.9	12.2 10.4 11.3b
	Mean	0.8x	1.0y	2.0z		1.3x	1.6y	2.72		5.7x	9.2y	9.2y 16.0z	
Mean	BTS		1	1.4 n			1	1.9 n			11	11.0 n	
cultivar	Seafarer		i,	1.1 m			1.	1.7 m			6	ш 9.6	
Interactions													
1) Dev. stage x cultivar 2) Dev. stage x photoper:	Dev. stage x cultivar Dev. stage x photoperiod		4.5	n.s.			•	* *			ď*	***	
	Cultivar x photoperiod		'n	n.s.			•	*			ď	n.s.	
<ol> <li>Dev. stage x cultivar x photoperiod</li> </ol>	x cultivar od		ď	n.s.			-	n.s.			*		

<sup>a</sup>Means within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.





Table A-6

Effects of Daylengths on Root Weight Per Plant, Stem Weight Per Plant and Total Plant Weight at 3 Stages of Development of Two Dry Bean Cultivars Grown Under Greenhouse Conditions in 1976<sup>a</sup>

		Root	Root Weight (g)/Plant	t (g)/	Plant	Stem	Stem Weight (g)/Plant	t (g)/	'Plant	To	tal Pl	Total Plant Wt.	(6)
Dev. Stage	Cultivar		Daylengths	ngths	-		Dayle	Daylengths			Dayl	Daylengths	
		80	13	18	Mean	80	13	18	Mean	00	13	18	Mean
Peak Flowering	Black Turtle Soup Seafarer Mean	0.7	0.8	1.1	0.9 1.0	6.0	1.3	1.3	1.0 1.3	9.29	5.2	6.3	4.5 5.1 4.8a
Pod Filling	Black Turtle Soup Seafarer Mean	0.5	8.00	1.8	0.9 1.1 1.0a	0.0	1.333	2.2	1.5 1.5 1.5b	5.0	6.7 7.6 7.1	12.7 13.7 13.2	7.8 8.8 8.3b
Maturity	Black Turtle Soup Seafarer Mean	0.0 8.0	0.8	1.1	0.7 1.0 0.9a	0.5	1.1	1.5 2.1 1.8	1.0 1.4 1.2a	3.6	7.7	10.5 12.5 11.5	6.9 8.1 7.5b
	Mean	0.5x	0.87	1.52		0.9x	1.13	1.92		3.9x	6.47	10.3z	
Mean	BTS		0.8	0.86 m			4	1.2 m			9	6.4 m	
cultivar	Seafarer		1.0	1.04 n			1.	1.4 n			7	7.3 m	
Interactions  1) Dev. stage x cultivar 2) Dev. stage x photoperion 3) Cultivar x photoperiod 4) Dev. stage x cultivar x photoperiod	Dev. stage x cultivar Dev. stage x brouperiod Dev. stage x photoperiod Cultivar x photoperiod Dev. stage x cultivar x photoperiod		u** u	n.s. n.s.			i∗ i i	n * s. s.			u* c c	*. *	

Ameans within the same parameter followed by similar letters are not significantly different at the 18 level by Duncan's Multiple Range Test.



## APPENDIX B

RELATIONSHIP OF POD ABSCISSION TO DIFFERENTIAL LEVELS

OF CARBOHYDRATES PRODUCED BY VARIATION

IN DAYLENGTH




Table B-1

Effects of Developmental Stages and Daylength on Free Sugar Levels in Vegetative Tissues of Two Phaseolus vulgaris L. Cultivars (data in mg glucose per gram dry weight)<sup>a</sup>

			Root	ot			St	Stem			Į,	Leaf		
Dev. Stage	Cultivars		Daylength (hrs)	th (hrs			Dayleng	Daylength (hrs)			Dayleng	Daylength (hrs)		
		8	13	18	Mean	80	13	18	Mean	ω	13	18	Mean	
Peak Flowering	Black Turtle Soup	11.8	16.0	17.0	14.9	81.4	95.2	88.6	4.88	49.3	63.1	7. 67	7 19	
,	Seafarer Mean	10.4	11.2	9.4	9.9 12.4a	53.9	67.6	61.3	74.60	38.9	58.6	50.3	49.3	
Pod	Black Turtle													
Filling	dnos	16.2	21.9	22.7	20.3	57.8	83.7	88.9	76.8	67.2	7.06	8.69	75.9	
	Seafarer Mean	13.6	18.5	13.3	14.0 17.1c	36.8	66.8	74.0	48.6 62.7b	55.9 61.5	73.4	56.7	56.3 66.1b	
Maturito	Black Turtle													
	dnog	9.0	11.2	19.8	13.3	21.6	35.7	58.5	38.6	1	,			
	Seafarer	13.2	13.9	15,3	14.1	21.0	34.9	38.0	31.3	1	1	1	1	
	Mean	11.1	12.5	17.5	13.7b	21.3	35.3	48.2	34.9a				1	
Mean: Day	Daylength	12.1x	14.9y	16.32		45.4x	61.2y	65.7z		52.8x	67.22	62.4y		
Mean	BTS		16.	16.2 n			67.	n 6.79			68.8	u 8		
cuttival	Seafarer		12.7	7 m			47.	47.0 m			52.8	E 8		
Interactions	suc													
1) Dev. st	Dev. stage x cultivar		•	*			•	:				*		
2) Dev. st	Dev. stage x daylength		•	**			•	**			*	*		
	Cultivar x daylength		•	**				*				*		
<ol> <li>Dev. stage x daylength</li> </ol>	Dev. stage x cultivar x daylength		ď	n.s.							*	*		
	The state of the s	-												١

Amens within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.




Table B-2

Effects of Developmental Stages and Daylength on Free Sugar Levels in Reproductive Tissues of Two Phaseolus vulgaris L. (data in mg glucose per gram dry weight)  $^{\rm a}$ 

			Seed	q			Pod Wall	Vall	
Dev. Stage	Cultivars		Daylength (hrs)	h (hrs)			Daylength (hrs)	th (hrs)	
		8	13	18	Mean	8	13	18	Mean
Peak Flowering	Black Turtle Soup		,	,		173	178	208	187
	Seafarer	1	ı	1	1	146	170	183	166
	Mean	,	,	ı	1	159	174	195	176c
Pod Filling	Black Turtle Soup	52.3	63.7	103.8	73.3	151	130	192	158
	Seafarer	46.1	58.4	75.2	59.9	140	113	120	124
	Mean	49.2	61.0	89.5	66.6a	145	122	156	141b
Maturity	Black Turtle Soup	57.8	69.4	74.2	67.1	21.4	21.2	25.4	22.7
To a month	Seafarer	70.8	76.4	80.1	75.8	32.2	27.3	29.1	29.5
	Mean	64.3	72.9	77.1	71.5a	26.8	24.2	27.2	26.la
Mean: daylength	ч	56.7x	¥6.99	83.3z		110.7x	106.7x	126.4y	
Mean	BTS		70.2	E			122	122.4 n	
cultivar	Seafarer		67.8	E 8			106.8	m 8.	
Interactions									
	2000		*				*	*	
	CULLIVAL		*				*	*	
2) Dev. stage x daylength	daylength		2	0			**	*	
<ol> <li>Cultivar x daylength</li> <li>Dev stade x cultivar</li> </ol>	lay tength								
			ď	n.s.			*		

 $^{\rm a}$ Means within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.




Table B-3

Effects of Developmental Stages and Daylength on Starch Levels in Vegetative Tissues of Two Phaseolus vulgaris L. Cultivars (data in mg glucose equivalent per gram dry weight) a

			Roc	Root			St	Stem			Le	Leaf	
Dev. Stage	Cultivars		Jayleng	Daylength (hrs)			Dayleng	Daylength (hrs)			Daylength (hrs)	th (hrs	
		8	13	18	Mean	8	13	18	Mean	8	13	18	Mean
Peak Flowering	Black Turtle Soup Seafarer	28.9	30.7	9.3	23.0	110.6	121.0	197.1	142.9	83.3	63.4	217.5	121.4
	Mean	25.5	28.1	7.8	20.4b	83.1	95.5	167.1	115.2b	75.5	47.3	146.8	89.9b
Pod Filling	Black Turtle Soup	5.0	15.7	25.0	15.2	62.9	162.0	242.2	156.7	15.2	53.6	30.2	33.0
	Seafarer Mean	19.9	19.3	13.1	17.5 16.4a	69.3	54.2	107.3	78.0 117.4b	18.8	39.5	14.5	24.3 28.7a
Maturity	Black Turtle												
	dnos	19.5	26.6	18.7	21.6	11.3	25.5	39.3	25.4	1 1	1 1	1 1	Ü
	Mean	18.3	26.2	20.9	21.8b	12.4	33.5	29.9	25.3a	1	ı	1	1
Mean: day	daylength	18.8x	23.9y	15.9x		54.9x		79.0y 123.9z		46.3x	46.9x	84.6y	
Mean	BTS		19.	ш 6.61			108	108.3 n			77.2 n	2 n	
cultivar	Seafarer		19.1	1 m			63	ез.е ш			41.3 m	3 m	
Interactions	suc												
	Dev. stade x cultivar			*			*	**			*	*	
2) Dev. st	Dev. stage x daylength		*	**			*	**			*	*	
	Cultivar x daylength		4	n.s.			*	*			*	*	
4) Dev. s	Dev. stage x cultivar x daylength		•	:			•	*			*	*	
- 711	and an												

\*Means within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.




Table B-4

Effects of Developmental Stages and Daylength on Starch Levels in Reproductive Tissues of Two Phaseolus vulgaris L. Cultivars (data in mg glucose equivalent per gram dry weight)  $^{\rm a}$ 

			Seed	gg			Pod Wall	a11	
Dev. Stage	Cultivars		Daylength (hrs)	th (hrs)			Daylength (hrs)	(hrs)	
		ω	13	18	Mean	80	13	18	Mean
Peak Flowering	Black Turtle Soup	,				36.3	58.8	9.69	54.9
	Seafarer	1	1	1	1	68.4	53.7	52.5	58.2
	Mean		1	1	ı	52.4	56.3	61.0	26.6b
Pod Filling	Black Turtle Soup	362.6	428.0	348.6	379.8	276.7	412.0	325.4	338.0
	Seafarer	372.1	456.7	366.0	398.3	341.8	424.3	366.6	377.5
	Mean	367.4	442.4	357.3	389.00	309.2	418.1	346.0	357.80
Maturity	Black Turtle Soup	283.7	371.1	386.3	347.0	8.4	30.3	64.2	34.3
	Seafarer	310.9	351.8	356.0	339.5	24.4	26.1	6.97	42.5
	Mean	297.3	361.4	371.2	343.3b	16.4	28.2	70.5	38.4a
Mean: daylength		332.3x	401.9z	364.2y		126x	167y	159y	
Mean	BTS		363.4	4 m			142	ш	
cultivar	Seafarer		368.9	ш 6			159 1	п	
Interactions									
The state of the s	******		2				*		
2) Dev. stage x currivat	daylength		*				*		
	laylength		*				*		
4) Dev. stage x cultivar x daylength	cultivar		n.s.				n.s.		

Ameans within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.




Table B-5

Effects of Developmental Stages and Daylength on Total Nonstructural Carbohydrate Levels in Vegetative Tissues of TWO Phaseolus <u>vulgaris</u> L. Cultivars (data in mg glucose per gram dry weight) <sup>a</sup>

			Root	ň			Stem	ma			Le	Leaf	
Dev. Stage	Cultivars		Daylength (hrs)	h (hrs)			Daylength (hrs)	h (hrs			Daylength (hrs)	h (hrs)	
		8	13	18	Mean	8	13	18	Mean	80	13	18	Mean
Peak Flowering	Black Turtle Soup Seafarer Mean	189.7 110.4 150.0	223.1 135.6 179.4	285.9 198.2 242.0	232.9 148.0 190.5b	40.4 31.1 35.8	46.5 34.2 40.3	20.9 13.6 17.2	35.9 26.3 31.1a	132.7 106.6 119.6	126.5 85.1 105.8	290.1 126.5 208.3	183.1 106.9 144.6b
Pod Filling	Black Turtle Soup Seafarer Mean	123.8 109.4 116.6	255.7 104.2 179.9	331.1 166.5 249.0	236.9 126.7 181.8b	22.4 30.1 26.3	36.8 31.2 34.0	43.5 26.1 34.8	34.2 29.1 31.7a	81.7 71.3 76.5	144.4 95.8 120.1	94.8 65.7 80.2	106.1 77.6 92.3a
Maturity	Black Turtle Soup Seafarer Mean	30.9 57.0 44.0	53.2 61.5 57.3	86.5 52.3 69.4	56.9 57.0 56.9a	27.9 26.9 27.4	37.3 37.9 37.6	36.9	34.1 34.8 34.4a	111		- cir	111
	Mean	103.6x	138.9y 186.7z	186.7z		29.8x	37.3y	30.1x		98.0x		112.9× 144.3y	
Mean	BTS		175.5 n	2 n			34.	34.7 n			145.	145.0 n	
cultivar	Seafarer		110.6	ш 9			30.1	1 m			91.	91.8 m	
Interactions 1) Dev. stag(2) Dev. stag(3) Cultivar 4) Dev. stag(4) Ev. stag(5)	eractions Dev. stage x cultivar Dev. stage x daylength Cultivar x daylength x daylength x daylength		*** *				**:*	* * * * *			*** *		

<sup>8</sup>Neans within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.





Table B-6

Developmental Stages and Daylangth Effects on Total Monstructural Carbohydrate Levels in Reproductive Tissues of Two Phasaclus vulgaris. L. Cultivurs (data are means of 10 plants, expressed as m g glucose per gram dry Weight!

			Se	Seeds			Pod	Pod Wall	
Dev. Stage	Cultivars		Dayleng	Daylength (hrs)			Dayleng	Daylength (hrs)	
		80	13	18	Mean	8	13	18	Mean
Peak Flowering	Black Turtle Soup	1	1	1	1	219	238	271	243
	Seafarer	1	1	,	1	206	221	244	224
	Mean	1	ı	1	1	212	229	258	233b
Pod Filling	Black Turtle Soup	435	491	445	457	426	541	510	492
,	Seafarer	409	514	463	462	485	536	499	202
	Mean	422	503	454	459c	455	538	504	499c
Maturity	Black Turtle Soun	341	438	458	412	29	51	88	26
To Tanana	Seafarer	380	428	434	414	64	53	106	74
	Mean	361	433	446	413b	46	52	9.7	65a
Mean: daylength		391x	468y	450y		238x	273y	286y	
Mean	BTS		43	434 m			26	264 m	
cultivar	Seafarer		43	439 m			26	268 m	
Interactions									
1) Dev. stage x cultivar	cultivar		u.	n.s.				*	
	davlength		*	**				**	
3) Cultivar x daylength	ylength		'n.	n.s.				*	
4) Dev. stage x cultivar	cultivar								
x daylength			ď	n.s.				n.s.	

 $^{4}$ Means within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.



## APPENDIX C

GROWTH AND POD ABSCISSION AS AFFECTED BY

CARBON DIOXIDE LEVELS




Table C-1

Effects of Carbon Dioxide Levels on Number of Flowers, Number of Pods and Percentage of Pod Abscission at 3 Stages of Development of Two Dry Bean Cultivars Grown in  $1975^{\rm d}$ 

Cultivars  CO1 Its  Black Turtle Soup 9 11 Seaferer 124 12 Mean 125 12 Seaferer 13 12 Seaferer 13 12 Seaferer 13 12 Seaferer 125 12 Seaferer 125 12 Seaferer 125 12 Seaferer 125 125 125 125 125 125 125 125 125 125			No.	No. of Flowers	wers	ž	No. of Pods	spo	8 PC	% Pod Abscission	ssion
Select Furtle Soup   9   11   10   10   10   10   10   10	Dev. Stages	Cultivars	CO2	Level	(mdd)	CO2	Level	(mdd)	00	2 Level	(mdd)
Black Turtle Soup   2   11   10   12   12   13   14   15   14   15   14   15   14   15   14   15   14   15   14   15   14   15   14   15   14   15   14   15   14   15   14   15   14   15   14   15   14   15   14   15   14   15   14   15   15			300	800	Mean	300	800	Mean	300	800	Mean
Seafarer   24   24   19   22   20	Peak Flowering	Black Turtle Soup	6	11	10	8	80	80	ı		
Black Turtle Soup 8 13 11 6 13 10 24.8     Seafarer		Seafarer Mean	24 16	24	24 17a	14	22 15	14c	1 1	1 1	
	Pod Filling	Black Turtle Soup	8	13	11	9	13	10	24.8	4.0	14.4
Black Turtle Soup   10   12   11   5   11   9   44.1   14.2   15   15   15   15   15   15   15   1		Seafarer Mean	20	21 17	21 16a	11	14	112	34.3	27.6	35.7 25.1a
	Maturity	Black Turtle Soup	10	175	11	ı,	#:	80 (	44.1	14.6	29.3
Mean   15x   17y   9x   14y   44.2y     Ultivar   BPS   10 m   9 m     Seafarer   22 n   14 n     Pactions   22 n   14 n     Pactions   22 n   14 n     Pactions   22 n   22 n     Pactions   22 n     Paction		Seafarer Mean	16	18	20 16a	o n	13	6 6 I	54.0	28.2	41.1b
Seafarer   22 n   14 n   9 m   9 m   9 m   9 m   9 m   14 n   1		Mean	15x	17y		×6	14y		44.2y	22.0x	
Seafarer   22 n   14 n	Mean	BTS		10 m						21.9 ш	
e x collitivar	cultivar	Seafarer		22 n			14 n			44.3 n	
bev. steps x CO <sub>2</sub> level n.s. ** Cultivar x CO <sub>2</sub> level n.s. n.s. n.s. n.s. v. n. level x cultivar n.s. n.s. n.s. n.s.	Interactions	cultivar		*			*			n.s.	
Cultivar x CO2 level n.s. n.s. n.s. n.s. n.s. n.s. n.s. n.s		COn level		n.s.			*			n.s.	
Dev. stage x cultivar		02 level		n.s.			n.s.			n.s.	
		cultivar		n.s.			n.s.			n.s.	

Ameans within a given parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.

Table C-2

Effects of Carbon Dioxide Levels on Number of Flowers, Number of Pods and Percentage of Pod Abscission at 3 Stages of Development of Two Dry Bean Cultivars Grown in 1976<sup>a</sup>

		No	No. of Flowers	owers	Ň	No. of Pods	spo	% PC	<pre>% Pod Abscission</pre>	ssion
Dev. Stages	Cultivar	002	CO <sub>2</sub> Level (ppm)	(mdd)	C02	CO2 Level (ppm)	(wdd	COS	CO2 Level (ppm)	(mdd)
		300	800	Mean	300	800	Mean	300	800	Mean
Peak Flowering	Black Turtle Soup	2	6	7	2	ω	7	1	1	
	Seafarer Mean	15	20	18 12a(*)	12 8	13	지기	ı	,	
Pod Filling	Black Turtle Soup	Ŋ	6	7	4	6	9	15.5	5.4	10.4
	Seafarer Mean	15	22 16	18 13a	919	15	10 8a	37.5	33.2	28.4a
Maturity	Black Turtle Soup	9	16	7	4 /	10	7	36.7	24.4	30.5
	Seafarer Mean	11	20	15b	øω	101	7a	55.5	39.1	47.3b
	Mean	10x	17y		×9	113		46.5y	29.2x	
Mean	BTS		8			ш 9			20.5 m	
cultivar	Seafarer		19 n			11 n			55.2 n	
Interactions										
remitting a court man ()	4000		n.s.			*			n.s.	
2) Dev. stage x	Curcion		n.s.			n.s.			n.s.	
	002 level		n.s.			n.s.			*	
4) Dev. stage x cultivar	cultivar		n.s.			n.s.			n.s.	

<sup>a</sup>Means within a given parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.

=




Table C-3

Effects of Carbon Dioxide Levels on Root Weight Per Plant, Stem Weight Per Plant and Total Plant Weight at 3 Stages of Development of Two Dry Bean Cultivars Grown in 1975<sup>a</sup>

		Root	Root Wt/Plant (g)	nt (g)	Stem	Stem Wt/Plant (g)	ıt (g)	Tota]	Total Plant Wt (g)	Wt (g)
Dev. Stage	Cultivar	C02	CO <sub>2</sub> Level (ppm)	(mdd)	002	CO2 Level (ppm)	(mdd)	8	CO <sub>2</sub> Level (ppm)	(mdd)
		300	800	Mean	300	800	Mean	300	800	Mean
Peak Flowering	Black Turtle Soup	1.3	1.7	1.5	1.6	2.0	1.8	6.4	9.0	7.7
	Seafarer	1.3	1.5	1.4	1.6	1.9	1.8	8.9	8.3	7.6
	Mean	1.3	1.6	1.4b	1.6	1.9	1.8b	9.9	9.8	7.6a
Pod Filling	Black Turtle Soup	6.0	1.5	1.2	1.4	2.8	2.1	7.8	15.4	11.6
	Seafarer	6.0	1.2	1.0	1.5	1.8	1.7	8.2	12.5	10.4
	Mean	6.0	1.3	1.1a	1.4	2.3	1.9b	8.0	14.0	11.0b
Maturity	Black Turtle Soup	0.7	1.5	1.1	6.0	2.1	1.5	8.9	15.7	11.2
	Seafarer	6.0	1.6	1.3	1.0	5.0	1.5	5.5	13.8	7.6
	Mean	0.8	1.6	1.2a	0.0	2.1	1.5a	6.2	14.8	10.5b
	Mean	1.0x	1.5y		1.3x	2.1y		x6.9	12.5y	
Mean	BIS		1.3 m			1.8 n			10.2 n	
cultivar	Seafarer		1.2 ш			1.6 m			9.2 ш	
Interactions										
1) Dev. stage x cultivars	cultivars		*			*			n.s.	
	CO3 level		*			*			*	
3) Cultivar x CO2 level	302 level		*			*			*	
	k čultivar									
x CO <sub>2</sub> level			n.s.			*			n.s.	
a Mo	age on thin the came narameter followed by similar letters are not significantly different at	eter fol	y bewo	v similar	letters	are no	t signifi	cantly di	fferent	at

Means within the same parameter followed by sli the 1% level by Duncan's Multiple Range Test.



Table C-4

Effect of Carbon Dioxide Levels on Root Weight Per Plant, Stem Weight Per Plant and Total Plant Weight at 3 Stages of Development of Two Dry Bean Cultivars Grown in  $1976^a$ 

		Root	Root Wt/Plant (g)	nt (g)	Stem	Stem Wt/Plant (g)	nt (g)	Total	Total Plant Wt (g)	Wt (g)
Dev. Stage	Cultivars	002	CO <sub>2</sub> Level (ppm)	(mdd)	CO2	CO2 Level (ppm)	(mdd)	8	CO2 Level (ppm)	(mdd)
		300	800	Mean	300	800	Mean	300	800	Mean
Peak Flowering	Black Turtle Soup Seafarer Mean	0.10	1.4	1.2 0.8 1.0a	0.0	1.6	1.2 0.7 1.0a	3.2	3.9	5.5 3.1 4.3a
Pod Filling	Black Turtle Soup Seafarer Mean	1.2	1.4	1.7 1.1 1.4b	0.8	2.3	1.8 1.6b	6.4 5.2	8 8 8 6 0 8	7.6 6.0 6.8b
Maturity	Black Turtle Soup Seafarer Mean	1.2	1.9	2.0 1.3 1.6c	0.9	2.9	1.8 1.9c	7.4 5.3 6.4	14.0 11.0 12.7	10.9 8.2 9.5c
	Mean	x6.0	1.7y		×6.0	2.0y		4.9x	8.9y	
Mean	BTS		1.6 n			1.6 n			8.0 n	
curtivar	Seafarer		1.1 m	u		1.4 m			5.8 m	
Interactions  1) Dev. stage x cultivars 2) Dev. stage x CO2 level 3) Cultivar x CO2 level 4) Dev. stage x cultivar x CO2 level	k cultivars k CO <sub>2</sub> level CO <sub>2</sub> level k cultivar		* * * * * * * * * * * * * * * * * * *			*** u u			n * n .s.	

<sup>a</sup>Means within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.



Table C-5

Effects of Carbon Dioxide Levels on Leaf Area Per Plant, Number of Leaf Per Plant and Leaf Weight Per Plant at 3 Stages of Development of Two Dry Bean Cultivars Grown in  $1975^a$ 

		Leaf A	ea/Pla	Leaf Area/Plant (cm <sup>2</sup> )	Number	of Lea	Number of Leaf/Plant	Leal	Leaf Wt/Plant (g)	nt (g)
Dev. Stage	Cultivar	CO2	CO2 Level (ppm)	(mdd)	CO2	CO <sub>2</sub> Level (ppm)	(mdd)	8	CO2 Level (ppm)	(mdd)
		300	800	Mean	300	800	Mean	300	800	Mean
Peak Flowering	Black Turtle Soup	715	802	758	12	24	18	2.9	4.5	3.7
	Seafarer Mean	710	740	725 742a	14	13	14 16b	2.0	4.0	3.2a
Pod Filling	Black Turtle Soup	620	1058	839	10	15	12	1.9	5.5	3.7
	Seafarer Mean	708	738 898	673 756a	11	14	13a	1.9	4.4	3.2a
	Mean	663x	835y		12×	16y		2.2x	4.2y	
Mean	BTS		n 66L			15 m			3.7 n	
cultivar	Seafarer		ш 669	E		13 ш			2.7 m	
Interactions										
	x cultivar		n.s.			* ;			***	
2) Dev. stage x CO <sub>2</sub> level	x CO <sub>2</sub> level		* *			*	o."		*	
	x cultivar		n.s.			*			*	

Ameans within the same parameter followed by similar letters are not significantly different at the 1% level by Dancan's Multiple Range Test.



Table C-6

Effects of Carbon Dioxide Levels Leaf Area Per Plant, Number of Leaf Per Plant and Leaf Weight Per Plant at 3 Stages of Development of Two Dry Bean Cultivars Grown in  $1976^a$ 

		Leaf Ar	ea/Pla	Leaf Area/Plant (cm <sup>2</sup> )	No.	No. Leaf/Plant	Lant	Leaf	Leaf Wt/Plant (g)	nt (g)
Dev. Stage	Cultivar	C02	CO2 Level (ppm)	(mdd)	CO <sub>2</sub>	CO <sub>2</sub> Level (ppm)	(mdd)	co	CO2 Level (ppm)	(wdd)
		300	800	Mean	300	800	Mean	300	800	Mean
Peak Flowering	Black Turtle Soup Seafarer	418	834	626 471	0101	12	##	2.0	4.1	3.0
	Mean	370	727	549a	10	12	11a	1.6	3.1	2.4a
Pod Filling	Black Turtle Soup	423	1252	837	11	17	14	3.1	8.4	8.
	Seafarer Mean	363	825 1039	594 716b	21	13	13b	2.5	6.4	4.4p
	Mean	381x	883y		10×	13y		2.1x	4.7y	
Mean	BTS		732 n			12 n			4.4 n	
cultivar	Seafarer		533 m			11 m			2.4 m	
Interactions										
1) Dev. stage x cultivar	x cultivar		n.s.			* :			* *	
<ol> <li>Dev. stage x CO<sub>2</sub> level</li> <li>Cultivar x CO<sub>2</sub> level</li> </ol>	x CO <sub>2</sub> level CO, level		* *			*			*	
4) Dev. stage x cultivar x CO <sub>2</sub> level	x ćultivar		n.s.			n.s.			:	

Ameans within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.



## APPENDIX D

RELATIONSHIP OF POD ABSCISSION TO DIFFERENTIAL LEVELS

OF CARBOHYDRATES PRODUCED BY VARIATION IN

CARBON DIOXIDE CONCENTRATION




Table D-1

Effects of Stages of Pod Development and Carbon Dioxide Levels on Free Sugar Concentration in Root, Stem and Leaf of the Two Dry Bean Cultivars (mg glucose per g dry weight)

			Root			Stem			Leaf	
Dev. Stage	Cultivar	002	CO2 Level (ppm)	(mdd)	CO2	CO <sub>2</sub> Level (ppm)	(mdd)	COS	CO <sub>2</sub> Level (ppm)	(mdd)
		300	800	Mean	300	800	Mean	300	800	Mean
paironola Veed	Black Turtle Soun	24.7	26.9	25.8	0.69	82.6	75.8	72.3	79.3	75.8
tean troughting	Seafarer	20.6	23.4	22.0	49.7	61.4	55.5	56.9	70.5	63.7
	Mean	22.7	25.1	23.9b	59.3	72.0	65.70	64.6	74.9	69.7a
pod pilling	Black Thrtle Sonn	19.8	26.4	23.1	63.4	64.3	63.9	61.8	92.2	77.0
Surrey pos	Conference	13.1	16.4	14.7	36.2	61.0	48.6	44.7	94.3	69.5
	Mean	16.4	21.4	18.9a	49.8	62.7	56.2b	53.3	93.2	73.2a
Maturity	Black Turtle Soup	22.0	25.6	23.8	21.3	73.5		1	1	,
Hacutter	Conform	13.4	22.6	18.0	16.2	44.2		1	1	,
	Mean	17.7	24.1	20.9a	18.7	58.8	38.8a		1	
Mean: CO <sub>2</sub> Level	1	18.9x	23.5y		42.6x	64.5y	7	58.9x	84.1y	
Mean	BTS		24.2 n			62.3 n			76.4 n	
cultivar	Seafarer		18.2 ш			44.8 1	ш		ш 9.99	
Interactions										
Dev.	stage x cultivar		n.s.			n.s.			n.s.	
2) Dev. stage x CO, level	CO2 level		n.s.			*			k ·	
	O2 level		n.s.			n.s.			* *	
4) Dev. stage x cultivar x CO <sub>2</sub> level	cultivar		n.s.			*			n.s.	

Ameans within the same parameter followed by similar letters are not significantly different at the 18 level by Duncan's Multiple Range Test.




Table D-2

Effects of Stages of Pod Development and Carbon Dioxide Levels on Free Sugar Concentration in Seed and Pod Wall of the Two Dry Bean Cultivars (mg glucose per g dry Weight) $^{\rm A}$ 

			Seed			Pod Wall	
Dev. Stage	Cultivar	CO2	CO2 Level (ppm)	(wdo	co	CO2 Level (ppm)	(mdd)
		300	800	Mean	300	800	Mean
Peak Flowering	Black Turtle Soup	1		-	226	293	259
	Seafarer	•	1	,	262	291	276
	Mean	1	1	1	244	292	268c
Pod Filling	Black Turtle Soup	89	115	92	108	118	113
5	Seafarer	09	118	68	102	140	121
	Mean	64	116	906	105	129	II la
Maturito	Black Turtle Soup	69	82	9/	20	26	38
- Touris	Seafarer	76	94	85	32 26	4 4 2 4 9 6 9	37 38a
	Medil			1			
Mean: CO, Level		x89	102y		125x	157y	
Mean	BTS		84 m			137 m	
cultivar	Seafarer		87 m			145 m	
Interactions							
	ıltivar					n.s.	
<ol> <li>Dev. stage x CO2 level</li> <li>Cultivar x CO2 level</li> </ol>	Jevel Jevel		n.s.			n.s.	
	ultivar		*			*	

 $^{8}$ Means within the same parameter followed by similar letters are not significantly different at the 1\$ level by Duncan's Multiple Range Test.




Table D-3

Effects of Stages of Pod Development and Carbon Dioxide Levels on Starch Concentration in Root, Stem and Leaf of the Two Dry Bean Cultivars (mg glucose per g dry weight)  $^{\rm a}$ 

			Root			Stem			Leaf	
Dev. Stage	Cultivar	C02	CO2 Level (ppm)	(mdd	CO2	CO <sub>2</sub> Level (ppm)	(mdd	CO2	CO2 Level (ppm)	(wdd)
		300	800	Mean	300	800	Mean	300	800	Mean
painono la deca	Black Turtle Soun	10.1	34.9	22.5	304	298	301	304	540	422
rear Fromering	Seafarer	13.8	27.3	20.6	91	82	87	146	430	288
	Mean	12.0	31.1	21.5b	198	190	194b	225	482	3550
Dod Filling	Black Turtle Soup	28.9	41.2	35.1	214	321	268	69	457	263
Four Fitting	Seafarer	15.6	26.2	20.9	65	113	89	38	260	149
	Mean	22.2	33.7	28.0c	140	217	178b	23	358	206a
Mathematical	Black Turtle Soup	0.3	9.4	4.8	28	170	66	,	1	1
Hacutter	Conform	8	5.4	4.6	33	37	35	1	ı	ı
	Mean	2.0	7.4	4.7a	31	103	67a	,		
Mean: CO, Level		12.1x	24.17		123x	170y		139×	422y	
Mean	BTS		20.8 n			223 n			342 n	
cultivar	Seafarer		15.3 m			70 m			218 1	E
Interactions										
10 00040	7621+1:00		*			*			n.s.	
	Curreyar		*			*			*	
3) Cultivar x CO2 level	0, level		n.s.			*			*	
	cultivar		n.s.			*			*	

 $^{a}$ Means within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.




Table D-4

Effects of Stages of Pod Development and Carbon Dioxide Levels on Starch Concentration in Seed and Pod Wall of the Two Dry Bean Cultivars (mg glucose per g dry weight)  $^{\rm a}$ 

			Seed			Pod Wall	1	
Dev. Stage	Cultivar	002	CO <sub>2</sub> Level (ppm)	(wdd	CO2	CO2 Level (ppm)	(wdd)	
		300	800	Mean	300	800	Mean	
Series Contracts	and offer your	,	,		200	117	158	
rean Fromering	Seafarer	1	1	1	201	84	142	
	Mean	1	,	1	201	100	150b	
Seillin Sou	died Sour	399	271	335	403	416	409	
For Fitting	Soafarer	411	312	361	398	408	403	
	Mean	405	292	348c	401	412	406c	
	dios elarin de la	285	275	280	32	09	46	
Maturity	Southern	265	252	258	27	21	24	
	Mean	276	263	269b	29	40	35a	
Mean: CO <sub>2</sub> Level		340y	277×		210y	184×		1
Mean	BTS		307 m			205 n		
cultivar	Seafarer		209 ш			190 ш	1	
Interactions								
1	3.0		*			n.s.		
	urcivar 10 Jane		**			*		
2) Dev. stage X CO2 Level	Tever		n.s.			*		
	in1+ivar							
			n.s.			n.s.		
7-2								1

<sup>a</sup>Means within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.




Table D-5

Effects of Stages of Pod Development and Carbon Dioxide Levels on Total Nonstructural Carbohydrate Concentration in Root, Stem and Leaf of the Two Dry Bean Cultivars (mg glucose per g dry weight) a

			Root			Stem			Leaf	
Dev. Stage	Cultivar	00	CO <sub>2</sub> Level (ppm)	(mdd	CO2	CO2 Level (ppm)	(mdd)	00	CO <sub>2</sub> Level (ppm)	(wdd)
		300	800	Mean	300	800	Mean	300	800	Mean
Peak Flowering	Black Turtle Soup	30	61	46	373	380	377	378	614	496
	Seafarer	24	44	34	141	153	147	206	499	352
	Mean	27	23	40c	257	267	262b	292	226	424b
Pod Filling	Black Turtle Soup	48	67	28	286	386	336	133	545	339
	Seafarer	30	42	36	102	194	148	85	347	216
	Mean	39	54	47b	194	290	242b	109	446	278a
Maturity	Black Turtle Soup	21	35	28	49	218	134	,	1	1
To Tanonii	Seafarer	17	29	23	20	84	49	1	1	1
	Mean	13	32	25a	49	151	100a			
Mean: CO <sub>2</sub> Level	1	28x	46y		167×	236Y		200x	501y	
Mean	BTS		44 n			282 n			418 n	
cultivar	Seafarer		31 ш			121 m			284 п	ш
Interactions										
The sample of the	Trans.		*			*			n.s.	
2) Dev. stade x	Correspond		*			*			*	
	02 level		n.s.			*			*	
4) Dev. stage x cultivar x CO level	cultivar		n.s.			*			*	

Ameans within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Multiple Range Test.





Table D-6


Effects of Stages of Pod Development and Carbon Dioxide Levels on Total Monstructural Carbohydrate Concentration in Seed and Pod Wall of the Two Dry Bean Cultivars (mg glucose per g dry weight) <sup>a</sup>

			Seed			Pod Wall	-	
Dev. Stage	Cultivar	8	CO2 Level (ppm)	(wdc	00	CO2 Level (ppm)	(mdd)	
		300	800	Mean	300	800	Mean	
Peak Flowering	Black Turtle Soup	,			433	406	420	
	Seafarer	1	,	1	429	372	401	
	Mean	1		1	431	389	410b	
Pod Filling	Black Turtle Soup	450	363	407	515	532	524	
	Seafarer	457	412	434	501	542	522	
	Mean	453	387	420b	208	537	5230	
Maturity	Black Turtle Soup	355	371	363	52	112	82	
-	Seafarer	328	342	335	29	91	75	
	Mean	341	357	349a	55	101	78a	
Mean: CO <sub>2</sub> Level		39.7⊻	372×		331x	343x		
Mean	BIS		385 ш	-		342 m		
cultivar	Seafarer		384 m			332 ш		
Interactions								
1) Dem stade v cultivar	l i war		*			n.s.		
2) Dev. stage x C	Con level		*			*		
3) Cultivar x CO, level	level		n.s.			n.s.		
	Sultivar		E.			S		
x CO <sub>2</sub> Tevel								

Ameans within the same parameter followed by similar letters are not significantly different at the 1% level by Duncan's Wultiple Range Test.









## SELECTED BIBLIOGRAPHY

- Abbott, D. L. 1960. The bourse shoot as a factor in the growth of apple fruits. Ann. Appl. Biol. 48: 434-438.
- Abeles, F. B. 1967. Mechanism of action of abscission accelerators. Physiol. Plant. 20: 442-454.
- Abeles, F. B., L. E. Craker and G. R. Leather. 1971. Abscission: The phytogerontological effects of ethylene. Plant Physiol. 47: 7-9.
- Abeles, F. B. and H. E. Gahagan. 1968. Abscission: The role of ethylene, ethylene analogues, carbon dioxide and oxygen. Plant Physiol. 43: 1255-1258.
- Adams, M. W. 1967. Basic of yield compensation in crop plants with special reference to the field bean, Phaseolus vulgaris L. Crop Sci. 7:505-510.
- Addicott, F. T. 1965. Physiology of abscission. In Encyclopedia of Plant Physiology, XV (Part 2); 1094-1126. Ed. W. Ruhland. Springer-Verlag, Berlin.
- Addicott, F. T. 1970. Plant hormones in the control of abscission. Biol. Rev. 45: 485-524.
- Addicott, F. T., H. R. Carns, J. L. Lyon, O. E. Smith and J. S. McMeans. 1964. On the physiology of abscisins. In Regulateurs naturels de la croissance végétale, 687-703. Ed. J. P. Nitsch, CNRS. Paris.
- Addicott, F. T. and J. L. Lyon. 1969. Physiology of abscisic acid and related substances. Ann. Rev. Plant Physiol. 20: 139-164.
- Addicott, F. T. and J. L. Lyon. 1973. Physiological ecology of abscission. In Shedding of Plant Parts, 85-124. Ed. T. T. Kozlowski, Academic Press.




- Addicott, F. T., J. L. Lyon, K. Ohkuma, W. E. Thiessen, H. R. Carns, O. E. Smith, J. W. Cornforth, B. V. Milborrow, G. Ryback and P. F. Wareing. 1968. Abscisic acid: A new name for abscisin II (Dormin). Science 159: 1943.
- Aung, L. H., F. Tognoni and A. A. De Hertogh. 1973. Changes in the carbohydrates of tulip bulbs during development. Hort. Sci. 8: 207-208.
- Batjer, L. P. 1965. Fruit thinning with chemicals. Agr. Information Bull. No. 289. U.S. Dept. Agr., Agr. Res. Serv.
- Binkley, A. M. 1932. The amount of blossom and pod drop on six varieties of garden beans. Proc. Amer. Soc. Hort. Sci. 29: 489-492.
- Blackwall, F. L. C. 1969. Effects of weather, irrigation, and pod removal on the setting of pods and the marketable yield of runner beans <a href="Phaseolus multi-florus">Phaseolus multi-florus</a>. J. Hort. Sci. 44: 371-384.
- Bornman, C. H. 1969. Laminal abscission in <u>Streptocarpus</u>.


  Abstracts. XI<sup>th</sup> Int. Bot. Congress. <u>Seattle</u>
  U.S.A. p. 18.
- Brown, K. J. 1971. Plant density and yield of cotton in northern Nigeria. Cotton Growing Rev. 48: 255-266.
- Brown, R. H. and R. E. Blaser. 1965. Relationships between reserve carbohydrate accumulation and growth rate in Orchardgrass and Tall Fescue. Crop Sci. 5: 577-582.
- Bukovac, M. J., F. Zucconi, R. P. Larsen and C. D. Kesner. 1969. Chemical promotion of fruit abscission in cherries and plums with special reference to 2-Chloroethanephosphonic acid. J. Amer. Soc. Hort. Sci. 94: 226-230.
- Carns, H. R. 1966. Abscission and its control. Ann. Rev. Plant Physiol. 17: 295-314.
- Cartter, J. L. and E. E. Hartwig. 1962. The management of soybeans. Adv. in Agron. 14: 360-412.
- Chandler, W. H. 1951. Deciduous Orchards. Lea and Febiger Publ., Philadelphia.




- Cooper, R. L. and W. A. Brun. 1967. Response of soybeans to a carbon dioxide enriched atmosphere. Crop Sci. 7: 455-457.
- Cooper, W. C. and W. H. Henry. 1967. The acceleration of abscission and colouring of Citrus fruit. Proc. Fla. St. Hort. Soc. 80: 7-14.
- Cordner, H. B. 1933. External and internal factors affecting blossom drop and set of pods in lima beans. Proc. Amer. Soc. Hort. Sci. 30: 571-576.
- Coyne, D. P. 1970. Genetic control of a photoperiod-temperature response for time of flowering in beans (Phaseolus vulgaris L.). Crop Sci. 10: 246-248.
- Coyne, D. P. and R. H. Mattson. 1964. Inheritance of time of flowering and length of blooming period in <u>Phaseolus vulgaris</u> L. Proc. Amer. Soc. Hort. Sci. 85: 366-373.
- Cracker, L. E. and F. B. Abeles. 1969. Abscission: Role of abscisic acid. Plant Physiol. 44: 1144-1149.
- Davis, L. A. and F. T. Addicott. 1972. Abscisic acid: Correlations with abscission and with development in the cotton fruit. Plant Physiol. 49: 644-668.
- Dela Fuente, R. K. and A. C. Leopold. 1968. Senescence processes in leaf abscission. Plant Physiol. 43: 1496-1502.
- Dickson, M. H. 1967. Diallel analysis of seven economic characters in snap beans. Crop Sci. 7: 121-124.
- Dornhoff, G. M. and R. M. Shibles. 1970. Varietal differences in net photosynthesis of soybean leaves. Crop Sci. 10: 42-45.
- Dreywood, R. 1946. Industr. Engng. Chem. Anal. ed. 18: 499.
- Dunlap, A. A. 1943. Low light intensity and cotton boll shedding. Science 98: 568-569.
- Eagles, C. F. and P. F. Wareing. 1964. The role of growth substances in the regulation of bud dormancy. Physiol. Plant. 17: 697-709.



- Eaton, F. M. and D. R. Ergle. 1953. Relationship of seasonal trends in carbohydrate and nitrogen levels and effects of girdling and spraying with sucrose and urea to the nutritional interpretation of boll shedding in cotton. Plant Physiol. 28: 503-520.
- Eaton, F. M. and D. R. Ergle. 1954. Effects of shade and partial defoliation on carbohydrate levels and the growth, fruiting and fiber properties of cotton plants. Plant Physiol. 29: 39-49.
- Eaton, F. M. and N. E. Rigler. 1945. Effects of light intensity, nitrogen supply and fruiting on carbohydrate utilization by the cotton plant. Plant Physiol. 20: 380-411.
- Edgerton, L. J., C. G. Forshey and G. D. Blanpied. 1967. Effect of summer applications of Alar (B 995) in reducing drop and improving color and firmness of apples. Proc. New York State Hort. Soc. 112: 204-206.
- Edgerton, L. T. and W. J. Greenhalgh. 1969. Regulation of growth, flowering and fruit abscission with 2-Chloroethanephosphonic acid. J. Amer. Soc. Hort. Sci. 94: 11-13.
- El-Antably, H. M. M., P. F. Wareing and J. Willman. 1967. Some physiological responses to d,l-abscisin (dormin). Planta. 73: 74-90.
- Evans, L. T. and R. W. King. 1974. Factors affecting flowering and reproduction in grain legume. TAC/FAO. Working Party on Grain Legumes, New Delhi, 1974.
- Ewing, E. C. 1918. A study of certain environmental factors and varietal differences influencing the fruiting of cotton. Mississippi Agr. Exp. Sta. Tech. Bull. 8: 1-95.
- Fisher, V. J. and C. K. Weaver. 1974. Flowering, pod set, and pod retention of lima bean in response to night temperature, humidity and soil moisture. J. Amer. Soc. Hort. Sci. 99: 448-450.
- Flinn, A. M. and J. S. Pate. 1970. A quantitative study of carbon transfer from pod and subtending leaf to the ripening seeds of the field pea (Pisum arvense L.). J. Expt. Bot. 21: 71-82.




- Ford, M. A. and G. N. Thorne. 1967. Effect of CO<sub>2</sub> concentration on growth of sugar beet, barley, kale and maize. Ann. Bot. N.S. 31: 629-644.
- Fukui, J., O. Mutsuo and I. Watanabe. 1965. Studies on the seed production of soybean. I. Effect of temperature on photosynthesis of soybean. Proc. Crop Sci. Soc. Japan 33: 432-436.
- Ghorashy, S. R., J. W. Pendleton, R. L. Bernard, and M. E. Bauer. 1971. Effect of leaf pubescence on transpiration, photosynthetic rate, and seed yield of three near-isogenic lines of soybeans. Crop Sci. 11: 426-427.
- Gifford, R. M., P. M. Bremner and D. B. Jones. 1973. Assessing photosynthetic limitation to grain yield in a field crop. Aust. J. Agric. Res. 24: 297-307.
- Goodman, A. 1955. Correlation between cloud shade and shedding in cotton. Nature 176: 39.
- Greenwood, E. A. N., P. Farrington, and J. D. Beresford.
  1975. Characteristics of the canopy, root system and grain yield of a crop of <u>Lupinus angustifolius</u> cv. Unicrop. Aust. J. Agric. Res. 26: 497-510.
- Griggs, W. H., B. T. Iwakiri, R. B. Fridley and J. Mehlaschau. 1970. Effect of 2-Chloroethylphosphonic acid and cycloheximide on abscission and ripening of "Bartlett" pears. Hort. Science 5: 264-266.
- Guinn, G. 1974. Abscission of cotton floral buds and bolls as influenced by factors affecting photosynthesis and respiration. Crop Sci. 14: 291-293.
- Hall, W. C., F. A. Herrero and F. R. H. Katterman. 1961. Leaf abscission in cotton. IV Effects of a natural promoter and amino acids on abscission in cotyledonary node explants. Botan. Gaz. 123: 29-34.
- Hardman, L. L. and W. A. Brun. 1971. Effect of atmospheric carbon dioxide enrichment at different developmental stages on growth and yield components of soybeans. Crop Sci. 11: 886-888.




- Hartmann, H. T. 1955. Induction of abscission of olive fruits by maleic hydrazide. Bot. Gaz. 123: 29-34.
- Hartmann, H. T., A. J. Heslop and J. Whisler. 1968. Chemical induction of fruit abscission in olives. Calif. Agric. 22: 14-16.
- Hartmann, H. T., A. Tombesi and J. Whisler. 1970. Promotion of ethylene evolution and fruit abscission in the olive by 2-Chloroethanephosphonic acid and cycloheximide. J. Amer. Soc. Hort. Sci. 95: 635-640.
- Hartung, W. and J. Witt. 1968. Uber der Einfluss der Bodenfeuchtigkeit auf den Wuchsstoffgehatt von Anastatica hierochuntica und Helianthus Flora (Jena). Abt. B 157: 603-614.
- Hawkins, R. S., R. L. Matlock, and C. Hobart. 1933. Physiological factors affecting the fruiting of cotton with special reference to boll shedding. Arizona Agri. Exp. Sta. Tech. Bull. 46: 361-407.
- Heinicke, A. J. 1917. Factors influencing the abscission of flowers and partially developed fruits of the apple (Pyrus malus). Cornell Univ. Agr. Exp. Sta. Bull. 393: 44-112.
- Heinze, P. H. and A. E. Murneek. 1940. Comparative accuracy and efficiency in determination of carbohydrates in plant material. Missouri Agri. Exp. Sta. Res. Bull. 314.
- Hendricks, S. B. and H. A. Borthwick. 1965. The physiological functions of phytochrome. In "Chemistry and Biochemistry of Plant Pigments." 405-436. Ed. T. W. Goodwin, Academic Press, New York.
- Hicks, D. R. and J. W. Pendleton. 1969. Effect of floral bud removal on performance of soybeans. Crop Sci. 9: 435-437.
- Hodge, J. E. and B. T. Hofrieter. 1962. Determination of reducing sugars and carbohydrates. In "Methods in Carbohydrate Chemistry." Vol. I, pp. 380-394. Ed. R. L. Whistler and M. L. Wolfrom. Academic Press, New York.
- Hodgson, G. L. and H. E. Blackman. 1956. An analysis of the influence of plant density on the growth of <u>Vicia faba</u>. I. The influence of density on the pattern of development. J. Expt. Bot. 7: 147-165.

- Itai, C. and Y. Vaadia. 1971. Cytokinin activity in water-stressed shoots. Plant Physiol. 47: 87-90.
- Jacobs, W. P. 1962. Longevity of plant organs: Internal factors controlling abscission. Ann. Rev. Plant Physiol. 13: 403-436.
- Jacobs, W. P. 1968. Hormonal regulation of leaf abscission. Plant Physiol. 43: 1480-1495.
- Jacobs, W. P., J. A. Shield, Jr., D. J. Osborne. 1962.
  Senescence factor and abscission of Coleus leaves.
  Plant Physiol. 37: 104-106.
- Jung, G. A. and D. Smith. 1960. Influence of extended storage at constant low temperature on cold resistance and carbohydrate reserves of alfalfa and medium red clover. Plant Physiol. 35: 123-125.
- Kambal, A. E. 1968. A study of the agronomic characters of some varieties of <u>Vicia faba</u>. Sudan Agric. J. 3: 1-10.
- Kato, I., S. Sakaguchi and Y. Naito. 1955. Anatomical observations on fallen buds, flowers and pods of soy-bean, Glycine Max. M. Tokai-Kinki Nat. Agr. Exp. Sta. Bull. 2: 159-168.
- Kiyosawa, S. and K. Kiyosawa. 1962. Effect of daylength on fruiting in soybean plant. Proc. Crop Sci. Soc. Japan 30: 342-345.
- Koch, R. B., W. F. Geddes and F. Smith. 1951. The carbohydrates of Gramineae. I. The sugars of the flour of wheat (<u>Triticum</u> <u>vulgare</u>) Cereal Chem. 28: 424-430.
- Krenzer, Jr., E. G. and D. N. Moss. 1975. Carbon dioxide enrichment effects upon yield and yield components in wheat. Crop Sci. 15: 71-74.
- Laidlaw, R. A. and S. G. Reid. 1952. Analytical studies on the carbohydrates of grass and clovers. I. Development of methods for the estimation of free sugar and fructosan contents. J. Sci. Fd. Agric. 3: 19-25.
- Lambeth, V. N. 1950. Some factors influencing pod set and yield of the lima bean. Univ. of Missouri Agr. Exp. Sta. Res. Bull. 466.



- Lee, T. A., D. L. Ketring and R. D. Powell. 1972. Flowering and growth response of peanut plants (Arachis hypogara. L.) at two levels of relative humidities. Plant Physiol. 49: 190-193.
- Leuty, S. J. and M. J. Bukovac. 1968. A comparison of the growth and anatomical development of naturally abscising, non-abscising and naphthaleneacetic acid-treated peach fruits. Phytomorphology 18: 372-379.
- Lipe, J. A. and P. W. Morgan. 1973. Ethylene, a regulator of young fruit abscission. Plant Physiol. 51: 949-953.
- Loomis, W. E. 1926. A study of the clearing of alcoholic plant extracts. Plant Physiol. 1: 179-189.
- Luckwill, L. C. 1953. Studies of fruit development in relation to plant hormones. I. Hormone production by the developing apple seed in relation to fruit drop. J. Hort. Sci. 28: 14-24.
- McAlister, D. F. and O. A. Krober. 1958. Response of soybeans to leaf and pod removal. Agron. J. 50: 674-677.
- McMichael, B. L., W. R. Jordan and R. D. Powell. 1972. An effect of water stress on ethylene production by intact cotton petioles. Plant Physiol. 49: 658-660.
- Mann, J. D. and E. G. Jaworski. 1970. Comparison of stresses which may limit soybean yields. Crop Sci. 10: 620-624.
- Martin, G. C. and C. Nishijima. 1972. Levels of endogenous growth regulators in abscising and persisting peach fruits. J. Amer. Soc. Hort. Sci. 97: 561-565.
- Mason, T. G. 1922. Growth and abscission in Sea Island cotton. Ann. Bot. 36: 457-483.
- Meadley, J. T. and G. M. Milbourn. 1971. The growth of vining peas. III. The effect of shading on abscission of flowers and pods. J. Agric. Sci. 77: 103-108.
- Milborrow, B. V. 1974. The chemistry and physiology of abscisic acid. Ann. Rev. Plant Physiol. 25: 259-307.



- Nelson, N. 1944. A photometric adaptation of the Somogyi method for the determination of glucose. J. Biol. Chem. 153: 375-379.
- Nitsch, J. P., C. Pratt, C. Nitsch, and N. J. Shaulis. 1960. Natural growth substances in concord and concord seedless grapes in relation to berry development. Amer. J. Bot. 47: 566-5
- Ohkuma, K., F. T. Addicott, O. E. Smith and W. E. Thiessen. 1965. The structure of abscisin. II. Tetrahedron letters 29: 2529-2535.
- Ojehomon, O. O. 1970. A comparison of the vegetative growth, development and seed yield of three varieties of cowpea, Vigna unguiculata (L.) Walp. J. Agric. Sci. 74: 363-374.
  - Ojehomon, O. O. 1970. Effect of continuous removal of open flowers on the seed yield of two varieties of cowpea, Vigna unguiculata (L.) Walp. J. Agric. Sci. 74: 375-381.
    - Ojehomon, O. O. 1972. Fruit abscission in cowpea, Vigna unguiculata. L. Walp. I. Distribution of I4C-assimilates in the inforescence, and comparative growth of ovaries from persisting and abscising open flowers. J. Expt. Bot. 23: 751-761.
      - Ojehomon, O. O., A. S. Rathjen and D. G. Morgan. 1968.

        Effects of daylength on the morphology and flowering of five determinate varieties of Phaseolus vulgaris. L. J. Agric. Sci. 71: 209-214.
        - Ojehomon, O. O., M. S. Zehni and D. G. Morgan. 1973.
          The effects of photoperiod on flower-bud development in <u>Phaseolus vulgaris</u>. Ann. Bot. 37: 871-884.
          - Ortega, Y. S. 1971. Inheritance of flowering time in French bean (Phaseolus vulgaris). Agronomic Tropical 21: 319-328.
            - Osborne, D. J. 1955. Acceleration of abscission by a factor produced in senescent leaves. Nature 176: 1161-1163.
              - Osborne, D. J. 1968. Defoliation and defoliants. Botan. Gaz. 123: 29-34.


Welson

(CDB-14


- Osborne, D. J. 1973. Internal factors regulating abscission. In Shedding of Plant Parts, 125-148. Ed. T. T. Kozlowski, Academic Press.
  - Palmer, R. L., H. Z. Hield and L. N. Lewis. 1969. Citrus leaf and fruit abscission. Proc. 1st. Int. Citrus Symp., 3: 1135-1143.
  - Parkerson, R. H. and F. W. Whitmore. 1972. A correlation of stem sugars, starch and liquid with wood formation in Eastern White pine. Forest Sci. 18: 178-183.
    - Powell, L. E. and C. Pratt. 1966. Growth promoting substances in the developing fruit of peach (<u>Prunus</u> persica L.). J. Hort. Sci. 41: 331-348.
      - Rasmussen, H. P. and M. J. Bukovac. 1969. A histochemical study of abscission layer formation in the bean. Amer. J. Bot. 56: 69-76.
        - Ratner, A., R. Goren and S. P. Monselise. 1969. Activity of pectin esterase and cellulase in the abscission zone of Citrus leaf explants. Plant Physiol. 44: 1717-1723.
          - Reynolds, J. H. and D. Smity. 1962. Trend of carbohydrate reserves in alfalfa, smooth bromegrass and timothy grown under various cutting schedules. Crop Sci. 2: 333-336.
            - Rowlands, D. G. 1961. Fertility studies in the field bean (Vicia faba L.) II. Inbreeding. Heredity 16: 497-508.
              - Rubinstein, B. and A. C. Leopold. 1962. Effects of amino acids on bean leaf abscission. Plant Physiol. 37: 398-401.
                - Rubinstein, B. and A. C. Leopold. 1964. The nature of leaf abscission. Quarterly Rev. Biol. 39: 356-372.
                  - Schlenk, H. and J. L. Gellerman. 1960. Esterification of fatty acids with diazomethane on a small scale. Anal. Chem. 32: 1412-1414.
                    - Shaffer, P. A. and A. F. Hartmann. 1921. The iodimetric determination of copper and its use in sugar analysis I. J. Biol. Chem. 45: 349-364.



- Shaffer, P. A. and A. F. Hartmann. 1921. The iodimetric determination of copper and its use in sugar analysis II. J. Biol. Chem. 45: 365-390.
- Shaffer, P. A. and M. Somogyi. 1933. Copper-Iodometric reagents for sugar determination. J. Biol. Chem. 100: 695-713.
- Smith, D. 1950. Seasonal fluctuations of root reserves in red colver <u>Trifolium pratense</u> L. Plant Physiol. 25: 702-710.
- Smith, D. 1968. Classification of several native North American grasses as starch or fructosan accumulators in relation to taxonomy. Jour. Brit. Grassl. Soc. 23: 306-309.
  - Smith, D. 1969. Removing and analyzing total nonstructural carbohydrates from plant tissue. Research Report #41. University of Wisconsin, 12 pp.
    - Smith, D. and L. F. Graber. 1948. The influence of top growth removal on the root and vegetative development of biennial sweet clover. J. Am. Soc. Agron. 40: 818-831.
      - Smith, O. E., J. L. Lyon, F. T. Addicott and R. E. Johnson. 1968. Abscission physiology of abscisic acid. In Biochemistry and Physiology of Plant Growth Substances; 1547-1560. Ed. F. Wightman and G. Setterfield. Runge Press, Ottawa.
        - Smith, D., G. M. Paulsen and C. A. Raguse. 1964. Extraction of total available carbohydrates from grass and legume tissue: Plant Physiol. 39: 960-962.
          - Somogyi, M. 1937. A reagent for the copper-iodometric determination of very small amounts of sugar. J. Biol. Chem. 117: 771-776.
            - Soper, M. H. R. 1952. A study of the principal factors affecting the establishment and development of the field bean (<u>Vicia faba</u>). J. Agric. Sci., Camb. 42: 335-346.
              - Sprague, V. G. and J. T. Sullivan. 1950. Reserve carbohydrates in orchardgrass clipped periodically. Plant Physiol. 25: 92-102.



- Sprague, V. G. and J. T. Sullivan. 1952. Reserve carbohydrates in orchard grass cut for hay. Plant Physiol. 28: 304-313.
- Swanson, Jr., B. T., H. F. Wilkins, C. F. Weiser, and I. Klein. 1975. Endogenous ethylene and abscisic acid relative to phytogerontology. Plant Physiol. 55: 370-376.
- Sweet, G. B. 1973. Shedding of reproductive structures in forest trees. In "Shedding of Plant Parts," 341-382. Ed. T. T. Kozlowski, Academic Press, New York.
- Swen, M. A. 1933. Factors affecting flower shedding in soybean. Ph.D. dissertation, University of Illinois, Urbana.
- Treshow, M. 1970. Environment and plant response.
  McGraw-Hill, Publ., New York.
- Tureskaya, R., V. Kefeli, M. Kutacek, K. Vackova, N. Tschumarovski and T. Krupnikova. 1968. Isolation and some physiological properties of natural plant growth inhibitors. Biol. Plantarum. 10: 205-221.
- Van Schaik, P. H. and A. H. Probst. 1958. Effects of some environmental factors on flower production and reproductive efficiency in soybeans. Agron. Jour. 50: 192-197.
- Van Schaik, P. H. and A. H. Probst. 1958. The inheritance of inflorescence type, peduncle length, flowers per node, and percent flower shedding in soybeans. Agron. J. 50: 98-102.
- Van Steveninck, R. F. M. 1957. Influence of pea-mosaic virus on the reproductive capacity of yellow lupins. Bot. Gaz. 119: 63-70.
- Van Steveninck, R. F. M. 1957. Factors affecting the abscission of reproductive organs in yellow lupins (Lupinus luteus L.) I. The effect of different patterns of flower removal. J. Expt. Bot. 8: 373-381.
- Van Steveninck, R. F. M. 1958. Factors affecting the abscission of reproductive organs in yellow lupins (<u>Lupinus luteus L.</u>) II. The effects of growth substances, defoliation, and removal of lateral growth. J. Expt. Bot. 9: 372-383.



- Van Steveninck, R. F. M. 1959. Abscission accelerators in lupins (Lupinus Luteus L.) Nature 183: 1246-1248.
- Van Steveninck, R. F. M. 1959. Factors affecting the abscission of reproductive organs in yellow lupins (Lupinus luteus L.) III. Endogenous growth substances in virus-infected and healthy plants and their effect on abscission. J. Expt. Bot. 10: 367-375.
- Wadleigh, C. H. 1944. Growth status of cotton plant as influenced by the supply of nitrogen. Arkansas Agri. Exp. Sta. Bull. 446. 138 pp.
- Waite, R. and J. Boyd. 1953. The water soluble carbohydrates of grasses. I. Changes occurring during the normal life cycle. J. Sci. Fd. Agric. 4: 197-204.
- Wargo, P. M. 1971. Seasonal changes in carbohydrate levels in roots of sugar maple. U.S. Forest Ser. Res. Pap. N.E. 213: 1-8.
- Webster, B. D. 1973. Anatomical and histochemical changes in leaf abscission. In Shedding of Plant Parts 45-84 Ed. T. T. Kozlowski, Academic Press.
- Weinmann, H. 1947. Determination of total available carbohydrates in plants. Plant Physiol. 22: 279-290.
- Wien, H. C., S. L. Altschuler, J. L. Ozbun, and D. H. Wallace. 1976. 14C-assimilate distribution in Phaseolus vulgaris L. during the reproductive period. J. Amer. Soc. Hort. Sci. 101: 510-513.
- Wittenbach, V. A. and M. J. Bukovac. 1974. Cherry fruit abscission. Evidence for time of initiation and involvement of ethylene. Plant Physiol. 54: 494-498.
- Wittwer, S. H. 1954. Control of flowering and fruit setting by plant regulators. In "Plant Regulators in Agriculture," 62-80. Ed. H. B. Tukey, Wiley, Publ., New York.
- Wittwer, S. H. and W. M. Robb. 1964. Carbon dioxide enrichment of greenhouse atmospheres for food crop production. Econ. Bot. 18: 34-56.

Van Branch . . . . naV

- Wright, S. T. C. 1956. Studies of fruit development on relation to plant hormones. III Auxins in relation to fruit morphogenesis and fruit drop in the black currant Ribes nigrum. J. Hort. Sci. 31: 196-211.
- Wright, R. D. and L. H. Aung. 1975. Carbohydrates in two rhododendron cultivars. J. Amer. Soc. Hort. Sci. 100: 527-529.
- Wright, S. T. C. and R. W. P. Hiron. 1970. The accumulation of ABA in plants during wilting and under other stress conditions. In "Plant Growth Substances," 291-298. Ed. D. J. Carr. Springer Verlag, Publ., New York.
- Yamaguchi, S. 1954. Some interrelations of oxygen, carbon dioxide, sucrose and ethylene in abscission. Ph.D. dissertation, University of California, Los Angeles.
- Yemm, E. W. 1935. The respiration of barley plants. I Methods for the determination of carbohydrates in leaves. Proc. Roy. Soc. B. 117: 483-504.
- Yemm, E. W. and A. J. Willis. 1954. Stomatal movements and changes of carbohydrate in leaves of <u>Chrysan-</u> themum maximum. New Phytol. 53: 373-396.
- Yoshida, S., J. H. Cock and F. T. Parao. 1972. Physiological aspects of high yields. In "Rice Breeding," IRRI, Los Banos, pp. 455-468.
- Zeevaart, J. A. D. 1971. Effects of photoperiod on growth rate and endogenous gibberellins in the long-day rosette plant spinach. Plant Physiol. 47: 821-827.
- Zeevaart, J. A. D. 1971. (+)-Abscisic acid content of spinach in relation to photoperiod and water stress. Plant Physiol. 48: 86-90.
- Zucconi, F. 1975. Reassessment of the relationship between hormonal and developmental changes during abscission with particular reference to peach (Prunus persica L. Batsch) fruit. Ph.D. dissertation, Michigan State University.
- Zucconi, F., R. Stosser and M. J. Bukovac. 1969. Promotion of fruit abscission with abscisic acid. BioScience, 19: 815-817.

