RARY

-----

#### ABSTRACT

#### THE RELATIONSHIP OF SOME LINEAR AND PHYSICAL MEASUREMENTS

### TO BEEF CARCASS COMPOSITION

by Deloran M. Allen

Eighty steer carcasses were selected for chilled carcass weight and 12th rib fat thickness (average of three measurements). Two weight groups (500 to 550 lb. and 700 to 750 lb.) of 40 carcasses each and four fat thickness ranges of 10 carcasses each (0.26 to 0.50 in., 0.51 to 0.75 in., 0.76 to 1.00 in. and 1.01 to 1.25 in.) were selected within each weight group. The carcasses were subjectively scored for each grade factor and some linear measurements of fat and muscle were recorded. External fat thickness probes were made 4, 8 and 12 in. off the dorsal midline of the left side of each carcass, perpendicular to the anterior edge of the 5th, 8th and 11th thoracic vertebrae, the 1st, 4th and 6th lumbar vertebrae and the 3rd and 5th sacral vertebrae.

The left side of each carcass was cut into wholesale cuts and physically separated into muscle, fat and bone with the exception of the round, from which the rump was removed and then the two parts were individually physically separated; and the rib which was cut into the 6-7-8, 9-10-11 and 12th rib sections and each section was physically separated. The right side of each carcass was cut into boneless, closely trimmed (approximately 0.3 in.) retail cuts by wholesale cut.

In most instances, highly significant (P < .01) correlations were found between the fat probes and separable components, retail and fat trim yields. As high as 96% of the variation in carcass separable fat, external fat trim from the round, loin, rib and chuck and total retail fat trim could be accounted for in either pounds or percent of these variables by combinations of fat thickness, probes and carcass weight.

Percent separable muscle and fat from the 9-10-11 rib section and the wholesale flank showed the highest and most consistent relationships to percent carcass separable muscle and fat of any of the wholesale cuts. The ease of separation and low economic value of the flank provided a more rapid yet accurate cut than those presently used for prediction of carcass composition. Eighty-eight percent of the variation in percent carcass separable muscle, 94% of the variation in percent carcass separable fat and 83% of the variation in percent carcass separable bone could be accounted for by a combination of flank separable components and objective carcass measurements.

Percent retail yield of the flank was also highly related (P < .01) to percent boned, trimmed round, loin, rib and chuck and total carcass retail yield (0.81 and 0.88, respectively) and was included in prediction equations to estimate both of these carcass yields.

Low, non-significant correlation coefficients were found between 1.

dorsi muscle potassium and sodium and 1. dorsi muscle weight, area and total carcass separable muscle (range, 0.00 to 0.30). Means of the organoleptic analyses were not consistent with compositional data.

# THE RELATIONSHIP OF SOME LINEAR AND PHYSICAL MEASUREMENTS TO BEEF CARCASS COMPOSITION

By North M. Allen

## A THESIS

Submitted to
Michigan State University
in partial fulfillment of the requirements
for the degree of

DOCTOR OF PHILOSOPHY

Department of Animal Husbandry

1966

142006 44451

### ACKNOWLEDGEMENTS

The author wishes to express his sincere appreciation and thanks to Dr. R. A. Merkel and Dr. W. T. Magee of the Animal Husbandry Department for the invaluable guidance, assistance, and encouragement they have extended during this study. Special appreciation is also expressed to Mr. Ken Kemp and Mrs. Bea Eichelberger for their invaluable help.

Appreciation is also expressed to all members of the guidance committee and to the Allen Packing Company of Charlotte, Michigan, and Van Alstine Packing Company of Okemos, Michigan.

Most of all the author wishes to thank his wife, Joyce, for encouragement and help during this study and a special thanks to his parents for their understanding and encouragement throughout his lifetime.

## TABLE OF CONTENTS

Page	1
INTRODUCTION	
REVIEW OF LITERATURE	
Early Beef Cattle Evaluation Studies 3	
Growth	
Fattening	
Physical Methods for Estimating Carcass Composition 7	
Physical Separation Studies	
9-10-11 Rib Section Separation	
12th Rib Separation	
Round, Chuck and Foreshank Separation 8	
Flank Separation	
Relationship of Individual and Groups of Muscles to	
Carcass Composition	
Carcass Composition	
Polotionship of Tipoer Mossurements to Carooss	
Relationship of Linear Measurements to Carcass	
Outpool20201	
Specific Gravity	
Chemical Methods for Estimating Carcass Composition 12	
Relationship Between Muscle Content of Potassium and	
Sodium and Carcass Composition	
Linear and Multiple Regression Equations for Predicting	
Carcass Composition	
Beef Carcass Retail Yield Studies	
Variability of Beef Carcasses Retail Yield 15	
Factors Influencing Variability of Retail Yield 16	
Influence of Fat Upon Retail Yield 16	
Influence of Carcass Weight Upon Retail Yield 19	
Influence of Conformation Upon Retail Yield 20	
Influence of Bone Upon Retail Yield	
Regression Equations for Measuring Differences in	
Retail Yield 24	

Pag	;e
EXPERIMENTAL PROCEDURE	)
Source of Material	)
Grouping	)
Slaughtering Procedure	,
Subjective Carcass Evaluation	,
Linear Carcass Measurements	•
Cutting Procedure       31         Left Side       33         Bone and Muscle Study       33         Right Side       33	} }
Tenderness Studies	ŀ
Chemical Analysis	j
Statistical Analysis	;
RESULTS AND DISCUSSION	<u>;</u>
Effect of Carcass Weight on Pounds and/or Percent Carcass Separable Components, Retail and Fat Trim Yields 36	<b>,</b>
Effect of Fat Thickness on Pounds and/or Percent Carcass Separable Components, Retail and Fat Trim Yields 41	
Carcass Weight and Fat Thickness Interaction Effects on Pounds and/or Percent Carcass Separable Components, Retail and Fat Trim Yields	-
Relationships Between Subjective Carcass Scores and Separable Components, Retail and Fat Trim Yields 42	:
Relationship Between Linear Fat Measurements and Separable Components, Retail and Fat Trim Yields 47	,
Relationships Between Some Linear Carcass Measurements and Separable Components, Retail and Fat Trim Yields 56	,

Page

	Page
Relationships Between Weight of Certain Entire Muscles and Muscle Groups and Weight of Total Separable Carcass Muscle and Bone	114
Relationships Between Weights of Certain Entire Bones and Bone Groups and Weight of Total Separable Carcass Muscle and Bone	114
Multiple Regression Analyses of Pounds and Percent Separable Carcass Muscle and Bone on Individual and Groups of Muscle and Bone Weights	117
Weights	117 119
Relationships Between Weight and Percent Total Muscle, Fat,	121
Relationships Between Wholesale Cut Retail Yield and Total	121
Carcass Separable Components, Retail and Fat Trim Yields	123
Relationships Between Pounds and Percent Wholesale Cut Fat Trim and Total Carcass Separable Components, Retail and Fat Trim Yields	127
Relationships Between Wholesale Cut Bone Yield and Total Carcass Separable Components, Retail and Fat Trim Yields	132
Multiple Regression Analyses of Pounds and Percent Retail Yield Measures on a Combination of Wholesale Cut Yields and Carcass Scores and Measurements	136
Yields and Objective Carcass Measurements  Multiple Regression Analyses of Pounds and Percent Boned, Trimmed Retail Round, Loin, Rib and Chuck Yield on Objective Carcass Measurements and Percent	136 it
Flank Retail Yield	139
Combination of Weights of Wholesale Cut Yields and Carcass Scores and Measurements	143
Carcass Scores and Measurements	146
ments and Percent Flank Retail Yield	147

	Page
Relationships Between <u>L</u> . <u>dorsi</u> Muscle Potassium and Sodium and <u>L</u> . <u>dorsi</u> Weight, Area and Total Carcass Separable Muscle	<b>!</b>
Means and Standard Deviations of Some Subjective Carcass Scores and Some Chemical and Organoleptic Characteris	
tics	. 152
SUMMARY	. 155
LITERATURE CITED	. 161
SUPPLEMENT	. 168
A PPENDTX	. 189

# LIST OF TABLES

Table		Page
1	Distribution of carcasses within weight and fat thickness groups	26
2	Characteristics and scores used in the subjective carcass evaluation	28
3	Means and standard deviations of weights of carcass separa	ble
	components, retail and fat trim yields, 1. dorsi muscle area and average 12th rib fat thickness	37
3 <b>a</b>		
		38
4	Means and standard deviations of percent carcass separable components, retail and fat trim yields	
4a		40
5	Simple correlation coefficients between subjective carcass scores and carcass separable components, retail and fat trim yields for the combined weight groups	
6	Simple correlation coefficients between subjective carcass scores and carcass separable components, retail and fat trim yields for the 500 to 550 lb. weight group	
7	Simple correlation coefficients between subjective carcass scores and carcass separable components, retail and fat trim yields for the 700 to 750 lb. weight group	
8	Simple correlation coefficients between linear fat measure ments and carcass separable components, retail and fat trim yields for the combined weight groups	
9	Simple correlation coefficients between linear fat measure ments and carcass separable components, retail and fat tri yields for the 500 to 550 lb. weight group	m
10	Simple correlation coefficients between linear fat measure ments and carcass separable components, retail and fat tri yields for the 700 to 750 lb. weight group	m
11	Simple correlation coefficients between some linear carcas measurements and carcass separable components, retail and fat trim yields for the combined weight groups	

Table Page

12	Simple correlation coefficients between some linear carcass measurements and carcass separable components, retail and fat trim yields for the 500 to 550 lb. and 700 to 750 lb. weight groups
13	Coefficient of determination between weight of carcass separable fat and a combination of objective carcass fat measurements, and the regression coefficient for each of the measurements (500 to 550 lb. weight group) 60
14	Coefficient of determination between percent carcass separable fat and a combination of objective carcass fat measurements, and the regression coefficient for each of the measurements (500 to 550 lb. weight group) 61
15	Coefficient of determination between weight of total carcass separable fat and a combination of objective carcass fat measurements, and the regression coefficient for each of the measurements (combined weight groups) 62
16	Coefficient of determination between percent carcass separable fat and a combination of objective carcass fat measurements, and the regression coefficient for each of the measurements (combined weight groups) 63
17	Coefficient of determination between weight of external fat trim from the round, loin, rib and chuck and a combination of objective carcass fat measurements, and the regression coefficient for each of the measurements (combined weight groups)
18	Coefficient of determination between weight of external fat trim from the round, loin, rib and chuck and a combination of objective carcass fat measurements, and the regression coefficient for each of the measurements (500 to 550 lb. weight group)
19	Coefficient of determination between weight of total retail fat trim and a combination of objective carcass fat measurements, and the regression coefficient for each of the measurements (combined weight groups)
20	Coefficient of determination between weight of total retail fat trim and a combination of objective carcass fat measurements, and the regression coefficient for each measurement (500 to 550 lb. weight group) 67
21	Coefficient of determination between percent total retail fat trim and a combination of objective carcass fat measurements, and the regression coefficient for each of the measurements (500 to 550 lb. weight group) 68

22	Coefficient of determination between weight of separable carcass muscle and combination of objective carcass measurements, and the regression coefficient for each of the measurements (combined weight groups)	
23	Coefficient of determination between percent separable carcass muscle and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (500 to 550 lb. weight group)	70
24	Coefficient of determination between weight of total carcas separable bone and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (combined weight groups)	
25	Coefficient of determination between percent carcass separable bone and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (combined weight groups)	72
26	Coefficient of determination between weight of boned, trimmed round, loin, rib and chuck retail yield and a combination of objective measurements, and the regression coefficient for each of the measurements (combined weight groups)	<b>.</b> 73
27	Coefficient of determination between weight of total carca- boned, trimmed retail yield and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (combined weight groups)	
28	Simple correlation coefficients between wholesale cut separable muscle and carcass separable components, retail and fat trim yields for the combined weight groups	76
29	Simple correlation coefficients between wholesale cut separable muscle and carcass separable components, retail and fat trim yields for the 500 to 550 lb. weight group.	77
30	Simple correlation coefficients between wholesale cut separable muscle and carcass separable components, retail and fat trim yields for the 700 to 750 lb. weight group.	78
31	Simple correlation coefficients between wholesale cut separable fat and carcass separable components, retail and fat trim yields for the combined weight groups	82
32	Simple correlation coefficients between wholesale cut separable fat and carcass separable components, retail	83

Table		Page
33	Simple correlation coefficients between wholesale cut separable fat and carcass separable components, retail and fat trim yields for the 700 to 750 lb. weight group.	84
34	Simple correlation coefficients between wholesale cut separable bone and carcass separable components, retail and fat trim yields for the combined weight groups	86
35	Simple correlation coefficients between wholesale cut separable bone and carcass separable components, retail and fat trim yields for the 500 to 550 lb. weight group.	87
36	Simple correlation coefficients between wholesale cut separable bone and carcass separable components, retail and fat trim yields for the 700 to 750 lb. weight group.	88
37	Simple correlation coefficients between percent separable components of some wholesale cuts and separable components of the carcass	s 91
38	Coefficient of determination between weight of carcass separable muscle and a combination of weights of wholesal cut separable components and objective carcass measurement and the regression coefficient for each of the variables (combined weight group)	
39	Coefficient of determination between percent carcass separable muscle and a combination of weights of wholesale cut separable components and objective carcass measurements, and the regression coefficient for each of the variables (combined weight group)	r- 95
40	Coefficient of determination between percent carcass separable muscle and a combination of weights of wholesale c separable components and objective carcass measurements, the regression coefficient for each of the variables (500 to 550 lb. weight group)	and
41	Coefficient of determination between weight of carcass seriable muscle and a combination of weights of wholesale comparable components and objective carcass measurements, the regression coefficient for each of the variables (700 750 lb. weight group)	ut and to
42	Coefficient of determination between weight of carcass ser arable muscle and a combination of weights of wholesale components, and the regression coefficient for each variable (combined weight group)	ut

43	coefficient of determination between percent carcass separable muscle and a combination of weights of wholesale cut separable components and the regression coefficient for each of the variables (combined weight groups) 99
44	Multiple regression equations for estimating pounds and/or percent total carcass separable muscle using objective carcass measurements and percent wholesale cut separable muscle as predictors, and the coefficient of determination for each equation (combined weight group) 100
45	Multiple regression equations for estimating pounds and/or percent total carcass separable muscle using objective carcass measurements and percent wholesale cut separable muscle as predictors, and the coefficient of determination for each equation (500 to 550 lb. weight group) 101
46	Multiple regression equations for estimating pounds and/or percent total carcass separable muscle using objective carcass measurements and percent wholesale cut separable muscle as predictors, and the coefficient of determination for each equation (700 to 750 lb. weight group) 102
47	Multiple regression equations for estimating pounds and/or percent total carcass separable muscle using objective carcass measurements and flank separable components as predictors, and the coefficient of determination for each equation (combined weight groups)
48	Multiple regression equations for estimating pounds and/or percent total carcass separable muscle using objective carcass measurements and flank separable components as predictors, and the coefficients of determination for each equation (500 to 550 lb. weight group) 104
49	Multiple regression equations for estimating pounds and/or percent total carcass separable muscle using objective carcass measurements and flank separable components as predictors, and the coefficient of determination for each equation (700 to 750 lb. weight group) 105
50	Multiple regression equations for estimating pounds and/or percent total carcass separable fat using objective carcass measurements and percent wholesale cut separable fat as predictors, and the coefficient of determination for each equation (combined weight group)
51	Multiple regression equations for estimating pounds and/or percent total carcass separable fat using objective carcass measurements and percent wholesale cut separable fat as predictors, and the coefficient of determination for each equation (500 to 550 lb. weight group)

52	Multiple regression equations for estimating pounds and/or percent total carcass separable fat using objective carcass measurements and percent wholesale cut separable fat as predictors, and the coefficient of determination for each equation (700 to 750 lb. weight group) 109
53	Multiple regression equations for estimating pounds and/or percent total carcass separable bone using objective carcass measurements and percent wholesale cut separable components as predictors, and the coefficient of determination for each equation (combined weight groups)
54	Multiple regression equations for estimating pounds and/or percent total carcass separable bone using objective carcass measurements and percent wholesale cut separable components as predictors, and the coefficient of determination for each equation (500 to 550 lb. weight group)
55	Multiple regression equations for estimating pounds and/or percent total carcass separable bone using objective carcass measurements and percent wholesale cut separable components as predictors, and the coefficient of determination for each equation (700 to 750 lb. weight group)
56	Simple correlation coefficients between some individual muscle and muscle group weights and carcass separable muscle and bone
57	Simple correlation coefficients between individual bone and bone group weights and carcass separable muscle and bone
58	Coefficient of determination between weight of carcass separable muscle and a combination of muscle and bone weights, and the regression coefficient for each of the variables (combined weight groups)
59	Coefficient of determination between percent carcass separable muscle and a combination of muscle and bone weights, and the regression coefficient for each of the variables (500 to 550 lb. weight group)
60	Coefficient of determination between weight of separable carcass bone and a combination of muscle and bone weights, and the regression coefficient for each of the variables (combined weight groups)
61	Coefficient of determination between percent carcass separable muscle and a combination of muscle and bone weights, and the regression coefficient for each of the variables (combined weight groups)

Table		Page
62	Simple correlation coefficients between weights and percents of total carcass separable muscle, fat and bone for the combined and individual weight groups	122
63	Simple correlation coefficients between wholesale cut retail yield and carcass separable components, retail and fat trim yields for the combined weight groups	124
64	Simple correlation coefficients between wholesale cut retail yield and carcass separable components, retail and fat trim yields for the 500 to 550 lb. weight group	
65	Simple correlation coefficients between wholesale cut retail yield and carcass separable components, retail and fat trim yields for the 700 to 750 lb. weight group	126
66	Simple correlation coefficients between wholesale cut fat trim and carcass separable components, retail and fat trin yields for the combined weight groups	
67	Simple correlation coefficients between wholesale cut fat trim and carcass separable components, retail and fat trin yields for the 500 to 550 lb. weight group	
68	Simple correlation coefficients between wholesale cut fat trim and carcass separable components, retail and fat trip yields for the 700 to 750 lb. weight group	
69	Simple correlation coefficients between wholesale cut bonyield and carcass separable components, retail and fat tryields for the combined weight groups	im
70	Simple correlation coefficients between wholesale cut bonyield and carcass separable components, retail and fat tryields for the 500 to 550 lb. weight group	im
71	Simple correlation coefficients between wholesale cut bonyield and carcass separable components, retail and fat tryields for the 700 to 750 lb. weight group	<b>i</b> m
72	Coefficient of determination between weight of boned, tripround, loin, rib and chuck retail yield and a combination of weights of wholesale cut yields and objective carcass measurements, and the regression coefficient for each of the variables (combined weight groups)	
73	Coefficient of determination between percent boned, trimmoround, loin, rib and chuck retail yield and a combination of percent wholesale cut yields and carcass scores and measurements, and the regression coefficient for each of the variables (combined weight groups)	

Table Page

74	Multiple regression equations for estimating pounds and/or percent boned, trimmed round, loin, rib and chuck retail yield using objective carcass measurements and percent flank retail yield as predictors, and the coefficient of determination for each equation (combined weight groups) 140
75	Multiple regression equations for estimating pounds and/or percent boned, trimmed round, loin, rib and chuck retail yield using objective carcass measurements and percent flank retail yield as predictors, and the coefficient of determination for each equation (500 to 550 lb. weight group)
76	Multiple regression equations for estimating pounds and/or percent boned, trimmed round, loin, rib and chuck retail yield using objective carcass measurements and percent flank retail yield as predictors, and the coefficient of determination for each equation (700 to 750 lb. weight group)
77	Coefficient of determination between pounds of boned, trimmed total carcass retail yield and a combination of weights of wholesale cut yields and objective carcass measurements, and the regression coefficient for each of the variables (combined weight groups)
78	Coefficient of determination between percent boned, trimmed total carcass retail yield and a combination of weights of wholesale cut yields and carcass scores and measurements, and the regression coefficient for each of the variables (combined weight groups)
79	Coefficient of determination between weight of boned, trimmed total carcass retail yield and a combination of weights of wholesale cut yields and objective carcass measurements, and the regression coefficient for each of the variables (500 to 550 lb. weight group)
80	Coefficient of determination between percent boned, trimmed total carcass retail yield and a combination of weights of wholesale cut yields and objective carcass measurements, and the regression coefficient for each of the variables (500 to 550 lb. weight group)
81	Multiple regression equations for estimating pounds and/or percent boned, trimmed total carcass retail yield using objective carcass measurements and percent flank retail yield as predictors, and the coefficient of determination for each equation (combined weight groups)

Table	Page
82	Multiple regression equations for estimating pounds and/or percent boned, trimmed total carcass retail yield using objective carcass measurements and percent retail yield as predictors, and the coefficient of determination for each equation (500 to 550 lb. weight group) 149
83	Multiple regression equations for estimating pounds and/or percent boned, trimmed total carcass retial yield using objective carcass measurements and percent flank retail yield as predictors, and the coefficient of determination for each equation (700 to 750 lb. weight group) 150
84	Simple correlation coefficients between muscle potassium and sodium and 1. dorsi muscle weight and area, and total carcass separable muscle
85	Means and standard deviations of several subjective carcass scores and some chemical and organoleptic characteristics 153

# LIST OF FIGURES

Figure		Page
1	Illustration showing fat thickness measurements taken at the 12th rib	29
2	Illustration showing the pattern of the fat probes taken 4, 8 and 12 in. from the midline on the carcass	30
3	Illustration showing the measurement of fat depth over the brisket	32

# LIST OF SUPPLEMENT TABLES

rable		Page
A	Coefficient of determination between weight of carcass separable fat and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (combined weight groups)	168
В	Coefficient of determination between percent carcass separable fat and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (combined weight groups)	168
С	Coefficient of determination between weight of carcass separable fat and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to 750 lb. weight group)	169
D	Coefficient of determination between percent carcass separable fat and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to 750 lb. weight group)	169
E	Coefficient of determination between percent carcass separable fat and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (500 to 550 lb. weight group)	170
F	Coefficient of determination between percent carcass separable fat and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to 750 lb. weight group)	170
G	Coefficient of determination between percent external fat trim from the round, loin, rib and chuck and a combination of objective carcass measurements, and the regression co- efficient for each of the measurements (combined weight groups)	n
н	Coefficient of determination between percent external fat trim from the round, loin, rib and chuck and a combination of objective carcass measurements, and the regression co- efficient for each of the measurements (500 to 550 lb. weight group)	
I	Coefficient of determination between percent external fat trim from the round, loin, rib and chuck and a combination of objective measurements, and the regression coefficient for each of the measurements (700 to 750 lb. weight group	

J	Coefficient of determination between percent total retail fat trim and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (combined weight groups)
К	Coefficient of determination between weight of total retail fat trim and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to 750 lb. weight group)
L	Coefficient of determination between percent total retail fat trim and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to 750 lb. weight group)
М	Coefficient of determination between percent carcass separable muscle and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (combined weight groups)
N	Coefficient of determination between weight of carcass separable muscle and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (500 to 550 lb, weight group)
0	Coefficient of determination between weight of carcass separable muscle and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to 750 lb. weight group) 175
P	Coefficient of determination between percent carcass separable muscle and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to 750 lb. weight group)
Q	Coefficient of determination between weight of carcass separable bone and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (500 to 550 lb. weight group) 175
R	Coefficient of determination between percent carcass separable bone and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (500 to 550 lb. weight group)
S	Coefficient of determination between weight of carcass separable bone and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to 750 lb. weight group) 176

T	Coefficient of determination between percent carcass separable bone and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to 750 lb. weight group) 1	.76
U	Coefficient of determination between weight of carcass separable muscle and a combination of objective carcass measurements and weight of flank separable components, and the regression coefficient for each of the variables (500 to 550 lb. weight group)	7
V	Coefficient of determination between percent carcass separable muscle and a combination of objective carcass measurements, flank separable component weights and kidney knob weight, and the regression coefficient for each of the variables (700 to 750 lb. weight group)	.77
W	Coefficient of determination between weight of carcass separable muscle and a combination of carcass weight and weights of flank separable components, and the regression coefficient for each of the variables (500 to 550 lb. weight group)	'7
X	Coefficient of determination between weight of carcass separable muscle and a combination of carcass weight and weights of flank separable components, and the regression coefficient for each of the variables (700 to 750 lb. weight group)	'8
Y	Coefficient of determination between percent carcass separable muscle and a combination of carcass weight and weights of flank separable components, and the regression coefficient for each of the variables (500 to 550 lb. weight group)	'8
Z	Coefficient of determination between percent carcass separable muscle and a combination of carcass weight and weights of flank separable components, and the regression coefficient for each of the variables (700 to 750 lb. weight group)	'8
AA	Coefficient of determination between percent carcass separable muscle and a combination of muscle and bone weights, and the regression coefficient for each of the variables (combined weight groups)	'9
AB	Coefficient of determination between weight of carcass separable muscle and a combination of muscle and bone weights, and the regression coefficient for each of the variables (500 to 550 lb. weight group)	<b>'</b> 9

Page

AM	rib and chuck retail yield and a combination of objective carcass measurements, foreshank bone weight and percent flank fat trim, and the regression coefficient for each of the variables (500 to 550 lb. weight group)
AN	Coefficient of determination between weight of round, loin, rib and chuck retail yield and a combination of carcass measurements, scores and percent flank yield, and the regression coefficient for each of the variables (700 to 750 lb. weight group)
AO	Coefficient of determination between percent round, loin, rib and chuck retail yield and a combination of carcass measurements, scores and percent flank yield, and the regression coefficient for each of the variables (700 to 750 lb. weight group)
AP	Coefficient of determination between weight of carcass retail yield and carcass weight, flank yield and fat trim, and the regression coefficient for each of the variables (700 to 750 lb. weight group)
AQ	Coefficient of determination between percent carcass retail yield and a combination of kidney knob weight and flank retail yield and fat trim weights, and the regression coefficient for each of the variables (700 to 750 lb. weight group)
AR	Coefficient of determination between weight of carcass retail yield and a combination of objective carcass measurements and percent flank yield, and the regression coefficient for each of the variables (combined weight groups) 186
AS	Coefficient of determination between percent carcass retail yield and a combination of carcass measurements, scores and percent flank yield, and the regression coefficient for each of the variables (combined weight groups) 186
AT	Coefficient of determination between weight of carcass retail yield and a combination of objective carcass measurements and percent flank fat trim, and the regression coefficient for each of the variables (500 to 550 lb. weight group) . 187
AU	Coefficient of determination between percent carcass retail yield and a combination of objective carcass measurements and percent flank fat trim, and the regression coefficient for each of the variables (500 to 550 lb. weight group). 187

Table		Page
AV	Coefficient of determination between weight of carcass retail yield and a combination of carcass weight and percent flank yield, and the regression coefficient for each variable (700 to 750 lb. weight group)	188
AW	Coefficient of determination between percent carcass retail yield and a combination of carcass weight, percent flank yield and percent kidney knob, and the regression coefficient for each of the variables (700 to 750 lb. weight group)	188

## LIST OF APPENDIX TABLES

Appendix		Page
IA	Carcass evaluation - subjective scores	189
I B	Carcass evaluation - objective fat measures	192
I C	Carcass evaluation - objective measurements	207
II A	Right side - wholesale cut yield	210
II B	Retail cutout - right side	216
III A	Physical separation of wholesale cuts - left side	222
III B	Separable muscle, fat and bone of 6-7-8th, 9-10-11th,	227
	and 12th rib sections	237
III C	Muscle and bone weights	240
IV	Chemical and organoleptic data	246
	List of variables on data cards	249

#### INTRODUCTION

Consumer research studies have repeatedly shown that the American consumer selects beef primarily for maximum quantity of muscle with a minimum amount of fat and/or bone. Today's retailer reflects consumer demand for well muscled, trim retail cuts in his purchases of carcass beef. Additionally, retail cut yield grade is an optional feature in the present U. S. Department of Agriculture beef grade specifications. Thus, the producer must market cattle with a high quantity of inherent muscling and a minimum of trimable fat to provide the retail cuts demanded by today's beef industry and consumer. Beef carcasses commensurate with these specifications should have at least minimum U. S. D. A. Choice quality, external fat thickness between 0.25 and 0.5 in. and yield a minimum of 65% boneless trimmed retail cuts. Such cattle exist today, but many beef carcasses have excessive fat or deficient muscling and combinations of these two factors.

Objective, yet practical, methods are needed to accurately identify and measure carcass quantitative and qualitative differences. When such methods become available, it is anticipated they would be applicable for characterizing these same differences in live cattle; thus, providing more objective criteria for selection of breeding stock to meet today's consumer demand.

The most accurate methods to date for measuring carcass quantitative differences necessitate destruction of at least a portion of the carcass (physical separation and chemical analyses). Non-destructive methods

currently used involve various combinations of objective and subjective criteria, but none accurately measures quantitative differences and/or value. Breidenstein (1962) reported as much as a \$13.54 per hundred-weight difference in value within the same carcass grade. Previous beef carcass composition studies have consistently indicated that degree of fatness has a very marked effect upon yield of boneless, trimmed retail cuts. Yet fat thickness as conventionally measured at the 12th rib has repeatedly been found to account for less than 40% of the variation in total carcass fat. Likewise, indices of muscling such as area of <u>longissimus dorsi</u> as well as other muscles account for less than 40% of the variation in yield of boneless, trimmed retail cuts.

Thus, this study was initiated in an attempt to more accurately predict total physically separable fat and muscle as well as yield of boneless, trimmed retail cuts using a rather comprehensive number of objective measurements. The specific objectives of this study were: 1. To develop a repeatable, accurate and practical method for measuring total separable fat and muscle in beef carcasses. 2. To study the variation in percent trimmed wholesale round, loin, rib and chuck from steer carcasses of approximately the same weight and degree of fatness. 3. To study the variation in actual boneless, trimmed retail yield among steer carcasses of approximately the same weight and degree of fatness from estimations by several existing regression equations. 4. To develop a regression equation in an attempt to more accurately account for the total variation in boneless, trimmed retail yield.

## REVIEW OF LITERATURE

## Early Beef Cattle Evaluation Studies

Research studies during the last half-century have revealed the tremendous variability in production traits and carcass characteristics which exists within the beef cattle population.

The investigations of Lawes and Gilbert (1860) and Henry and Sanborn (1883-1890) were among the first reported studies of carcass composition following the growing and fattening period. Trowbridge et al. (1918) conducted an exhaustive study of the chemical composition of gains made by three year old steers during the fattening process to determine what changes take place in the form of the animal. They observed a marked increase in fat during the fattening period.

Robert Bakewell is credited with being the first great improver of cattle. He developed a low-set, blocky, quick-maturing type of cattle through selection. As quoted by Ensminger (1955), "His objective was to breed cattle that would yield the greatest quantity of good beef rather than to obtain great size."

In the ensuing years, "beef-type" cattle have become identified with selection for increased quantity of muscle relative to other cattle types.

Wilson and Curtis (1893) reported limited data comparing performance traits and carcass qualities of beef- and dairy-type steers. Dairy- and beef-type steers had approximately equal gains and percent of high priced cuts. These workers concluded, however, that the carcasses from the

dairy-type steers were deficient in conformation or shape of the high priced cuts, as well as having less marbling. Moreover, they reported excessive internal fat in the carcasses from some of the dairy-type steers.

Knapp and Nordskog (1946a, b), using data from the U. S. Range
Livestock Experiment Station, presented the first known heritability
estimates of quantitative traits in beef cattle. Heritability estimates
have been determined for most of the economically important performance
traits in beef cattle (Black and Knapp, 1936; Black, 1938; Knapp et al.,
1941; Knapp and Black, 1942; Knapp and Clark, 1950; Warwick, 1958; Carter
and Kincaid, 1959; Shelby et al., 1960; Gregory et al., 1961). These
early studies were primarily concerned with production traits and type
studies (Knox and Koger, 1946). More recently, other traits such as
length of calving interval, maternal ability, dressing percent, 1. dorsi
muscle area, tenderness, and other carcass quantitative and qualitative
characteristics have been studied. Heritability has been established for
many of these traits (Knapp and Clark, 1950; Clark, 1954; Warwick, 1958;
Gregory et al., 1961).

## Growth

The recent emphasis upon dietary weight control as well as the controversial incrimination of animal fats in cardiovascular diseases has precipitated much interest in body composition studies. Before attempting to study composition, a knowledge of animal body growth and development is necessary.

Meek (1901), Brody and Ragsdale (1924) and Lush (1928) were among the early workers who studied post-natal growth. They studied external body measurements and live weight and concluded that the latter increased at a faster rate than any other single body measurement. In addition, they found that linear skeletal measurements such as measurements of the skull and height measurements over the shoulder and rump increased at a slower rate than did measurements of fat and muscle mass. These workers determined that the skeleton was better developed at birth than were muscle and fat masses which constitute the greatest proportion of total body mass at normal slaughter weight. The conclusions of these early workers were supported by the findings of Hammond (1932), McMeekan (1941), Wallace (1948) and Palsson and Verges (1952) resulting from complete dissection of swine and sheep carcasses. These latter workers concluded that a primary growth wave occurs from the cranium to the facial parts of the head and posteriorally to the lumbar region. A secondary growth wave starts in the metacarpals and metatarsals and continues down toward the digits and upward along the limbs to the lumbar region. These findings led to the measurement of area of the 1. dorsi muscle in the middle of the back as an index of muscling. Palsson (1939) reported that muscle development could most accurately be determined from the cross-sectional area of the 1. dorsi muscle at the last rib, since this area of the animal body is the last to reach full development.

Luitingh (1962) observed from dissection studies of cattle that the cut, "the loin" was not the latest developing part of the animal. This

agrees with the findings of Butterfield (1963b) who concluded from his studies with beef cattle that muscles along the spine were constant in their rate of development when compared with the rate of development of total carcass lean. In an extensive dissection study, he determined the rate of proportional increase in muscle groups in different breeds of cattle using as a control the proportion of these muscle groups from pre-natal, but near full-term calves. He classified the different muscles as early developing, late developing, very late developing and average developing muscles. Muscles which he classified as early developing were those muscles which at birth had a weight-relation to total carcass muscle that was greater than the same weight-relation in the mature animal. These were the intrinsic muscles of the forelimb and the distal intrinsic muscles of the hind limb. Late developing muscles were those which had a weight-relation to total carcass muscle at birth that was less than that in the mature animal. Late developing muscles were abdominal muscles and poximal muscles of the hind limb. Very late developing muscles were those which had a weight-relation to total carcass muscle at birth that was less than that in the mature animal and which do not increase relative to total muscle until late in life. The intrinsic muscles of the neck and thorax and muscles of the neck and thorax which are attached to the thoracic limb were classified as very late developing. Average developing muscles were those whose weight-relation to that of total carcass muscle did not change during post-natal life and included only those muscles surrounding the spinal column.

## Fattening

Hankins and Titus (1939) stated that in young growing animals weight gains are composed largely of protein and water; whereas, those of the mature or nearly mature animal consist primarily of fat. These workers reported that one of the best known and most obvious changes which accompanies growth and fattening is the increase in the ratio of carcass weight to the weight of the entire body (dressing percent). Warner et al. (1934) reported that as the hog grows and fattens, the percent ham, loin, shoulder and head decrease. They also reported an increase in the percent bacon and fat trim. Hankins and Titus (1939) found that in beef, the percent rib, short loin, plate and flank increased as the animal fattened and the percent round, sirloin and foreshank decreased. The chuck and rump showed very little change.

Physical Methods for Estimating Carcass Composition

Physical Separation Studies. Physical separation data of entire beef carcasses are limited because such studies are laborious, time consuming and involve economic loss of product. However, physical separation of wholesale cuts or parts thereof has been used rather extensively to measure beef carcass composition.

9-10-11 Rib Section Separation. The most widely used method of estimating beef carcass composition is physical separation of the 9-10-11 rib section as described by Hankins and Howe (1946). They reported correlation coefficients between the percent separable muscle, fat and bone from

the 9-10-11 rib section with the same components from the entire carcass of 0.85, 0.93 and 0.83, respectively. The conclusions of these workers were supported by the findings of Crown and Damon (1959) who reported correlation coefficients of 0.94, 0.98 and 0.73 for muscle, fat and bone, respectively, between these components in the 9-10-11 rib section and the same components in the carcass.

12th Rib Separation. Crown and Damon (1959) also reported correlation coefficients between the percent separable carcass muscle, fat and bone and the same separable components of the 12th rib section of 0.82, 0.96 and 0.75, respectively. They suggested using the 12th rib section to predict carcass composition for greater economy of time and labor since results were comparable to those from the 9-10-11 rib section.

Round, Chuck and Foreshank Separation. Cole et al. (1960) reported high relationships between separable muscle of the round, chuck and foreshank and total separable muscle of the carcass (0.95, 0.93, and 0.81, respectively). Separable muscle of these wholesale cuts was associated with 90, 87 and 66%, respectively, of the variation in total separable muscle of the carcass. These workers suggested the use of either the round or foreshank as an index of carcass muscling due to their high predictive value and relative ease of separation.

Flank Separation. Hankins and Howe (1946), Hedrick et al. (1963) and Miller et al. (1965) reported that retail yield or separable muscle and fat of the flank is highly related to total beef carcass muscle and

fat. Hankins and Howe (1946) reported that the correlation between percent separable fat in the flank with percent ether extract in the carcass was high (0.95). Miller et al. (1965) reported that the percent yield of muscle from the wholesale flank showed the second highest relation to percent total carcass retail yield, the round being the most highly related. These workers concluded that yield of muscle from the flank was relatively constant but that it was a fat depot and the amount of fat in the flank was highly related to total fat trim or total separable fat.

Relationship of Individual and Groups of Muscles to Carcass Composition. The relationship between several individual muscles and total carcass muscle has been shown to be quite high by several workers (Orme et al., 1960; Dumont et al., 1961; Butterfield, 1963a; and Miller et al., 1965). These workers reported a high correlation between 1. dorsi muscle weight and total separable carcass muscle (approximately 0.90). Weight of the biceps femoris muscle was reported by Orme et al. (1960) and Butterfield (1963b) to be highly correlated to total carcass muscle (0.96).

Orme et al. (1960) also reported high correlation coefficients between total separable carcass muscle and the semimembranosus plus adductor (0.92), semitendinosus (0.83), quadriceps muscle group (0.93), psoas major (0.82), triceps brachii (0.84) and infraspinatus (0.80) muscles.

Relationship of Individual and Groups of Bones to Carcass Composition.

Orme et al. (1959) reported that live weight, chilled carcass weight,

primal cut weight and estimated carcass muscle (from 9-10-11 rib physical

separation) in almost all instances were significantly and positively related to measures (weight and linear measurements) of the fore and hind cannon bones. These workers found that when the effects of live weight were removed, width and circumference measurements of the hind and fore cannons were associated with 15 to 25% of the existing variation in area of  $\underline{1}$ .  $\underline{dorsi}$  muscle. These same workers concluded that even though cannon bone measurements and radiographs of the lumbar vertebrae are related to muscling, these relations are too low to be of any predictive value. Henderson  $\underline{et}$   $\underline{al}$ . (1966) reported significant correlations (P < .01) between total carcass bone and percent separable muscle (0.77), and percent separable muscle from the four major wholesale cuts (0.73).

Butterfield (1963a) observed high correlations between the weight of various bones and total carcass bone weight in beef steers. He reported correlation coefficients between total bone weight and weight of the scapula (0.96), humerus (0.99), radius and ulna (0.97), radius and ulna plus carpus (0.98), or coxae (0.97), femur (0.98), tibia (0.97) and the tibia plus tarsus (0.96). Orme (1963) reported highly significant correlations (P < .01) between total lamb carcass muscle weight and weight of the following bones: radius and ulna, metacarpal, metatarsal, tibia and the femur. He also reported highly significant correlations (P < .01) between these same bones and total carcass bone weight (r = 0.80) as well as between weight of bones from various wholesale cuts of lamb and total carcass bone weight (total carcass bone with leg-bone 0.90, loin-bone 0.52, rack-bone 0.69, shoulder-bone 0.82 and foreshank bone 0.84).

Relationship of Linear Measurements to Carcass Composition. White and Green (1952) reported that various live animal measurements were highly correlated with weights of major beef cuts (0.5 to 0.6) and linear measurements of the live animal were of value for estimating weight of various wholesale cuts. Cook et al. (1951), Green et al. (1955), Dawson et al. (1955), Tallis et al. (1957) and Orme (1958) reported similar results. Orme (1958) and Allen (1963) reported simple correlation coefficients (0.3 to 0.8) between live animal measurements and weight of various wholesale cuts. Orme (1958) reported that when the effect of live weight was held constant, most of these relationships were markedly reduced. Cole et al. (1962) reported that carcass weight, carcass length and 1. dorsi muscle area were related to total carcass separable muscle (r = 0.75, 0.23 and 0.39, respectively). With carcass weight held constant, Cole et al. (1960) stated that 1. dorsi muscle area accounted for 5% of the variation in pounds of separable lean in the carcass.

Specific Gravity. Pearson (1965) stated that the major problem in determining specific gravity is that of accurately measuring volume. He stated that this could be done by one of two methods: 1) gaseous displacement and 2) water displacement. Morales et al. (1945) reported data for guinea pigs which provided the basic information for the calculation of equations giving the density of the fat-free body. Brown et al. (1951) and Whiteman et al. (1951) used the underwater specific gravity method for estimating the composition of carcasses and cuts from farm animals. Gnaedinger et al. (1963) using the helium dilution technique

for determining specific gravity of market weight pigs, obtained poor agreement between this value and chemical analysis.

Chemical Methods for Estimating Carcass Composition

Research workers have used various chemical methods for the determination of body composition. Among these are the antipyrene (Brodie et al., 1949) and tritium (Pace et al., 1947) methods. Moore (1946) presents an excellent discussion of the principles involved in the isotope dilution techniques for measuring total body water.

Relationship between Muscle Content of Potassium and Sodium and Carcass Composition. Kirton and Pearson (1963a,b) reported that potassium and sodium content of muscle tissue, as determined by flame photometry, was significantly related to carcass composition in sheep and hogs. With a group of 10 lamb carcasses, these workers found that the potassium content of muscle tissue when measured by flame photometry was significantly related to carcass composition but not when measured from K<sup>40</sup> content. In another group of 20 lots of ground pork and 15 lots of ground lamb, potassium content by both methods studied by these workers gave approximately the same results. However, the flame photometry method was more closely related to % water, fat and protein content of the samples. These workers concluded that a degree of precision at least equal to the flame photometry method was needed in a non-destructive method before it would become useful. Similar conclusions were made by

Kirton et al. (1961) for the K<sup>40</sup> method of predicting carcass composition in live lambs. Correlations determined by these workers were higher between measured potassium content of muscle and total muscle than between measured sodium content of muscle and total muscle. Kirton and Pearson (1963a) reported a species difference between sheep and swine with the relation between potassium and total muscle being higher in swine than in lambs. Gillett et al. (1965) further reported a significant difference in potassium and sodium content among various muscles of the pig. They also reported a significant difference in potassium and sodium content between the two breeds of swine studied. It was therefore concluded by these workers that due to the variations in potassium and sodium content (especially potassium), constancy does not exist between the muscle-element ratio and this is an important source of error when using this method of estimating composition.

Linear and Multiple Regression Equations for Predicting Carcass Composition. Numerous linear and multiple regression equations for predicting
carcass composition have been developed. Hankins and Howe (1946) were
among the early workers to develop linear regression equations from the
percent separable muscle, fat and bone of the 9-10-11th rib section for
predicting beef carcass composition.

Cole et al. (1960) developed regression equations to predict total separable muscle in beef carcasses from the pounds of separable muscle in each of several wholesale cuts (round, sirloin, shortloin, rib, chuck and foreshank).

Butterfield (1963a) calculated multiple regression equations for the estimation of total muscle, fat and bone weight from separable components as well as from some linear measurements.

Orme et al. (1960) developed linear regression, prediction equations for total separable carcass muscle using weights of individual muscles or groups of muscles as predictive factors. The latter authors stated that the fraction of the total sum of squares for separable muscle in the beef carcass associated with the degree of freedom for linear regression ranged from 63 to 91%. They reported that the four heavier muscles seemed to be the best predictors but the four lighter muscles also account for a large fraction of the variation in total carcass muscle.

Cole et al. (1962) developed a "simplified method for predicting pounds of muscle in beef carcasses". They reported that carcass weight was more closely related to total separable muscle than any other single measure and therefore should be included in the prediction equation. When they combined one fat measurement taken at the 12th rib (described by Ramsey et al., 1962) with carcass weight in a regression analysis, they were able to account for over 70% of the variation in total separable carcass muscle.

Kirton and Pearson (1963b) presented regression equations for the prediction of % water, ether-extract, and protein in empty pig carcasses using potassium content of ground samples as predicting factors. They concluded, however, that the accuracy with which composition could be predicted from potassium content was unsatisfactory under the conditions used.

#### Beef Carcass Retail Yield Studies

In recent years, numerous studies have been conducted to determine the variability and factors influencing variability of beef carcass retail yield. These have been concerned with quantitative factors as they may be related to carcass value.

Variability of Beef Carcasses Retail Yield. Research data reported by Butler (1957), Cole et al. (1960), Murphy et al. (1960) and Butler et al. (1961) have shown that great variability occurs in the retail yield of beef carcasses. These data indicate that carcass value differences exist within as well as between grades.

Kropf and Graf (1959) reported that total carcass boneless, closely trimmed retail yield varied from a high of 68.71% to a low of 57.63% among U. S. Choice, Good, Commercial (Standard) grade steer, heifer and cow carcasses ranging in weight from 400 to 900 lb.

Breidenstein (1962) found a range of 19% in retail yield and \$13.55 per hundredweight value difference among 105 steer sides and both sides of 94 heifer carcasses within the U. S. Good and Choice grades. He eliminated the extremes from the study above, thus leaving a range of 14.4% between the low and high yielding carcasses which still included 95% of the original sample. This represents a value difference between the low and high yielding carcasses of \$10.32 per hundredweight or approximately \$60.00 for a 600 pound carcass.

Brungardt and Bray (1963) conducted a similar study on 99 left sides of U. S. Choice steers. These were selected at random within three weight

groups (260 to 288 pounds, 300 to 325 pounds and 332 to 360 pounds). The light to heavy weight groups yielded an average percent boneless, trimmed retail cuts from the round, loin, rib and chuck of 50.8%, 49.7% and 48.5%, respectively. The range in percent retail yield was 47.0 to 54.6% in the light weight group, 46.3 to 53.0% in the middle weight group and 45.5 to 51.5% in the heavy weight group. The ranges included only those 22 of the 33 sides in each group closest to the average retail cut-out for the group. The ranges in value per hundredweight were \$6.56, \$5.52 and \$5.25 for the light, middle and heavy weight groups, respectively. While these values were not as large as those reported by Briedenstein (1962), this study included only the four major wholesale cuts of beef and one-third of the extremes were not included in the ranges reported.

Factors Influencing Variability of Retail Yield. Since carcasses are composed of three major tissue components, muscle, fat and bone; the proportion of these three components influences retail yield. Retail cuts as referred to in these studies consist of muscle plus fat trimmed to approximately 0.3 in. Thus, any increase in the amount of fat and/or bone results in a decrease in the percent retail yield.

Influence of Fat upon Retail Yield. Kropf and Graf (1959) reported total carcass fat trim ranged from 10.78 to 27.16% among U. S. Choice, Good and Commercial (Standard) grade steer, heifer and cow carcasses ranging in weight from 400 to 900 lb.

Ramsey et al. (1962) reported that external fat thickness at the 12th rib varied from 0.1 to 1.1 in., kidney fat from 1.8 to 8.9% and

u. S. Choice, Good and Standard grades. Brungardt and Bray (1963) reported that among carcasses grading U. S. Good and Choice, fat thickness at the 12th rib varied from 0.35 to 1.60 in. and kidney fat from 2.4 to 7.8%. Cole et al. (1962) stated that, holding carcass weight constant, external fat thickness at the 12th rib accounted for 34% of the variation in separable carcass lean.

Lewis et al. (1964) reported correlation coefficients between the third fat measurement over the 12th rib (described by Ramsey et al., 1963) and percent retail cuts and bone-in, closely trimmed retail cuts of -.26 and -.04, respectively. These workers also reported that a number of probes of subcutaneous fat over the rump and clod showed little relationship to the third fat measurement over the 12th rib, yet were negatively but highly significantly correlated with both percent bone-in and boneless, closely trimmed retail cuts. Carpenter et al. (1965) reported that each of several measurements of fat taken on the exposed surface between the 12th and 13th rib in lamb carcasses were significantly related (P < .01) to percent fat trim. These same measurements were also highly, negatively correlated with the yield of leg, loin, rack and shoulder. Hedrick et al. (1963) and Miller et al. (1965) reported significant (P < .01) negative correlations between fat thickness at the 12th rib, and several other carcass fat thickness measurements and retail yield. The latter workers reported higher correlations between subcutaneous fat thickness measurements and percent retail cuts than with weight of retail cuts. They found

that subcutaneous fat thickness measurements (probes) were nonsignificantly correlated with weight of retail cuts except those measurements and probes over the 11th to 12th thoracic vertebrae which were significant (P < .05). All correlations between fat thickness and percent boneless and partially boneless retail cuts were negative and highly significant (P < .01). These same workers reported that probes taken in the 11th to 12th thoracic vertebrae area were more closely related to retail yield than fat thickness measurements at the 12th thoracic vertebra. They stated that degree of fatness has a greater influence upon retail yield than  $\underline{1}$ .  $\underline{dorsi}$  muscle area and an increase of 0.16 in. in fat thickness at the 12th rib resulted in a 1% decrease in partially boneless retail cuts.

Zinn et al. (1963) reported a negative correlation (-.81) between percent fat trim and boneless round, loin, rib and chuck in beef carcasses. With multiple regression analysis, they found that each 1% increase in carcass fat trim, resulted in a corresponding decrease of 0.34% boneless round, loin, rib and chuck. Miller et al. (1965) reported negative and significant correlation coefficients (P < .01) between fat trim from the right and left sides with retail yield of the round, loin, rib and chuck and with total carcass retail yield. The latter workers concluded that variation in percent fat trim accounted for more of the variation in boneless and partially boneless retail yield of the carcass than any other variable studied. They stated that an increase of approximately 1.10% in fat trim decreased partially boneless retail yield 1%. Butterfield

(1965) reported that when separable fat is increased above 20%, the proportion of expensive muscles in a beef carcass is markedly reduced.

Brumgardt and Bray (1963) reported correlations between external fat thickness measurements taken at various points on the carcass and percent retail yield from the round, loin, rib and chuck ranging from -.63 to -.73. Retail yield was also negatively correlated (-.54) with percent kidney and pelvic fat. Percent kidney and pelvic fat were also reported to be negatively correlated to retail yield by Butterfield (1963a).

Murphy et al. (1960) reported high negative correlation coefficients between retail yield and a single fat thickness measurement at the 12th rib and with percent kidney knob (-.83 and -.66, respectively). Brown et al. (1962) reported on a study involving 453 steer carcasses. They observed a high negative correlation coefficient between estimated (physical separation of 9-10-11 rib) separable lean and fat (-.84). Breidenstein (1962) reported that for each one pound increase in kidney fat there was a decrease in partially boneless retail yield of 0.37% for steers and 0.44% for heifers. Kropf and Graf (1959) also found more trim fat among heifer carcasses than for the same weight and grade steer carcass.

The above results agree with those of Callow (1947) who reported that fat is the most significant variable in beef carcass composition.

Influence of Carcass Weight Upon Retail Yield. Many research studies have shown that carcass weight is negatively related to retail yield, Cole et al. (1960), Butterfield (1963a), Breidenstein (1962), Cole et al. (1962),

Brungardt and Bray (1963) and Swiger et al. (1964). Kropf and Graf (1962) observed that increased carcass weight had a significant depressing effect upon total carcass retail yield.

Cole et al. (1960) reported that carcass weight was more closely related to total separable muscle (0.77) than any other single variable studied. Butterfield (1963a) also found that carcass weight was more closely related to pounds of separable carcass muscle than any other variable. Swiger et al. (1964) reported that the simple correlation coefficient between carcass weight and percent retail yield was -.48. These workers stated that carcass weight alone accounted for 93% of the variation in retail yield.

Brungardt and Bray (1963) reported that heavier carcasses contained significantly more fat per unit of carcass weight than lighter carcasses. Cole et al. (1962) found that as carcass weight increased, the average percent steaks decreased and the percent waste increased. Breidenstein (1962) reported that a 100 lb. increase in carcass weight resulted in a reduction in retail yield of 1.42%. Murphy et al. (1960) also reported a a negative relationship between carcass weight and retail yield.

Influence of Conformation Upon Retail Yield. Briskey and Bray (1964) reported that although conformation has long been included in grading standards, its inclusion is based upon the supposition that it is related to yield of retail cuts, especially those from the round, rib, loin and chuck. They stated that this trait is related primarily to the shape and fullness of the round, loin, rib and chuck. They concluded that conformation is related to muscle development, i.e., thickness, depth and length,

but that it is also largely influenced by fat deposition both subcutaneous and intermuscular.

Breidenstein (1962) reported no significant relation between conformation score and yield of retail cuts in steers grading primarily U. S. Good and Choice. However, they observed that a one-third increase in conformation score in heifer carcasses was accompanied by an increase of 0.34% in partially boneless retail yield.

Butler (1957) reported data from beef carcasses with a wide range of conformation scores, but the percent of individual wholesale cuts was very similar. This agrees with findings of Wilson and Curtis (1893) and Branaman et al. (1962) comparing retail yield of dairy- and beef-type steers. Brungardt and Bray (1963) reported that while variation in untrimmed wholesale cut yield is small, when the cuts were trimmed to approximately 3/8 in. external fat, the yields of these trimmed wholesale cuts between beef carcasses were markedly different and closely related to boneless, trimmed retail cut yields.

In a study involving 96 carcasses, Zinn et al. (1961) reported significant (P < .01) correlations between conformation score and carcass fat thickness at the 12th rib (0.50) and with percent trimable fat (0.69). This agrees with results of Miller et al. (1965) who reported that dressing percent and conformation score were not related to retail yield but slightly related to fat thickness at the 12th rib and to trimable fat. The latter workers concluded that higher conformation scores were due largely to excess fat which was subsequently trimmed off. These data agree with

Hedrick et al. (1963) who concluded that it is difficult for superior muscle development to compensate for excess fat deposition.

In a study involving British, Zebu and dairy breeds of cattle, Cole et al. (1964) reported that Holstein steer carcasses had the highest percent separable muscle in all wholesale cuts, except the chuck and plate. Brahman steer carcasses had the highest percent separable muscle in these two wholesale cuts. Holstein carcasses had the lowest percent total separable carcass fat and were lowest in percent separable fat in all wholesale cuts, except the chuck where Brahman carcasses were lower. These authors found that the British breeds yielded the lowest percent muscle in all wholesale cuts except the foreshank where they yielded a higher percent muscle than Zebu cattle. The British breeds yielded the highest percent total carcass separable fat as well as percent separable fat from all wholesale cuts. These workers concluded from their results, that any effect conformation might exert upon percent yield of separable muscle (wholesale cuts or carcass) was overcome by the depressing effect of fat upon yield of separable muscle.

Kirton (1964) stated that intensive selection of breeding ewes for and against conformation had little influence upon the amount and value of the meat produced by their lambs. He concluded that carcasses with better conformation had less muscle and more fat than carcasses with poor conformation.

Briskey and Bray (1964) stated that it is difficult to determine muscular development in heavily fatted carcasses, since a heavily finished carcass is more likely to be scored higher in conformation than one with

less finish. They stated that from the literature available it seemed safe to conclude that conformation influenced retail yield to a far lesser extent than did degree of fatness, especially external fat.

Influence of 1. dorsi Muscle Area Upon Retail Yield. The ease with which this measurement can be obtained has led to extensive use since it is recorded in almost all beef carcass research studies.

Cole et al. (1960), Brungardt and Bray (1963), Butler et al. (1961), Hedrick et al. (1963), Gottsch et al. (1961), Cahill et al. (1961) and Breidenstein (1962) have reported significant, positive, correlation coefficients (0.40 to 0.60) between 1. dorsi muscle area and retail yield. However, these same authors showed that on a carcass weight and fat constant basis the correlation coefficients were significantly reduced. Since area of 1. dorsi muscle is at least partially a function of weight, this would be expected. Field et al. (1963) reported that although the area of 1. dorsi muscle-weight relationship existed, it was a nonlinear relationship in their study. Brungardt and Bray (1963) reported that 20% of the variation in retail yield could be accounted for by differences in area of the 1. dorsi muscle. They also reported that with carcass weight, percent kidney fat and a single fat thickness measurement at the 12th rib held constant, the standard partial regression coefficient of boneless, closely trimmed retail yield from the round, loin, rib and chuck on area of the 1. dorsi muscle was only 0.16.

Briskey and Bray (1964) concluded that although the influence of area of  $\underline{1}$ .  $\underline{dorsi}$  muscle upon retail yield is small compared to that of fat,

emphasis upon size of this muscle may be justified because it comprises a large proportion of two of the high priced cuts of the beef carcass.

<u>Influence of Bone Upon Retail Yield</u>. Wythe <u>et al</u>. (1961) and Henderson <u>et al</u>. (1966) reported positive relationships between the amount of bone and retail yield.

Wythe et al. (1961) studied the weight, length, and length-weight ratio of the metacarpus, metatarsus, tibia, femur and the radius-ulna in relation to the yield of trimmed retail round, loin, rib and chuck. They observed significant (P < .01) and positive relationships between bone measurements and retail yield. Henderson et al. (1966) reported significant correlations (P < .01) between percent total carcass bone and percent total retail yield (0.64) and percent total retail yield of the four major wholesale cuts (0.68). Good et al. (1961) reported significant correlation coefficients between circumference of cannon bone and muscling score (-.32), circumference of round (0.30) and 1. dorsi area (0.13).

Regression Equations for Measuring Differences in Retail Yield. Murphy et al. (1960) developed a regression equation from data of 169 cattle varying widely in grade and weight. This equation was developed to predict the estimated percentage of boneless, closely trimmed retail cuts from the round, loin, rib and chuck. They compared predicted retail yield with actual retail yield and obtained a correlation of 0.906 with a standard error of the estimate of 1.9%. Palmer et al. (1961) conducted studies with 138 cattle varying in slaughter weight, breeding and quality grade. They reported a correlation coefficient between actual and estimated retail yield (Murphy et al., 1960) of 0.76 which accounted for slightly less than 60% of the total variation in retail cut yield.

Comparison of actual boneless, trimmed retail yields with those estimated by the equation of Murphy et al. (1960) ranges from a low of 35% of the variation in retail yield accounted for to a high of 80%, Briskey and Bray (1964). In a review of the literature, the latter authors stated that the equation of Murphy et al. (1960) probably accounts for 50 to 70% of the variation in the boneless, trimmed retail yield from the four major wholesale cuts.

Brungardt and Bray (1963) developed a regression equation designed to measure variation in percent boneless, trimmed retail yield from the round, loin, rib and chuck. Using this equation, they were able to account for 81% of the variation in boneless, trimmed retail yield.

Kropf and Graf (1959) and Breidenstein (1962) reported marked retail yield differences between steer and heifer carcasses and the latter author developed a separate regression equation for each sex. He was able to account for 72% of the variation in retail yield of steer carcasses but that for heifers, although similar, was slightly less accurate.

### EXPERIMENTAL PROCEDURE

Source of Material: Eighty steer carcasses were purchased from several beef packing companies in central Michigan. Selection was made pre-slaughter to insure procurement of steers of the three major British beef breeds. Carcass selection involved the following two criteria:

1) chilled carcass weight and 2) average fat thickness (12th rib measurement) as described by Naumann (1952).

Grouping: In an attempt to minimize the effects of carcass weight and fat thickness upon composition and cutability, the carcasses were equally divided into two weight ranges; 500 to 550 pounds and 700 to 750 pounds. The two weight ranges were further subdivided into four fat thickness (av. at 12th rib) ranges: .26 to .50 in., .51 to .75 in., .76 to 1.0 in. and 1.01 to 1.25 in. Ten steers were selected within each of the resulting eight groups as shown in table 1.

Table 1. Distribution of carcasses within weight and fat thickness groups.

	Av	erage fat thic	kness (12th ri	b)
Carcass weight	.26" to .50"	.51" to .75"	.76" to 1.0"	1.01" to 1.25"
	Group I	Group II	Group III	Group IV
500 to 550 lb.	10 carcasses	10 carcasses	10 carcasses	10 carcasses
	Group V	Group VI	Group VII	Group VIII
700 to 750 lb.	10 carcasses	10 carcasses	10 carcasses	10 carcasses

Slaughtering Procedure. Conventional slaughter procedures were followed except the carcasses were not scribed or the necks pinned.

Subjective Carcass Evaluation. Following selection of the carcasses, each was subjectively scored for the characteristics shown in table 2.

Conformation was scored with a conscious effort to evaluate degree of muscling irrespective of quantity of fat.

#### Linear Carcass Measurements

Linear Fat Measurements at the 12th Rib. The fat thickness measurements obtained at the 12th rib are shown in Figure 1. In addition to average fat thickness as previously described (av. of A, B and C), four other fat thickness measurements were taken at the 12th rib (D, E, F and G). The site for measurement D was determined by extending a perpendicular from the ventral tip of line W-X to the fat seam over the 1. dorsi muscle. A perpendicular line was drawn to the outer edge of the subcutaneous fat from the point at which the previous perpendicular bisected the fat seam. Measurement of the latter perpendicular was recorded as fat measurement D. Line Y-Z which is the same length as line W-X, was drawn from the point where the perpendicular of measurement D bisected the fat seam to the fat seam at point Z. Measurements E, F and G were made from perpendiculars drawn from the outer edge of subcutaneous fat to points located on the fat seam at one-fourth, one-half and three-fourths the length of line Y-Z.

<u>Fat Probes</u>. Probes of subcutaneous fat were made on the left side of each carcass at the sites shown in Figure 2. The following vertebrae

Table 2. Characteristics and scores used in the subjective carcass evaluation.

1 4 7 10 - 1	ore  Av.  2 5 8 11  Av. 2	High 3 6 9 12 +	a — a — a — b — c — a — d — d
Sc Low 1 4 7 10 -	Av. 2 5 8 11	3 6 9 12	a a b c a
Sc Low 1 4 7 10	Av. 2 5 8 11	3 6 9 12	a b c a
1 4 7 10 - 1	Av. 2 5 8 11	3 6 9 12	b c a
1 4 7 10 - 1	Av. 2 5 8 11	3 6 9 12	c a
1 4 7 10 - 1	Av. 2 5 8 11	3 6 9 12	a
1 4 7 10 - 1	Av. 2 5 8 11	3 6 9 12	
1 4 7 10 - 1	Av. 2 5 8 11	3 6 9 12	u
1 4 7 10 - 1	Av. 2 5 8 11	3 6 9 12	
1 4 7 10	2 5 8 11	3 6 9 12	
4 7 10 -	5 8 11	6 9 12	
4 7 10 -	5 8 11	6 9 12	
7 10 	8 11	9 12	
10	11	12	
<u>-</u> 1			
_	Av.	+	
_			
	2	3	
4	5	6	
7	8	9	
_	Av.	+	
1	2	3	
4			
7	8	9	
10	11	12	
13	14	15	
16	17	18	
19	20	21	
22	23	24	
25	26	27	
28	29	30	
	7 10 13 16 19 22 25	4 5 7 8 10 11 13 14 16 17 19 20 22 23 25 26	1       2       3         4       5       6         7       8       9         10       11       12         13       14       15         16       17       18         19       20       21         22       23       24         25       26       27




Figure 1. Illustration showing fat thickness measurements taken at the  $12\,\mathrm{th}$  rib.




Figure 2. Illustration showing the pattern of the fat probes taken 4, 8 and 12 in. from the midline on the carcass.

were used as skeletal reference points for the location of these sites:

The 5th, 8th and 1lth thoracic, the 1st, 4th and 6th lumbar and the 3rd and 5th sacral. Fat probes were made with a scalpel and metal ruler perpendicular to the anterior edge of these reference vertebrae, 4, 8, and 12 in. laterally from the dorsal tip of the vertebral cartilage. All probes were recorded to the nearest mm.

Length and Circumference of Round. These measurements were made with a flexible steel tape in accordance with the procedures described by Naumann (1952).

Depth at 10th and 12th Thoracic Vertebrae. These measurements were taken with a sliding T-square as described by MacLeod (1964).

Depth of Fat over the Brisket. This measurement was taken, with a sliding T-square, perpendicular to the first sternebra as shown in Figure 3.

## Cutting Procedure

Both right and left side of each carcass were cut into conventional wholesale cuts according to the procedure described by Wellington (1953), with the following two exceptions. The plate was removed from the wholesale rib cut by measuring 10 in. ventrally from the spinal column, at both the loin and blade ends of the rib, and removing the plate along a line connecting these two points. The brisket was removed from the chuck, by extending the cut where the foreshank was removed, on a line parallel with the dorsal side of the chuck.



Figure 3. Illustration showing the measurement of fat depth over the brisket.

Left Side. The wholesale cuts from the left side were individually physically separated into the components: muscle, fat and bone with two exceptions. The wholesale round and rib were further sub-divided. The rump was removed from the round according to the procedure described by Wellington (1953), and the wholesale rib was cut into three sub-parts; the 6-7-8 rib section, the 9-10-11 rib section and the 12th rib section following the procedure described by Hankins and Howe (1946). All three rib sections and the rump and rumpless round were then individually separated into muscle, fat and bone. All physically separated components were weighed and recorded to the nearest 0.05 lb.

Bone and Muscle Study. Weights of the following individual and bone groups from the left side were recorded to the nearest 0.05 lb.: radius plus ulna, humerus, scapula, tibia plus fibula and the femur.

Muscle weights were recorded for the following individual and muscle groups: semitendinosus, semimembranosus, biceps femoris, rectus femoris, triceps brachii, supraspinatus, psoas major, longissimus dorsi, semimembranosus plus adductor, and the quadriceps muscles.

Right Side. The four major wholesale cuts from the right side (round, rib, loin and chuck) were trimmed of external fat to approximately 0.3 in. Weight of untrimmed and trimmed wholesale cut and fat trim from each was recorded to the nearest 0.05 lb. All wholesale cuts were then cut into closely trimmed (approximately 0.3 in. external fat) boneless retail cuts. Weight of retail cuts, fat trim, and bone from each wholesale cut was recorded to the nearest 0.05 lb.

#### Tenderness Studies

The section of the left 1. dorsi muscle from the shortloin of each carcass was frozen and stored at approximately -10°F for subsequent chemical and organoleptic analysis. The frozen muscles were removed from the freezer 24 hr. prior to organoleptic analysis and four, 1 1/4 in. steaks were removed from the anterior end of the muscle and numbered 1 through 4 with 1 designated the most anterior steak. Steak number 1 was rewrapped and replaced in the freezer for subsequent chemical analysis. Steaks 2, 3 and 4 were then allowed to thaw overnight at approximately 38°F.

A thermometer was placed in the steak through the dorsal edge of the muscle to the approximate center of the cross-sectional area of the steak. The steaks were cooked in deep fat (lard) at a temperature of 300 ± 2°F. The steaks were removed and allowed to cool for 20 min. Four, one inch cores were removed from each steak parallel to the muscle fibers. The four cores from steak number 2 were used for Warner-Bratzler shear values. One shear value was determined from each core and the four values were averaged and recorded as shear value for each steak. Cores from steaks 3 and 4 were cut into two equal portions perpendicular to the orientation of the muscle fibers and served to a trained, eight-member taste panel. Evaluation of flavor, juiciness and tenderness was scored by the panel on a nine-point hedonic scale with nine designated as most desirable. Position of the cores within each steak was kept constant for each panel member.

# Chemical Analysis

Steak number 1 was removed from the freezer, allowed to thaw overnight at 38°F and then ground to obtain a sample for chemical analysis as outlined by 0rme (1958). Percent moisture and protein were determined by procedures as outlined by Benne et al. (1956). Ether extract was determined by the method described by 0rme (1958).

Potassium and sodium analysis were determined on each muscle sample using the TCA extraction procedure of Mounib and Evans (1957) as modified by Kirton and Pearson (1963). A Beckman DU spectrophotometer with a model 9220 flame attachment was used for the analysis. Potassium and sodium content of each muscle was calculated by use of a standard curve determined by plotting the percent transmittance against the ppm of these electrolytes in standard solutions.

## Statistical Analysis

The statistical procedures followed were described by Snedecor (1956). A 2 x 4 x 10 factorial analysis was calculated to determine treatment and interaction effects on several measures of carcass composition and cutability. The multiple range test of Duncan (1955) was employed to evaluate significance of individual differences. Simple correlation coefficients were calculated on a total, within weight and within weight-fat thickness group basis. Where applicable, multiple correlation coefficients were calculated on a total and within weight group basis.

#### RESULTS AND DISCUSSION

The means and standard deviations for the carcass traits which were correlated with all objective and subjective measurements are presented in tables 3, 3a, 4 and 4a. These means are presented for the combined and individual weight groups, within fat thickness ranges and within weight and fat thickness groups.

Effect of Carcass Weight upon Pounds and Percent Carcass Separable Components, Retail and Fat Trim Yields. Carcass weight had a highly significant (P < .01) effect upon pounds of all the carcass separable components, retail and fat trim yields (table 3). Carcass weight also had a highly significant (P < .01) effect upon percent retail and fat trim yields but not on the percent separable carcass fat, muscle and bone (table 4).

It is interesting to note that the 200 lb. increase in carcass weight from the 500 to 550 lb. weight group to the 700 to 750 lb. weight group resulted in an 88 lb. increase in separable muscle and a 94 lb. increase in total retail yield. Bone accounted for approximately 20 lb. of the total 200 lb. difference in carcass weight. Fat accounted for the remaining portion of this weight difference. Carcasses of the 500 to 550 lb. weight group were obtained from steers weighing approximately 850 lb. alive and those from the 700 to 750 lb. group from 1150 lb. steers. Thus, less than one-third of this live weight difference between these two carcass weight groups was separable muscle or total retail cut yield.

It is also of interest that the percents separable fat, muscle and bone as well as percents retail and fat trim yields between the two carcass weight groups (table 4) are markedly similar.

trim yields, 1. dorsi muscle area and average 12th rib fat thickness for the combined and individual weight groups and fat thickness ranges. Means and standard deviations of weights of carcass separable components, retail and fat Table 3.

Individu	ar weign	C groups a	individual weight groups and rat thickness ranges	ıcknessı	anges.	Dot this olonoon		
		weight	500-	<b>-00</b> 2		combined welcht	ness ranges eloht proms)	(8)
Trait		groups	550 1b.	750 lb.	0.26-0.50	0.51-0.75 0.76-1.00	0.76-1.00	1.01-1.25
Separable muscle,	Mean	159.60	137.57 <sup>a</sup>	181.64 <sup>b</sup>	171.35 <sup>c</sup>	161.01 <sup>d</sup>	154.34 <sup>d</sup> ,e	151.72 <sup>e</sup>
1b.	S.D.1	25.19	11.12	12.89	25.24	25.97	22.02	24.51
Separable fat,	Mean	113.78	94.93 <sup>a</sup>	132.63b	94.08c	111.54 <sup>d</sup>	123.01 <sup>e</sup>	126.47 <sup>e</sup>
1b.	S.D.	25.29	17.05	16.61	24.11	21.75	21.77	21.40
Separable bone,	Mean	35.21	30.32 <sup>a</sup>	40.11 <sup>b</sup>	38.75 <sup>c</sup>	36.47 <sup>c</sup>	33.19 <sup>d</sup>	32,43 <sup>d</sup>
1b.	S.D.	6.35	3.78	4.26	5.28	6.08	6.28	5,98
Retail yield from RLRC <sup>2</sup> , 1b.	Mean	158.24	138.52ª	177.97 <sup>b</sup>	164.14 <sup>c</sup>	160.57 <sup>c</sup>	156.46 <sup>c, d</sup>	151.81 <sup>d</sup>
	S.D.	22.46	10.31	10.82	20.03	20.94	21.78	26.34
Total retail	Mean	192.35	168.82a	215.88 <sup>b</sup>	199.63 <sup>c</sup>	195.93c	189.90c, d	183.92 <sup>d</sup>
yield, lb.	S.D.	27.24	13.72	13.38	23.16	24.67	25.95	33.41
Total fat trim,	Mean	62.54	49.58ª	75.50 <sup>b</sup>	48.27c	57,59 <b>đ</b>	68.25 <sup>e</sup>	76.04e
1b.	S.D.	19.49	16.53	12.33	18.65	18.88	16.50	11.67
External fat trim	Mean	11.73	8.27a	15.19 <sup>b</sup>	6.59c	9.47 <sup>c</sup>	12.96 <sup>d</sup>	17.90e
from RLRC, 1b.	S.D.		6.04	5.44	4.74	6.67	5.28	4.01
L. dorsi muscle area, sq. in.	Mean S.D.	10.93 1.67	10.00	11.86	11.62	10.79	10.73 1.06	10.58
Av. 12th rib fat thickness, in.	Mean S.D.	0.75	0.74	0.77	0.42	0.64	0.86	1.08
Carcass wt., 1b.	Mean S.D.	625.98 93.85	534.42 16.51	717.52	614.55 13.24	628.30 16.96	630.05 13.22	631.00 21.87
IS.D. = Standard deviation	eviation	١.			Round. loin.		chuck.	

LS,D. = Standard deviation. a,bMeans on the same line having different superscripts are highly significantly (P < .01) different

due to the effects of carcass weight.  $^{\rm c,d,e}_{\rm Means}$  on the same line having different superscripts are highly significantly (P < .01) different due to the effects of fat thickness.

		1000
		,

Means and standard deviations of weights of carcass separable components, retail and fat trim yields, 1. dorsi muscle area and 12th rib fat thickness for the individual fat thickness ranges within each weight group. Table 3a.

TUGINIGARI IRC		LUICKNESS	ranges		each weight group.	ic group.	;		
		-000	200-220 IB.	- 1	group	-00/	/00-750 Ib.	weight group	roup
		Group	Group	Group	Group	Group	Group	Group	$\operatorname{Group}$
		Н	II	III	ΙΛ	^	ΙΛ	VII	VIII
		0.26"-	0.51"-	0.76"-	1.01"-	0.26"-	0.51"-	0.76"-	1.01"-
Trait		0.50"	0.75"	1.00"	1.25"	0.50"	0.75"	1.00"	1.25"
•	;						1		
Separable muscle,	Mean	148.98	136.92	134.68	124./0	193.72	185.10	174.00	173.74
lb.	S.D.	10.92	6.70	6.54	10.26	10,65	9.47	11.05	9.24
Senarable fat	Mean	73.48	93.36	104.87	108.00	114.69	129,72	141,14	164,94
1b.	S.D.	12.26	9.35	9.54	10.82	11,55	13.29	13.36	9.54
Separable bone,	Mean	34.60	31.40	28.06	27.22	42.91	41.58	38.32	37.63
15.	s.D.	3.75	2.24	1.59	1.54	2,53	3.93	4.73	3.60
Retail vield from	Mean	146.78	141.90	137.44	127.96	181.50	179.25	175.48	175.66
RLRC <sup>2</sup> , 1b.	S.D.	7.55	5.26	9.74	8.21	10.99	11.07	10.11	11.52
Total retail	Mean	179.74	174.18	167.32	154.04	219.53	217.72	212.47	213.80
yield, 1b.	s.D.	7.76	7.47	13.21	10.76	13.86	13.27	10.73	16.02
•	;	0	•	1		•	6		,
Total fat trim,	Mean	32.02	•	55.96	68.44	64.52	73.29	80.54	83.64
1b.	S.D.	6.52	6.58	13.28	8.55	10.25	12.73	7.94	9.32
External fat trim	Mean	2.64	4,88	9.70	15.88	10.53	14.06	16.24	19.92
from RLRC, 1b.	S.D.	1.79	2.78	4.04	4.04	3.10	6.28	4.33	2.93
		· •		•	:	•	•	<u>:</u>	)
L.dorsi muscle	Mean	10.08	9.92	10.08	9.93	13.17	11.66	11.38	11.22
area, sq. in.	S.D.	1.63	1.83	09.0	1.14	1,39	1.31	1.38	0.88
:	;	•	;	•	•	•	,	•	,
Av. 12th rib fat	Mean	0.40	0.64	0.86	1.06	0.45	0.65	0.87	1.10
thickness, in.	S.D.	0.0	0.04	0.07	90.0	0.0	90.0	0.09	0.10
Carcass wt., 1b.	Mean	520.40	532,60	546.30	538.40	708.70	724.00	713.80	723.60
	S.D.	10.48	13.86	10.18	19.48	15.52	19.58	15.68	24.03
IS.D. = Standard deviation.	eviation		7	ZRLRC = R	= Round, loin,		rib and chuck	•	

Means and standard deviations of percent of carcass separable components, retail and fat trim yields for the combined and individual weight groups and fat thickness ranges. Table 4.

		Combined				Fat thickness ranges	ess ranges	
		weight	200-	700-		(combined weight groups)	ight groups	
Trait		groups	550 1b.	750 lb.	0.26-0.50	0.51-0.75	0.76-1.00	1.01-1.25
Separable muscle,	Mean	51.22	51.67 <sup>a</sup>	50.77ª	55.93 <sup>c</sup>	51.53d	49.10e	48.33e
. %	s.D.1	4.11	4.55	3,63	3,46	2.62	2.65	2.75
Separable fat, %	Mean	36.29	35.54a	37.04a	30.29c	35.59d	39.01e	40.27e
	S.D.	5.21	5.86	4.42	4.44	3.22	3,38	2.92
Separable bone, %	Mean	11.30	11.40a	11.20ª	12.70 <sup>c</sup>	11.69d	10.52 <sup>e</sup>	10.30e
•	S.D.	1.37	1.58	1.13	1.25	96.0	96.0	0.71
Retail_yield from	Mean	51.58	52.64a	50.53 <sup>b</sup>	54.72 <sup>c</sup>	52.22d	50.64 <sup>d</sup> , e	48.76e
RLRC <sup>2</sup> , %	S.D.	3.54	4.06	2.59	3,48	2.60	3.04	2.02
Total retail	Mean	62.72	64.15ª	61.28 <sup>b</sup>	66.60°	63.77d	61.48 <sup>d</sup> , e	59.02e
yield, %	S.D.	4.57	5.37	3.04	4.36	3.49	3.71	2.95
Total fat trim, %	Mean	20.11	18.78a	21.45 <sup>b</sup>	15.54c	18.32 <sup>d</sup>	21.92e	24.68 <sup>f</sup>
	S.D.	5.13	60.9	3.54	4.20	3.99	3.86	3.18
External fat trim	Mean	3.72	3.13a	4.31 <sup>b</sup>	2.04c	2.91c	4.12d	5.81e
from RLRC, %	S.D.	2.03	2.29	1.54	1.31	1.79	1.43	1.30

a, bMeans on the same line having different superscripts are highly significantly (P < .01) different due to the effects of carcass weight.  $c,d,e,f_{Means}$  on the sameline having different superscripts are highly significantly (P < .01) ZRLRC = Round, loin, rib and chuck. LS.D. = Standard deviation.

different due to the effects of fat thickness.

Means and standard deviations of percents of carcass separable components, retail and fat trim yields for the individual fat thickness ranges within each weight group. Table 4a.

		20	500-550 1b.	weight group	dn		700-750 1b.	weight group	۵
		Group		Group	Group	Group	Group	Group	Group
		Н	II	III	IV	Λ	VI	VII	VIII
		0.26"-	0.51"-	0.76"-	1.01"-	0.26"-	0.51"-	0.76"-	1.01"-
Trait		0.50"	0.75"	1.00"	1.25"	0.50"	0.75"	1.00"	1.25"
Serenation of Asserta		57 17	51 59	67 67	48 41	54.68	51 47	17 87	76 87
% S.D.1		3.66	2.61	2.61	3.46	2.90	2.76	2.77	1.99
Separable fat, %	Mean	28.22	35.12	38.51	40.29	32.36	36.05	39.51	40.26
	s.D.	4.75	3.00	3.28	3.43	3.10	3.52	3.57	2.49
Separable bone. %	Mean	13.28	11.83	10.31	10.17	12.11	11.55	10.72	10.44
	S.D.	•	0.92	09.0	0.58	99.0	0.98	1.22	0.83
Retail vield from	Mean	57.00ª	53,86b	51.40b, c,	d 48.28d	52,43b,c	50.58c,d	49.88c,d	49.23c,d
$RLRC^2$ , %	S.D.	•	1.21	3.56 1.98	1.98	2.48	2.62	2.37	2.04
Total retail	Медп	69 80 <b>8</b>	66.11b	62,57b,c		63.41b,c		60,39c,d	59,91c,d
yield, %	S.D.	2.88	1.90	4.73	2.79	3.03	3.16	2.00	2.98
Total fat trim %	Мев	12 428	15,928,b	20 92c.d	978 2C	18 67b.c	20 71c.d	22 91d,e	23 51d,e
o, (m., 151 151)	S.D.	2.43	2.55	4.91	3.11	3,11	3.78	2.28	2.95
External fat trim	Mean		1.86a,b	3,62c,d	6.02f	3.05 <sup>b</sup> , c	3,97c,d	4.62d,e,f	5.60e,f
from RLRC, % S.D.	S.D.	0.69	1.06	1.49	1.62	0.92	1.78	1.25	0.91

LS,D. = Standard deviation.  $^{2}RLRC = Round$ , loin, rib and chuck.  $^{3}S, ^{2}S, ^$ different due to the interaction of carcass weight and fat thickness. Effect of Fat Thickness upon Pounds and Percent Carcass Separable Components, Retail and Fat Trim Yields. Fat thickness had a highly significant (P < .01) influence upon pounds (table 3) and percent (table 4) carcass separable components, retail and fat trim yields. These results were as expected since the fatter carcass yielded significantly more separable fat, and fat trim than trimmer carcasses and lower yields of separable muscle, bone and retail cuts.

Even though the range within each of the four fat thicknesses was equal (0.25 in.) variation in yield (pounds and percent) of separable muscle, fat and bone was greater between the first two groups (0.26 to 0.50 in. and 0.51 to 0.75 in.) than between the second and third (0.50 to 0.75 in. and 0.76 to 1.00 in.) and was least between the third and fourth fat thickness ranges (0.76 to 1.00 in. and 1.01 to 1.25 in.). These data provide further evidence that 12th rib fat thickness is not a good indicator of total separable carcass fat. Similar findings were reported by Cole et al. (1962), Ramsey et al. (1962), Brungardt and Bray (1963), Hedrick et al. (1963), Lewis et al. (1964), Butterfield (1965), Carpenter et al. (1965) and Miller et al. (1965).

Carcass Weight and Fat Thickness Interaction upon Pounds and Percent Carcass Separable Components, Retail and Fat Trim Yields. Only the percent retail and fat trim yields (table 4a) were significantly (P < .01) effected by carcass weight and fat thickness interaction.

The mean 12th rib fat thickness of the eight carcasses within corresponding groups (I and V, II and VI, III and VII, and IV and VIII) (table

3a) are similar but percents separable muscle and bone were higher, while fat was lower, for the 500 to 550 lb. weight group than the corresponding fat thickness ranges of the 700 to 750 lb. weight group (table 4a). As previously discussed, these separable component (pounds and percent) differences were greatest between the first two fat thickness ranges (I and II, and V and VI) than between the third and fourth fat thickness ranges (III and IV, and VII and VIII). Pounds and percent total retail yield varied more widely within the 500 to 550 lb. weight group (approximately 25 lb. and 11.5%) than in the 700 to 750 lb. weight group (approximately 5.75 lb. and 3.2%) (tables 3a and 4a).

Relationships between Subjective Carcass Scores and Separable Components,

Retail and Fat Trim Yields

Correlation coefficients between subjective carcass scores and carcass separable components, retail and fat trim yields for the combined and individual weight groups appear in tables 5, 6 and 7. In general, correlations between these characteristics were very similar for pounds and percent comparisons.

Low negative correlation coefficients were found between subjective conformation scores and pounds and percents separable muscle and measures of retail beef yield in the combined weight groups and within the two weight groups (range, -.11 to -.38). Significant (P < .05) negative correlations were found between percent total carcass separable muscle and percent retail yields with marbling score and carcass grade in the combined weight groups and the 500 to 550 lb. weight group (range, -.39 to -.51).

Simple correlation coefficients between subjective carcass scores and carcass separable components, retail and fat trim yields for the combined weight groups. Table 5.

	Senerah 1a	eh le	Congrable	-ahla	Samerahla	ah1a	Roned trimmed	rimmed	Roned +rimmed	rimmed	Total fat	fat	Fotomal fat	1
Subjective	muscle	le	fa	fat	bone	a Dice	retail	retail RLRC1	retail yield	yield	trim	, F	trim from RLRC	om RLRC
score	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%
Conformation of:	of:													
Carcass	12	27	27 0.24	0.36	34	56	12	23	13	24	24 0.27	0.36	0,31	0.36
Hindquarter	10	25	0.22	0.33	30	51	07	17	09	18	0.22	0.29	0.24	0.27
Forequarter	03	29	29 0.32	0.39	26	55	05	26	07	-,31	0.36	0.42	0.41	0.44
Round	12	21	21 0.18	0.30	-,34	52	09	14	10	14	14 0.19	0.26	0.21	0.24
Carcass:														
Maturity	0.64	0.18	0.18 0.26	18	0.62	0.21	0.64	0.10	0.65	0.05 0.19	0.19	09	0.08	06
Marbling	0.09	39	39 0.47	0.48	12	58	0.10	39	0.08	43	0.47	0.46	0.37	0.37
Grade	06	45	45 0.42	0.54	0.5428	67	05	43	06	44	0,43	0.50	0.38	0.41
Correlations > 0 220 are along floant (D < 05)	0 220	0 40	font flo	d) tue	(50 /									

Correlations > 0.220 are significant (P < .05). Correlations > 0.287 are significant (P < .01). 1RLRC = Round, loin, rib and chuck.

Simple correlation coefficients between subjective carcass scores and carcass separable components, retail and fat trim yields for the 500 to 550 lb, weight group. Table 6.

Subjective	Separab	Separable muscle	Separab	rable :	Separable bone	able e	Boned trimmed retail RLRC1	rimmed RLRC1	Boned trimmed retail yield	rimmed vield	Total : trim	Total fat trim	External fat trim from RLRC	fat m RLRC
score	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%
Conformation of:	of:													
Carcass	10	22	22 0.36	0.3454		57	18	22	24	27	27 0.37	0.36	0.38	0.38
Hindquarter	09	20	20 0.33	0.30	48	52	09	16	14	21 0.31	0.31	0.30	0.31	0.30
Forequarter	14	26	26 0.40	0.37	51	55	23	28	27	-,31	31 0.40	0.40	0.44	0.43
Round	05	16	16 0.30	0.27	49	52	04	11	09	14	14 0.25	0.24	0.23	0.22
Carcass maturity	0.14	0.15	20	22	22 0.34	0.33	0.17	0.10	0.20	0.13	0.1310	13	17	18
Marbling	30	51	51 0.66	09.0	66	74	25	45	30	48	48 0.56	0.52	0.46	0.45
Carcass grade	32	51	51 0.66	0.62	0.62717828	78	28	44	4433	48	0.55	48 0.55 0.53 0.49	0.49	0.47
Country of 319 and almost thou	213		1001 640		10 / 02)									

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01).

RLRC = Round, loin, rib and chuck.

Simple correlation coefficients between subjective carcass scores and carcass separable components, retail and fat trim yields for the 700 to 750 lb. weight group. Table 7.

Subjective score	Separable muscle 1b. %	able	Separal fat 1b.	arable fat	Separable bone 1b. %	able e %	Boned trimmed retail RLRC1 1b. %	rimmed RLRC1 %	Boned trimmed retail yield 1b. %	rimmed yield %	Total trim 1b.	Total fat trim b. %	External fat trim from RLRC 1b. %	l fat om RLRC %
Conformation of:	of:													
Carcass	38	36	36 0.40	0.43	0.4354	55	27	24	25	24	24 0.38	0,40	0.37	0.38
Hindquarter	31	35	35 0.41	0.40	47	51	15	20	13	18	18 0.34	0.33	0.29	0.29
Forequarter	37	31	0.35	0.40	57	56	37	25	39	30	30 0.42	0.48	07.0	0.44
Round	38	32	32 0.34	0.39	55	53	24	20	22	20	20 0.35	0.37	0.33	0.34
Carcass maturity	0.54	0.45	- 38	45	45 0.37	0.33	0.54	0.51	0.54	0.54	0.5452	56	40	42
Marbling	25	21	0.20	0.25	3.2536	35	27	23	26	23	23 0.23	0.26	0.09	0.12
Carcass grade	36	-, 33	33 0.33	0.38	7.3847	47	36	35	33	32	32 0.36	0.40	0.21	0.24

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01).

RLRC = Round, loin, rib and chuck.

Correlation coefficients between marbling score and carcass grade with percents separable muscle and retail yields for the 700 to 750 lb. weight group were considerably lower than in the 500 to 550 lb. weight group (range, -.21 to -.35). The relationships between pounds of separable muscle and retail beef yields were also negative but in most instances lower than for percents, on both a combined and within weight group basis. These low relationships between conformation scores, carcass grade and marbling with retail yields support the findings of Cole et al. (1964) and Kirton (1964).

Positive correlation coefficients were observed between conformation scores and both weights and percents of separable carcass fat and fat trim yields in the combined and individual weight groups (range, 0.18 to 0.48). While carcass and forequarter conformation scores were more highly related to percent separable fat and fat trim yields than round and hind-quarter conformation scores, little difference in the magnitude of these correlation coefficients existed between the two weight groups and combined weight groups. These data indicate that conformation score is influenced by degree of fat even though a deliberate attempt was made in this study to evaluate degree of muscling without regard for fat. This was accomplished on the ribbed carcass after measuring fat thickness at the twelfth rib. These results substantiate the statement of Briskey and Bray (1964) that evaluation of conformation is influenced by fatness, especially in very fat carcasses. These results also agree with the findings of Cole et al. (1964) and Kirton (1964) and support the statement of

Hedrick et al. (1963) who stated that "for a carcass to have a high cutout, it is difficult to compensate for excess fat deposition with superior muscle development."

Carcass grade and marbling score were also positively correlated with carcass separable fat and fat trim yields in the combined and individual weight groups (range, 0.09 to 0.66). Highly significant (P < .01) correlations between these characteristics were found in the 500 to 550 lb. weight group (range, 0.45 to 0.66).

Carcass maturity score was positively related to measures of carcass muscle and bone (range, 0.10 to 0.64); whereas, it was negatively related to percent separable carcass fat and fat trim yields in the combined weight groups and to both pounds and percent separable carcass fat and fat trim yields within weight groups (range, -.06 to -.56).

Relationships Between Linear Fat Measurements and Separable Components,

Retail and Fat Trim Yields

Simple correlation coefficients between the fat probes and fat measurements at the 12th rib with pounds and percent separable components, retail beef and fat trim yields for the combined weight groups as well as within weight groups appear in tables 8, 9 and 10.

Correlation coefficients between fat measurements and percent separable components and retail yields were considerably higher, in most instances, than weight comparisons in the combined weight groups (table 8). This difference in magnitude of correlations was not observed for pounds and percent between fat trim yields and the linear fat measurements.

Simple correlation coefficients between linear fat measurements and carcass separable components, retail and fat trim yields for the combined weight groups. Table 8.

retail	and fat trim yields	for	the combined	combined weight groups	comps.					
	· ·			Boned,	را د	Boned,	d,		E	
	Separable muscle	Separable fat	Separable bone	trimmed retail RLR	$rac{ ext{mmed}}{ ext{RL} ext{RC}^1}$	trimmed retail yield	med yield	Total rat trim	External rat trim from RLRC	L rat om RLRC
Fat probes	1b. %	1b. %	1b. %	1b.	%	1b.	%	15. %	1b.	%
5th Thoracic										
4 in.			2576	0.00	64	01	63	0.65 0.69	0.64	9
8 in.	1242	97.0	2046	07	38	07	36	34	0.35	0.38
12 in.		0.54	0.0646	0.22	49	0.20	50	29	0.59	S
8th Thoracic										
4 in.		0.33 0.61	4462	30	52	30	51	41 0.	0.45	0.52
8 fn.	0458	0.56 0.60		0.03	56	0.04	52	0.58 0.57	0.62	09.0
12 in.	0.0755	0.60 0.59	0656	0.12	59	0.13	56	63 0.	0.67	0.63
11th Thoracic										
4 in.	2258	0.6		14	53	15	52	48	0.54	0.59
8 fn.	1063	0.56 0.66	2262	05	61	04	57	0.59 0.62	0.66	0.67
	ŧ	9.0		0.12	63	0.13	59	89	0.72	0.67
1st Lumbar										
4 in.	1160	0.53 0.63			2	-`0	- 60	0.59 0.64	99.0	0.68
	0957	0.50 0.60	1449	03	54	05	58	27	0.61	0.63
12 in.		0.75 0.64			66	0.23	66	76 0.	0.75	0.69
4th Lumbar		•				. ,				
4 fn.		0.53 0.58		0.00	- 60	0.00	58	o.	99.0	99.0
	2752	0.34 0.54	3350	23	52	22	48	0.42 0.52	0.47	0.51
12 fn.		0.61 0.61	0.0046	90.0	68	90.0	99	·	0.74	0.70
6th Lumbar										
4 fn.		9.0			60	0.12	61	68 0.	0.70	0.65
8 fn.	0961	0.52 0.61	1249	05	<b>-</b> .61	-`06	58	0.57 0.60	0.57	0.58
12 in.		0.4	80		47	0.18	49	57 0.	09.0	0.55

Simple correlation coefficients between linear fat measurements and carcass separable components, retail and fat trim yields for the combined weight groups. (continued) Table 8.

retall	retall and rat trim yleids	- 1	ror the combined weight groups.	Weignt 2	roups.	(continued)	uea)			
				Boned,	, td.	Boned,	ď,			
	Separable	Separable	Separable	trimmed	med	trimmed	med	Total fat	External	l fat
	sc1	fa	one	retail RLRC1	RLRC1	retail yield	yie1d	대	trim from	M RLRC
Fat probes	Ib. %	1b. %	1b. %	Ib.	%	Ib.	%	1b. %	Ib.	%
3rd Sacral										
4 tn	- 03 - 65	C	27 - 50 -	0	52	000	5.	0.57.0.53	0 63	85.0
		•			1 1		) L			
8 In.	79 10	0.28 0.28	0.0137	0.08	•	20.0	04	0.26 0.33	0.00	0.56
12 fn.	3961	·		35	57	34	52	0.37 0.56	0.44	0.53
5th Sacral										
4 fn.	0.0450	ö		0.04	55	0.04	52	52	0.53	0.50
8 in.	0.0458	0.59 0.56	0.0241	0.10	53	0.0	52	0.58 0.53	09.0	0.56
12 in.	0151	·		0.04	48	0.04	-,45	φ <b>τ</b>	0.53	0.49
12th Rib Fat										
Measurements										
¥	0753	o.		0.04	46	0.03	45	0.48 0.48	0.55	0.54
Ф	29 60	ö		24	55	24	54	0.46 0.61	0.51	0.58
ပ	3368	ö		26	60	26	57	0.51 0.66	0.59	99.0
Д	1462	0.53 0.64	2156	06	56	06	54	0.55 0.58	0.64	0.64
ы	1560	0		05	51	06	51	0.51 0.56	0.59	0.61
ĽΨ	1557	ö	2556	02	39	÷. 03	40	0.42 0.47	0.49	0.51
ტ	i	0.5'55	1355	0.10	44	0.0	43	0.52 0.50	0.57	0.54
Av. A, B, C.	2673	ö		16	78	17	64	0.59 0.71	0.68	0.72
Deptn or brisker fat	0.0652	0.56 0.55	0.0142	0.04	59	0.02	60	0.60 0.58	0.57	0.54
3		•		•						

Correlations > 0.220 are significant (P < .05). Correlations > 0.287 are significant (P < .01). IRLRC = Round, loin, rib and chuck.

Simple correlation coefficients between linear fat measurements and carcass separable components, retail and fat trim yields for the 500 to 550 lb. weight group. Table 9.

retail	and tat trim yleids	- 1	tor the 500 to 550 lb. Weight group	DO ID. We	TRUE RE	•	-			1
	01100000	01402000	Characa	boned,	α, 110,1	boned,	a, mod	Total fat	Twtorns1 fat	
	Separable	Separable fat	para bone	$ m retail~RLRC^1$	$R_{ m LRC}^{ m L}$	retail	Z, E	trim	trim from RLRC	ည္က
Fat probes	1b. %	1b. %	1b. %	1b.	%	1b.	%	1b. %	1b. %	
		C	5	0	. 72		7/2	0 77		
	0/ - 00 -	; ;	7 (		7/.	70.	† !	; c		
8 in.	•	0	5962	35	47	36	47	0.52 0.50	0.45 0.44	
12 fn.	5865	0.67 0.68	6464	56	<b>-</b> . 63	<b>-</b> .58	64	63 0.		
8th Thoracic										
4 in.	5258	ö	6465	47	58	50	60	63 0.	58 0.	
8 in.	5959	0.55 0.58	4846	54	56	53	55	0.53 0.54	0.58 0.58	
12 in.	5862	<u>.</u>		54	63	54	62	62 0.	65 0.	
11th Thorseic										
4 fn.	55 58	o	65 63	52	58	52	59	61 0.	63 0.	
	•	0.62 0.63	6162	54	62	52	- 60	0,60 0,60	0.62 0.61	
	4652	ö		54	60	53	59	60 0.	62 0.	
1st Lumbar										
4 in.		ö	6059	62	64	61	63	60 0.	0	
	i	0.68 0.68	6162	53	61	60	67	0.69 0.69	0.69 0.69	
12 in.		o.	6465	62	71	64	72	72 0.	0	
4th Lumbar										
4 in.	ı	47 0.	ï	59	57	56	55	52 0.	Ö	
8 fn.	5564	0.68 0.67	6063	57	67	-, 60	70	0.70 0.69	99	
12 fn.	i	67 0.	5655	67	74	99	73	68 0.	70 0.	
6th Lumbar										
4 fn.	5764	Ö	ï	54	62	56	-, 63	61 0.	58 0.	
		0.62 0.61	5658	54	62	-,49	57	0.59 0.58	0.55 0.54	
12 in.		ö	i	40	48	43	51	52 0.	52 0.	
										١

Simple correlation coefficients between linear fat measurements and carcass separable components, (continued) retail and fat trim yields for the 500 to 550 lb. weight group. Table 9.

	External fat	from	b. %		•	44	46 0.	.60 0.59				0.34 0.34					0.64 0.63					6
	Total fat Ext	rim	. % 1b.		77 0 77	44 0.44	49 0.51	65 0.64 0.			0.42	0,36		0, 60	0.70	0.75	0.58 0.57 0	0.58	0.51	0.50	0.84	( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( )
þ		į;	% 1b.		•	Š	Ö	67 0.			45 0.						58 0.			47		· ·
Boned	trin	retail yield	1b.		7.1	14	58	62		47	40	37		62	51	56	48	52				;
Boned,	trimmed	retail RLRCl	%					68			67 1			57			58					i
Bo		retai	1b.					63			43			59	550			67		940		Š
	Separable	bone	1b. %					3435			4445			5551	7276		5356	5252		60 58		
	Separable	fat	1b. %		c	ċ	ö	0.59 0.60		ö	0.53 0.55	ö		ö	ö	ö	0.63 0.62	Ö	0	ö	0	
	Separable	3c1	1b. %			•	6560	6264		60 58	5457	5054	urement	5048	5367	6071	5260	5356	5459	5657	6676	;
		,	Fat probes	3rd Sacral	7	4 In.		12 in.	5th Sacral	4 in.	8 fn.	12 in.	12th Rib Fat measurement	¥	Ф	ပ	Q	ы	Ŀ	9	Av. A, B, C	Depth of brisket

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01). IRLRC = Round, loin, rib and chuck.

Simple correlation coefficients between linear fat measurements and carcass separable components, retail and fat trim yields for the 700 to 750 lb. weight group. Table 10.

148331	Separable Separab	Separable fat	Separable bone	Boned, trimmed		Boned, trimmed	id ield	Total fat trim	External tron	nal fat from RLRC
Fat probes	1b. %	16. %	16. %		ı	1b.	%	1b. %		%
5th Thoracic	;	•				ć	Č	ì	i.	ï
4 in.	5959 - 31 - 36	0.61 0.64	6367		43	28	- 36	0.54 0.53	0.51	0.51
2 in. 12 in.	2934	; o	22 27		18	01	14	33	0.38	0.36
8th Thoracic										
4 fn.	6158	0.56 0.61	5759	. 65*-	57	45	52	0.61 0.61	0.59	0.60
8 fn.		0.58 0.61			51	28	43	63 <b>0.</b>	0.62	0.61
12 in.	4748	o.	4953		46	18	37	0.55 0.50	0.58	0.56
11th Thoracic										
4 fn.		0.61 0.59			49	14	45	62 0.	09.0	0.57
8 in.	67 66	0.64 0.69	6063		56	-,35	47	0.65 0.64	0.73	0.73
12 fn.		0.67 0.70			61	30	52	67 0.	0.71	0.70
1st Lumbar										
4 fn.	5764	0.66 0.66			63	-,34	54	0.73 0.69	0.76	0.75
8 in.		0.44 0.46	2930	-, 25	41	22	39		0.52	0.51
12 fn.	5464	0.64 0.62			49	22	47		0.59	0.57
4th Lumbar										
4 fu.	6462	0.59 0.63	5254	- 46	-,56	39	51	0.62 0.60	0.71	0.70
8 in.	4849	ö			53	27	44	99	0.53	0.52
12 in.	5051	·			57	38	<b>-</b> .53	66 0.	0.71	0.71
6th Lumbar										
4 in.	6668	ö		36	51	33	49	0.67 0.64	0.70	0.70
8 fn.	5460	0.57 0.57	2833		54	41	57	65 0.	0.59	0.59
12 fn.	4641	ö			31	27	29	36 0.	0.48	0.48

fficients between linear fat measurements and carcass separable components,	
nd carcas	Continue
: measurements a	1de for the 700 to 750 1h weight aroun (continued)
linear fat	750 18
between	-he 700 +
coefficients	a wielde for
correlation coef	otes 1 and for trim who
Simple	10+01
Table 10.	

retail	retail and tat trim yields for the /00 to /50 lb. weight group.	m yields tor	the /00 to	/50 Ib. W	elght g		(continued)	(par		
				Boned,	,	Boned,				
	Separable	Separable	Separable	trimmed	ped	trimmed	pe	Total fat	External	fat
•	8c1	fat	one	딥	뵚	retail yield	vield	Lrim	trim from RLRC	n RLRC
Fat probes	1b. %	1b. %	1b. %	Ib.	%	Ib.	%	1b. %	1b.	%
3rd Sacral										
4 in.	7068	0.64 0.69		4	52	39	-,49			7.71
8 in.	62 64	0.61 0.63	4550	26	4	22	40	0.55 0.50	0.63	0.61
12 in.	65 66	0.66 0.69		36	52	21	37			), 62
5th Sacral										
4 in.	4738	0.34 0.41	4340	54	46	51	45	0.45 0.51		.54
8 in.		0.56 0.58	3437	43	52	44	56	0.65 0.64		0.67
12 fn.	5245	0.44 0.50	5453	47	47	39	40	0.54 0.56	0.65	99.0
12th Rib fat										
measurements:										
A	5360	0.60 0.59	3745	07	33	02	28			7,44
Ф	5252	0.55 0.57	5761	32	44	28	45			. 63
U	6065	0.69 0.70	60 68	29	47	26	45			99.(
Д	67 64	0.62 0.66	5658	41	54	36	51			89.
ធ	6564	0.63 0.67	5962	29	43	25	39			.55
ഥ	5454	0.56 0.58	5356	12	24	10	21			.47
9	5857	0.56 0.59	5154	20	32	14	27	0.49 0.46	0.56	0.55
Av. A, B, C	6570	0.72 0.73	60 67	25	48	20	44			.67
Dept. of brisket										
fat	5046	0.42 0.46	3736	59	54	56	54	0.54 0.58	0.46	0.45
Correlations > 0.312 are significant (P	312 are ston	fficant (P <	05)							

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01).

1RLRC - Round, loin, rib and chuck.

These data suggest that linear fat measurements, in similar fat thickness groups but different weight groups, were similar in magnitude; whereas, the pounds separable components and retail beef yields were widely different between weight groups (table 3). However, fat trim yields were similar between weight groups and the relationships between the linear fat measurements and pounds and percent fat trim yields were more constant.

In contast to the combined weight groups, the within weight group (50 lb. weight range) correlation coefficients (tables 9 and 10) between linear fat measurements and pounds and percent separable components, retail and fat trim yields were similar. These differences in correlations between weights and percents on a combined weight group basis agree with the results of Orme (1958), Orme (1963), Allen (1963), Hedrick et al. (1963) and Miller et al. (1965).

The fat measurements most highly related to percent separable components and retail yields for the combined weight groups were: the 5th thoracic fat probe at four inches, fat measurement C over the 12th rib and the average of fat measurements A, B and C (range, -.57 to 0.78). These three linear measurements of fat also showed the highest relationship to both pounds and percent separable components and retail yields in the 500 to 550 lb. weight group (range, 0.69 to -.86). However, in the 700 to 750 lb. weight group, most of the relationships between these fat measurements were markedly reduced (range, -.36 to 0.73), but all correlations involving percents and most for weights remained highly significant. These findings support those of Lewis et al. (1964) who

reported significant (P < .05) correlations between fat probes and measurements and percent retail yield of the round, loin, rib and chuck. Since correlations were higher within the 500 to 550 lb. weight group than in the 700 to 750 lb. weight group, it appears that linear measurements of fat are more accurate indicators of separable muscle, fat, bone and retail yields for lighter carcasses than for heavier, fatter carcasses.

Within the 700 to 750 lb. weight group, the fat probes in the lumbar region were more highly correlated (range, -.22 to 0.76) with separable components than the same probes within the 500 to 550 lb. weight group (range, -.34 to -.74). These data indicate that in the heavier carcasses, covariances between fat probes in the lumbar region and separable components, retail and fat trim yields were greater in most instances than those in the 500 to 550 lb. weight group. Correlation coefficients between fat probes and measurements and fat trim yields for the combined weight groups as well as within weight groups were highly significant (P < .01) in most instances (range, 0.27 to 0.84).

Depth of brisket fat was negatively correlated (P < .01) with percent separable muscle, bone and retail yields for the combined weight groups (range, -.42 to -.60), but positively correlated with pounds of these components (range, 0.01 to 0.06). Within the two weight groups the depth of brisket fat and separable muscle and bone and retail yields were negative but significantly (P < .05) correlated when expressed as either pounds or percent. Depth of brisket fat was highly significantly correlated (P < .01) with pounds and percent separable fat and fat trim yields for the combined as well as the two individual weight groups (range, 0.42 to 0.63).

Relationships Between Some Linear Carcass Measurements and Separable

Components, Retail and Fat Trim Yields

Simple correlation coefficients between linear carcass measurements taken and separable components, retail and fat trim yields for the combined weight groups and for each weight group appear in tables 11 and 12. Correlation coefficients between depth of rib measurements and separable components, retail and fat trim yields were generally lower (range, 0.00 to 0.36) than those between other measurements and the same components. This was true in the combined weight groups as well as the two weight groups.

Length of round was highly correlated to pounds of separable muscle, bone and retail yields in the combined weight groups (0.82 to 0.92). However, these relationships were reduced when calculating correlations between this measurement and percentages or for pounds and/or percentages on a within weight group basis (range, 0.02 to 0.61), except for percent bone. In the case of bone, the correlation coefficients were still large enough to be useful for predictive purposes (range, 0.43 to 0.82). Length of round was negatively related to pounds and percent separable fat and fat trim yields within weight groups (range, -.47 to -.70). These findings are similar to those of 0rme (1963) who reported measurements of carcass length and length of fore or hind legs in lambs was negatively related to measures of carcass fat. Cole et al. (1962) also reported a negative relationship between length of carcass and separable carcass fat.

Circumference of round generally showed a higher relationship to measurements of carcass fat than to bone or carcass muscling for the

Simple correlation coefficients between some linear carcass measurements and carcass separable components, retail and fat trim yields for the combined weight groups. Table 11.

				Boned,	Boned,	ìd,			
	Separable muscle	Separable fat	Separable bone	$ ext{trimmed}$ $ ext{retail}$ $ ext{RLRC}^1$	trimmed trimmed:	ned yield	Total fat trim	External fat	External fat trim from RLRC
Linear measurement	1b. %	1b. %	1b. %	1b. %	1b.	%	1b. %	1b.	%
Depth 10th rib	0.2515	0.35 0.18	0.1521	0.3504	0.36	04	0.22 0.07	0.18	0.08
Depth 12th rib	0.2303	0.22 0.05	0.1806	0.33 0.06	0.32	0.04	0.13 0.00	90.0	02
Length of round	0.82 0.19	0.3723	0.92 0.43	0.83 0.02	0.82	0.03	0.2616	0.11	11
Circ. <sup>2</sup> of round	0.5329	0.69 0.32	0.4325	0.5640	0.55	40	0.66 0.39	0.58	0.42
L.dørsi muscle area	0.67 0.26	0.2517	0.4801	0.62 0.12	09.0	0.07	0.2005	0.10	02

Correlations > 0.220 are significant (P < .05). Correlations > 0.287 are significant (P < .01).

RELRC = Round, loin, rib and chuck.

Circ. = circumference.

ents between some linear carcass measurements and carcass separable compon-	lelds for the 500 to 550 lb, and 700 to 750 lb, weight groups.	
ass measurements	and 700 to 750 1	Boned,
some linear carca	500 to 550 lb.	Boned,
fficients between a	rim yields for the	
Table 12. Simple correlation coefficien	ents, retail and fat trim yie	
Table 12.		

				Boned,	Boned,		
	Separable	Separable	Separable	trimmed	t	Total fat	External fat
	muscle	£	pone	retail RLRC	retail yield	trim	trim from RLRC
Linear measurement	1b. %	1b. %	1b. %	1b. %	1b. %	16. %	1b. %
		5	00 to 550 lb.	500 to 550 lb. weight group	•		
		1			1		
Depth 10th rib	0.0903	0.12 0.08	1420		0.11	0.03 0.02	0707
Depth 12th rib	0.13 0.07	0.0003	0306		0.16	0.0002	
Length of round	0.36 0.48		0.80 0.82		0.53	7069	
Circ. <sup>2</sup> of round	1312	0.14 0.16	29 26	1313		0.22 0.23	0.22 0.22
L.dorsi muscle area	0.44 0.31	1318	1116		0.25	1214	
		71	00 to 750 lb.	700 to 750 lb. weight group	ol.		
Depth 10th rib	1224		1120		0.31	0.0306	0.13 0.09
Depth 12th rib	0.0413		0.0902		0.45	0415	0.0104
Length of round	0.52 0.43	4755	0.82 0.81	0.46 0.38		4853	4952
Circ. of round	4851		3034		32	0.50 0.47	
L.dorsi muscle area	0.45 0.50		0.25 0.30		0.31	5047	4140

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01).

RLRC = Round, loin, rib and chuck.

2Circ. = circumference.

combined weight groups. In the 500 to 550 lb. weight group circumference of round showed low relationships to both weight and percent separable components, retail and fat trim yields (range, -.12 to -.29).

L. dorsi muscle area was highly significantly (P < .01) correlated with pounds of separable muscle and retail yields for the combined weight groups (range, 0.60 to 0.67). Correlations between 1. dorsi muscle area and percent separable components and retail yields on a combined weight group basis as well as within weight group correlations were markedly lower. Cole et al. (1962), Field et al. (1963) and Henderson et al. (1966) show the same relationships between 1. dorsi muscle area and weight and percent measures of carcass muscling.

Multiple Regression Analyses of Objective Carcass Measurements on Separable

Components, Retail and Fat Trim Yields

Only those multiple correlations which were high enough to be useful for predictive purposes and include a practical number of independent variables will be reported in the results and discussion portion of this manuscript. Other regression analyses which were calculated will be found in the supplement.

Multiple Regression Analyses of Total Carcass Separable Fat on Objective Carcass Measurements. In the combined weight groups, 86% and 94% of the variation in percent and pounds, respectively, of separable carcass fat could be accounted for with a combination of carcass weight and seven probes and fat thickness measurements (suppl. p. 168).

In the 500 to 550 lb. weight group, 90% of the variation in weight of separable fat and 89% of the variation in percent separable fat could be accounted for using the combination of variables in tables 13 and 14.

Coefficient of determination between weight of carcass separable fat and a combination of objective carcass fat measurements, and the regression coefficient for each of the measurements (500 to 550 lb. weight group)

	Regression	Beta	Level of	R <sup>21</sup>
Independent variables	coefficients	weights	sig.	<u>deletes</u>
Constant	-143.153		0.00	
Carcass wt.	0.348	0.338	0.00	0.82
Probe-5th Thor4 in. <sup>2</sup>	1.826	0.445	0.00	0.81
Probe-6th Lumb8 in.3	0.383	0.200	0.01	0.88
Probe-6th Lumb12 in.	379	175	0.03	0.88
Fat depth over brisket	4.718	0.207	0.01	0.87
Fat measurement D	0.699	0.223	0.01	0.87

Standard error of estimate = 5.91

 $<sup>^{1}</sup>$ Value of the coefficient of determination if that variable was deleted from the analysis.

Thor. = Thoracic vertebra.

<sup>&</sup>lt;sup>3</sup>Lumb. = Lumbar vertebra.

Table 14. Coefficient of determination between percent carcass separable fat and a combination of objective carcass fat measurements, and the regression coefficient for each of the measurements

(500 to 550 lb. weight group).					
Independent variables	Regression coefficients	Beta weights	Level of sig.	R <sup>21</sup> deletes	
Independent Variables	COETITCIENCS	weights	SIE.	deletes	
Constant	-8.412		0.54		
Carcass wt.	0.045	0.127	0.09	0.88	
Probe-5th Thor4 in.2	0.742	0.538	0.00	0.78	
Probe-1st Lumb8 in.3	178	254	0.03	0.87	
Probe-3rd Sac12 in.4	0.180	0.271	0.00	0.84	
Fat depth over brisket	1.812	0.232	0.00	0.86	
Fat measurement D	0.400	0.371	0.00	0.83	

 $R^2 = 0.89$  Standard error of estimate = 2.10

In the 700 to 750 lb. weight group, 88% of the variation in pounds of separable carcass fat could be accounted for using nine variables (suppl., p. 169). These same nine variables were responsible for 89% of the variation in percent separable carcass fat (suppl., p. 169). These results indicate that accuracy can be attained with a combination of objective carcass fat measurements to predict either percent or pounds of separable carcass fat. The standard error of estimates, for the prediction of percent separable fat, in the combined weight groups, 500 to 550 lb. weight group and 700 to 750 lb. weight group, were 2.10, 2.10 and 1.66%, respectively. These compare favorably with the standard error of estimate

<sup>&</sup>lt;sup>1</sup>Value of the coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Thor. = Thoracic vertebra.

 $<sup>^{3}</sup>$ Lumb. = Lumbar vertebra.

<sup>&</sup>lt;sup>4</sup>Sac. = Sacral vertebra.

of 2.34% found by Hankins and Howe (1946) for their prediction equation for percent separable carcass fat from percent separable fat of the 9-10-11 rib section.

In the combined weight groups, using only the five variables listed in table 15, 92% of the variation in weight of total carcass separable fat was accounted for. It can be observed from this table that carcass weight has the greatest effect in this prediction equation. Using essentially the same variables minus carcass weight and including one more fat measurement, 81% of the variation in percent carcass separable fat was accounted for (table 16). Using only these five independent variables

Table 15. Coefficient of determination between weight of total carcass separable fat and a combination of objective carcass fat measurements, and the regression coefficient for each of the measurements (combined weight groups).

measurements	(combined weight	groups).		
	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	-35.266		0.00	
Carcass wt.	0.159	0.590	0.00	0.68
Probe-5th Thor4 in. <sup>2</sup>	1.686	0.271	0.00	0.87
Probe-1st Lumbar-12 in.	0.424	0.123	0.02	0.91
Probe-Av. 3rd Sacral <sup>3</sup>	0.914	0.178	0.00	0.90
Fat measurement D	0.588	0.133	0.00	0.91

 $R^2 = 0.92$  Standard error of estimate = 7.58

<sup>1</sup> Value of the coefficient of determination if that variable was deleted from the analysis.

 $<sup>^{2}</sup>$ Thor. = Thoracic vertebra.

<sup>3</sup>Av. of 3 probes taken at 4, 8 and 12 in. off the carcass dorsal midline.

Table 16. Coefficient of determination between percent carcass separable fat and a combination of objective carcass fat measurements, and the regression coefficient for each of the measurements (combined weight groups).

(combined weight groups).					
Independent variables	Regression coefficients	Beta weights	Level of sig.	R <sup>21</sup> deletes	
Constant	21.678		0.00		
Probe-5th Thor4 in.2	0.626	0.488	0.00	0.67	
Probe-8th Thor8 in.	282	262	0.00	0.79	
Probe-1st Lumbar-12 in.	0.134	0.189	0.01	0.79	
Probe-Av. 3rd Sacral3	0.393	0.372	0.00	0.74	
Fat measurement D	0.276	0.303	0.00	0.77	

 $R^2 = 0.81$  Standard error of estimate 2.34

resulted in a standard error of estimate of 2.34% carcass separable fat, which is the same standard error of estimate reported by Hankins and Howe (1946) using their prediction equation for carcass separable fat involving separable fat of the 9-10-11 rib section. This same multiple regression analysis in the 500 to 550 lb. weight group was responsible for 85% of the variation in percent carcass separable fat with a standard error of estimate of 2.46% (suppl., p.170). In the 700 to 750 lb. weight group, 83% of the variation in percent carcass separable fat could be accounted for using these five fat measurements with a standard error of estimate of 1.93% (suppl., p. 170).

<sup>&</sup>lt;sup>1</sup>Value of the coefficient of determination if that variable was deleted from the analysis.

<sup>2</sup>Thor: = Thoracic vertebra.

<sup>&</sup>lt;sup>3</sup>Av. of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

Multiple Regression Analyses of External Fat Trim from the Round, Loin, Rib and Chuck on Objective Carcass Measurements. In the combined weight groups, 79% of the variation in external fat trim from the round, loin, rib and chuck (RLRC) was accounted for using two fat probes, carcass weight and length of round (table 17). It can be observed from the data in this table that weight is the most important variable in the regression equation. These same four variables, in the combined weight groups, accounted for 71% of the variation in percent external fat trim from the RLRC (suppl., p. 171).

Table 17. Coefficient of determination between weight of external fat trim from the round, loin, rib and chuck and a combination of objective carcass fat measurements, and the regression coefficient for each of the measurements (combined veight groups)

<u>clent for each</u>	of the measure	ments (comb	ined weight	
	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	11.933		0.08	
Carcass wt.	0.050	0.700	0.00	0.64
Probe-4th Lumbar-12 in.	0.320	0.332	0.00	0.73
Probe Av. 3rd Sacra12	0.416	0.307	0.00	0.73
Langth of round	-1.498	434	0.00	0.73

 $R^2 = 0.79$  Standard error of estimate = 3.13

In the 500 to 550 lb. weight group, 83% of the variation in the external fat trim from these same four wholesale cuts could be accounted for

<sup>&</sup>lt;sup>1</sup>Value of the coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Av. of 3 probes taken at 4, 8 and 12 in. off the carcass dorsal midline at the 3rd Sacral vertebra.

using the independent variables shown in table 18. It should be noted from the data in this table that when the analysis was calculated within weight groups, carcass weight had little effect upon the weight of external fat trim from the RLRC. Ninety-three percent of the variation in percent external fat trim from the RLRC in the 500 to 550 lb. weight group could be accounted for using 15 variables, all of which had a significant (P < .05) effect upon percent fat trim (suppl., p. 171).

Table 18. Coefficient of determination between weight of external fat trim from the round, loin, rib and chuck and a combination of objective carcass fat measurements, and the regression coefficient for each of the measurements (500 to 550 lb. weight

group).				
Independent variables	Regression coefficients	Beta weights	Level of sig.	R <sup>21</sup> deletes
Constant	-8.324		0.62	
Carcass wt.	0.004	0.10	0.88	0.83
Probe-3rd Sacral-12 in.	0.199	0.290	0.00	0.77
Fat measurement A	0.238	0.297	0.01	0.79
Fat measurement C	0.509	0.494	0.00	0.74
Fat measurement E	0.367	0.526	0.00	0.77
Fat measurement F	320	490	0.01	0.79

 $R^2 = 0.83$  Standard error of estimate = 2.71  $^1$ Value of coefficient of determination if that variable was deleted from the analysis.

In the 700 to 750 lb. weight group, 96% of the variation in pounds or percent of external fat trim was accounted for using the 22 variables given in the supplement p. 172.

Multiple Regression Analyses of Total Retail Fat Trim on Objective Carcass Measurements. The six independent variables given in table 19 were responsible for 87% of the variation present in weight of total retail fat trim for the combined weight groups. Eighty-one percent of the variation in percent of this same trait could be accounted for using the 12 independent variables presented in the suppl., p. 173.

Table 19. Coefficient of determination between weight of total retail fat trim and a combination of objective carcass fat measurements, and the regression coefficient for each of the measure-

ments (combined weight groups). R21 Level of Regression Beta Independent variables coefficients weights sig. deletes Constant -2.572 0.87 Carcass wt. 0.158 0.762 0.00 0.72 Probe-4th Lumbar-12 in. 0.01 0.438 0.156 0.85 Probe-Av. of 3rd Sacral<sup>2</sup> 0.709 0.179 0.00 0.85 Fat depth over brisket 0.169 0.00 3.810 0.85 Length of round -2.748 -.273 0.00 0.85 0.02 Fat measurement C 0.562 0.161 0.86

Within the 500 to 550 lb. weight group, the coefficient of determination for weight of total retail fat trim was 0.85 using the five independent variables presented in table 20. Eighty-four percent of the variation

 $R^2 = 0.87$  Standard error of estimate = 7.41

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Av. of 3 probes 4, 8 and 12 in. off the carcass dorsal midline at the 3rd Sacral vertebra.

Table 20. Coefficient of determination between weight of total retail fat trim and a combination of objective carcass fat measurements, and the regression coefficient for each measurement

(500 to 550 lb. weight group).				
	Regression	Be <b>ta</b>	Level of	R21
Independent variables	coefficients	weights_	sig.	<u>deletes</u>
Constant	42.998		0.51	
Carcass wt.	0.119	0.119	0.15	0.84
Probe-Av. 3rd Sacral <sup>2</sup>	0.725	<b>0.3</b> 85	0.00	0.73
Length of round	-3.243	237	0.02	0.83
Fat measurement A	0.663	0.302	0.00	0.79
Fat measurement B	0.432	0.282	0.01	0.82

 $R^2 = 0.85$  Standard error of estimate - 6.80

in percent total retail fat trim could be accounted for in the 500 to 550 lb. weight group (table 21) using five carcass measurements. Carcass weight showed little effect on either pounds or percent total retail fat trim within weight groups.

In the 700 to 750 lb. weight group, 80% of the variation in weight of total retail fat trim was accounted for using eight variables (suppl., p. 173). Using six variables (suppl., p. 174), 73% of the variation in percent total retail fat trim could be accounted for in this weight group.

These results indicate that estimating yields of separable fat or fat trim from objective measurements was more difficult in the 700 to 750 lb. weight group than in the 500 to 550 lb. carcasses. However, comparable

<sup>1</sup> Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Av. of 3 probes 4, 8 and 12 in. off the carcass dorsal midline at the 3rd Sacral vertebra.

Table 21. Coefficient of determination between percent total retail fat trim and a combination of objective carcass fat measurements, and the regression coefficient for each of the measurements

(300 20 330 1	<ul><li>b. weight group)</li><li>Regression</li></ul>	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes_
Constant	34.189		0.16	
Carcass wt.	0.012	0.032	0.70	0.84
Probe-3rd Sacral-12 in.	0.272	0.393	0.00	0.72
Length of round	-1.246	247	0.02	0.82
Fat measurement A	0.264	0.327	0.00	0.77
Fat measurement B	0.168	0.296	0.01	0.80

R<sup>2</sup> = 0.84 Standard error of estimate = 2.58

<sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

accuracy to the 500 to 550 lb. carcasses could be attained in the 700 to 750 lb. weight group by increasing the number of variables in most instances.

Multiple Regression Analyses of Total Separable Carcass Muscle on Objective Carcass Measurements. For the combined weight groups, 93% of the variation in weight of total separable carcass muscle could be accounted for using the five independent variables listed in table 22. As can be seen from the data in this table, carcass weight was the most important variable in predicting separable carcass muscle. This agrees with the findings of Cole et al. (1962). Carcass weight, 5th thoracic fat probe 4 in. off the carcass dorsal midline, average of 3 fat probes at the 3rd sacral vertebra

Table 22. Coefficient of determination between weight of separable carcass muscle and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (combined weight groups).

	<del></del>	7 1 6	R21
coefficients	вет <b>а</b> weights	sig.	deletes_
25.362		0.00	
0.241	0.896	0.00	0.49
779	126	0.01	0.92
-1.175	230	0.00	0.91
501	111	0.01	0.93
1.407	0.093	0.03	0.93
	Regression coefficients  25.362  0.241 779  -1.175 501	Regression Beta weights  25.362  0.241  0.896 779 126  -1.175 230 501 111	coefficients     weights     sig.       25.362     0.00       0.241     0.896     0.00      779    126     0.01       -1.175    230     0.00      501    111     0.01

 $R^2 = 0.93$  Standard error of estimate = 6.78

and fat measurement C were responsible for 73% of the variation in percent separable carcass muscle (suppl., p. 174). These data indicate that carcass weight contributed much less to determining percent separable carcass muscle than to weight.

In the 500 to 550 lb. weight group, three variables accounted for 71% of the variation in weight of total separable carcass muscle (suppl., p. 174). These same three variables accounted for 78% of the variation in percent separable carcass muscle (table 23). The multiple correlation coefficient between these three objective measurements and percent separable carcass muscle (0.88) compares favorably with the correlation of 0.85 between percent separable muscle of the 9-10-11 rib section and percent separable carcass muscle determined by Hankins and Howe (1946).

<sup>1</sup> Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Av. of 3 probes 4, 8 and 12 in. off the carcass dorsal midline at the 3rd sacral vertebra.

Table 23. Coefficient of determination between percent separable carcass muscle and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (500 to 550 lb. weight group).

to 550 lb. wei	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	<b>8</b> 6.511		0.00	
Carcass wt.	<b>04</b> 6	168	0.08	0.76
Probe-5th thoracic-4 in.	436	407	0.00	0.70
Probe-Av. 3rd sacral2	437	491	0.00	0.63

 $R^2 = 0.78$  Standard error of estimate = 2.20

In the 700 to 750 lb. weight group, low coefficients of determination were observed for pounds and percent separable carcass muscle using objective carcass measurements as the independent variables. Three variables; l) carcass weight, 2) probe 4 in. off the carcass dorsal midline at the 5th thoracic vertebra (figure 2), and 3) the average of 3 probes taken at the 3rd sacral vertebra were responsible for 68% of the variation in pounds and 65% of the variation in percent separable carcass muscle (suppl., p. 175).

These data indicate that it was more difficult to predict pounds or percent separable carcass muscle in the 700 to 750 lb. weight group than in the 500 to 550 lb. carcasses. This may possibly be attributed to greater intermuscular fat deposition in the 700 to 750 lb. weight group than in the 500 to 550 lb. carcasses.

<sup>1</sup> Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Av. of 3 probes 4, 8 and 12 in. off the carcass dorsal midline at the 3rd Sacral vertebra.

Multiple Regression Analyses of Total Carcass Separable Bone on Objective Carcass Measurements. In the combined weight groups, 91% of the variation in pounds of separable carcass bone could be accounted for using the independent variables listed in table 24. This equation is a practical and accurate method for prediction of separable carcass bone as it employs four easily taken measurements and has standard error of estimate of 1.94 lb.

Table 24. Coefficient of determination between weight of total carcass separable bone and a combination of objective carcass measurements, and the regression coefficient for each of the measurements.

ments (combined weight groups).				
	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	-21.423		0.00	
Carcass wt.	0.032	0.468	0.00	0.87
Probe-5th Thoracic-4 in.	292	187	0.00	0.90
Length of round	1.499	0.457	0.00	0.87
Fat measurement C	197	173	0.00	0.90

R<sup>2</sup> = 0.91 Standard error of estimate = 1.94

1 Value of coefficient of determination if that variable was deleted from the analysis.

Using these same four variables, 80% of the variation in percent separable carcass bone was accounted for in the combined weight groups (table 25).

Table 25. Coefficient of determination between percent carcass separable bone and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (com-

bined weight	groups).			
Independent variables	Regression coefficients	Bet <b>a</b> weights	Level of sig.	R <sup>21</sup> deletes
Constant	4.209		0.02	
Carcass weight	008	575	0.00	0.73
Probe-5th Thoracic-4 in.	103	306	0.00	0.76
Length of round	0.503	0.712	0.00	0.70
Fat measurement C	063	<b>2</b> 56	0.00	0.77

 $R^2 = 0.80$  Standard error of estimate = 0.63% <sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

In the 500 to 550 lb. weight group, 78% of the variation in pounds of separable carcass bone was accounted for using carcass weight, 5th sacral fat probe - 4 in. (figure 2) and the length of round measurement (suppl., p. 175. These same three variables accounted for 83% of the variation in percent separable carcass bone in the 500 to 550 lb. weight group (suppl., p. 176). Carcass weight, average of 3 fat probes at the 3rd sacral (figure 2) and length of round accounted for 80% of the variation in pounds of separable carcass bone (suppl., p. 176) and 77% of the variation in percent separable carcass bone (suppl., p. 176) in the 700 to 750 lb. weight group.

On a combined or within weight group basis, the largest standard error of estimate for the prediction equations presented was 0.68%. This compares favorably to the standard error of estimate of 1.29% associated

with the prediction equation for percent separable bone of Hankins and Howe (1946).

Multiple Regression Analyses of Retail Yield Measures on Objective Carcass

Measurements. In the combined weight groups, 91% of the variation in

pounds of boned, trimmed round, loin, rib and chuck retail yield was

accounted for using the variables listed in table 26.

Table 26. Coefficient of determination between weight of boned, trimmed round, loin, rib and chuck retail yield and a combination of objective measurements, and the regression coefficient for each of the measurements (combined weight groups).

	Regression	Beta	Level of	R <sup>21</sup>
Independent variables	coefficients	weights	sig.	deletes
Constant	-5.504		0.76	
Carcass weight	0.206	0.860	0.00	0.76
Probe-5th Thoracic-4 in.	303	055	0.33	0.91
Probe-8th Thoracic-8 in.	0.242	0.521	0.34	0.91
Probe-4th Lumbar-12 in.	579	178	0.00	0.89
Probe-Av. 3rd Sacra1 <sup>2</sup>	746	164	0.00	0.90
Length of round	1.925	0.166	0.03	0.90

 $<sup>\</sup>mathbb{R}^2 = 0.91$  Standard error of estimate = 7.06

Using these same variables, 88% of the variation in total carcass boned, trimmed retail yield could be accounted for in the combined weight groups (table 27). On a percent basis in the combined and within weight

<sup>1</sup> Value of coefficient of determination if that variable were deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Av. of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

Table 27. Coefficient of determination between weight of total carcass boned, trimmed retail yield and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (combined weight groups).

of the measure	ments (combined	weight gro	ups).	
	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	-2.973		0.87	
Carcass weight	0.247	0.851	0.00	0.74
Probe-5th Thoracic-4 in.	510	<b>0</b> 76	0.23	0.88
Probe-8th Thoracic-8 in.	0.425	0.076	0.22	0.88
Probe-4th Lumbar-12 in.	739	188	0.00	0.87
Probe-Av. 3rd Sacra12	867	157	0.01	0.87
Length of round	2.262	0.161	0.07	0.88

 $R^2 = 0.88$  Standard error of estimate = 9.66

groups, accuracy was poor in the prediction of percent of either retail yield from the round, loin, rib and chuck or total carcass retail yield.

Also, on a within weight group basis, prediction accuracy of pounds of either of the retail yield measures was poor using only objective carcass measurements as independent variables.

Relationships Between Wholesale Cut Separable Muscle and Carcass Separable

Components, Retail and Fat Trim Yields

In this study, pounds of separable components, retail and fat trim yields were correlated with pounds of separable wholesale cut components, retail, fat trim and bone yields. Percent separable carcass components,

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Av. of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

retail and fat trim yields were correlated with percent wholesale cut separable components, retail, fat trim and bone yields. Percent wholesale cut separable components, retail, fat trim and bone yields were calculated as a percentage of each individual cut rather than as a percentage of the entire carcass.

Simple correlation coefficients between pounds and percent separable wholesale cut muscle and pounds and percent of total carcass separable components, retail and fat trim yields for each weight group as well as for the combined weight groups appear in tables 28, 29 and 30. Correlations between pounds and percent separable wholesale cut muscle and pounds and percent total carcass separable muscle, bone and retail yields were highly significant (P < .01) for the combined weight groups (range, 0.48 to 0.98) except those for pounds and percent of brisket muscle in most instances and percent foreshank muscle (range, 0.02 to 0.61). While within weightgroup correlations were lower than in the combined weight groups, those between pounds of separable rump, foreshank and flank muscle with pounds of total carcass separable muscle, bone and retail yields were markedly reduced.

Correlations between pounds and percents total carcass separable muscle and pounds and percents wholesale round, loin, rib and chuck separable muscle indicate a high relationship (P < .01) exists in the combined and individual weight groups (range, 0.67 to 0.98). Pounds of separable chuck muscle showed the highest relationship to pounds of total carcass separable muscle for combined weight groups (0.98) as well as within

Table 28. Simple ents,	Simple correlation coeffi ents, retail and fat trim	on coefficien fat trim yie	ients between w	en wholesale cut separable the combined weight groups.	separable muht groups.	scle and carcas	cients between wholesale cut separable muscle and carcass separable componingles for the combined weight groups.
Wholesale cut separable			arab	Boned, trimmed	Boned, trimmed	Ω	External fat
	1b. %	1b. %	1b. %	1b. %	İ	% 1b. %	16. %
Rumpless round	0.93 0.84	0.2980	0.90 0.55	0.84 0.61	0.82 0	0.57 0.2957	0.1362
Rump	0.56 0.66	0.1867	0.56 0.48	0.66 0.75	0.68 0	0.760272	J.1072
Intact round	0.97 0.83	0.3180	0.93 0.57	0.92 0.65	0.91 0,	0.62 0.2662	0.0865
Loin	0.94 0.93	0.4290	0.84 0.67	0.87 0.70	0.85 0	0.68 0.4069	0.2469
Ríb	0.88 0.95	0.3693	0.71 0.69	0.84 0.76	0.82 0	0.73 0.2776	0.1173
Chuck	0.98 0.92	0.3788	0.88 0.64	0.91 0.67	0.89 0	0.60 0.3164	0.1662
Foreshank	0.78 0.29	0.4023	0.7305	0.76 0.24	0.75 0	0.23 0.3417	0.1625
Brisket	0.12 0.02	0.1203	0.0603	0.17 0.18	0.61 0	0.47 0.4747	0.0017
Plate	0.90 0.87	0.3787	0.78 0.66	0.86 0.73	0.86 0	0.73 0.2875	0.0971
Flank	0.86 0.91	0.4592	0.75 0.76	0.85 0.79	0.86 0	0.78 0.3180	0.1673
			,				

Correlations > 0.220 are significant (P < .05). Correlations > 0.287 are significant (P < .01). IKLRC = Round, loin, rib and chuck.

Table 29. Simple ents.	Simple correlation coefficie ents, retail and fat trim yi	on coefficien	nts between welds for the	ents between wholesale cut separable musclelds for the 500 to 550 lb, weight group.	eparable muscle weight group.	and carcass	Simple correlation coefficients between wholesale cut separable muscle and carcass separable components, retail and fat trim yields for the 500 to 550 lb. weight group.
Wholesale cut				Boned,	Boned,		
separable	Separable	Separable	Separable	trimmed	trimmed	Total fat	External fat
muscle	muscle 1b. %	fat 1b. %	bone 1b. %	retail RLRC1 lb. %	retail yield 1b. %	trim 1b. %	trim from KLRC 1b. %
Rumpless round	0.80 0.84	8176	0.71 0.42	0.60 0.67	0.57 0.64	6055	5654
Rump	0.58 0.76	60 74	0.54 0.52	08.0 89.0	0.68 0.78	68 74	6575
Intact round	0.88 0.89	9082	0.79 0.50	0.79 0.77	0.77 0.74	99 62	7466
Loin	0.78 0.94	6290	0.42 0.65	0.46 0.74	0.42 0.72	3870	4167
R1b	0.67 0.96	3794	0.09 0.68	0.50 0.82	0.45 0.79	3577	3572
Chuck	0.94 0.93	7288	0.62 0.59	0.68 0.74	0.59 0.66	5764	5561
Foreshank	0.34 0.09	3003	0.4220	0.46 0.17	0.46 0.15	3004	3614
Brisket	02 0.64	0.0859	30 0.37	20 0.46	25 0.44	0.2236	0.2829
Plate	0.65 0.87	5588	0.40 0.69	0.60 0.75	0.60 0.74	5573	64 68
Flank	0.57 0.91	3094	0.30 0.84	0.56 0.80	0.59 0.79	4678	4670
Correlations $> 0.312$ are significant $(P < .05)$	).312 are sis	enificant (P	< .05).				

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01). IRLRC - Round, loin, rib anc chuck.

Table 30. Simpl ents,	e correlation retail and	on coefficien fat trim yie	nts between welds for the	Simple correlation coefficients between wholesale cut separable muscle and carcass ents, retail and fat trim yields for the 700 to 750 lb. weight group.	sparable muscle weight group.		separable compon-
Wholesale cut				Boned,	Boned,		
separable	Separable	Separable	Separable	trimmed	trimmed	Total fat	External fat
muscle	muscle 1b. %	fat 1b. %	bone 1b. %	retail RLRC <sup>1</sup> 1b. %	retail yield 1b. %	trim t 1b. %	trim from RLRC 1b. %
Rumpless round	0.79 0.89	8689	0.74 0.73	0.35 0.62	0.27 0.57	65 68	6577
Runp	0.18 0.49	0.0951	0.18 0.40	0.55 0.58	0.58 0.62	2563	1958
Intact round	0.90 0.83	8483	0.48 0.69	0.63 0.56	0.56 0.54	7965	7670
Loin	0.76 0.92	63 90	0.64 0.71	0.51 0.65	0.47 0.62	5069	4874
Rib	0.68 0.93	5991	0.45 0.70	0.58 0.72	0.56 0.70	68 80	68 80
Chuck	0.92 0.92	7793	0.69 0.75	0.64 0.69	0.58 0.64	7277	69 78
Foreshank	0.35 0.59	5153	0.30 0.20	0.08 0.39	0.08 0.42	4343	97 67
Brisket	0.40 0.88	4085	0.19 0.60	0.07 0.64	0.11 0.62	3573	3267
Plate	0.73 0.88	5585	0.48 0.59	0.55 .0.66	0.56 0.69	6278	6072
Flank	0.62 0.91	3589	0.44 0.61	0.57 0.76	0.60 0.76	4383	4476
, the state of the	201 0 20	4, 64, 54, 70	7 057				

Correlations > 0.321 are significant (P < .05). Correlations > 0.403 are significant (P < .01).

RLRC = Round, loin, rib and chuck.

weight groups (0.94 and 0.92 for the 500 to 550 and 700 to 750 lb. weight groups, respectively). Orme et al. (1960) also found that of all the wholesale cuts, pounds of separable chuck muscle showed the highest relationship to pounds of total separable carcass muscle. Correlations between wholesale round separable muscle and total carcass separable muscle were next highest (0.97, 0.88 and 0.90 for the combined, 500 to 550 lb. and 700 to 750 lb. weight groups, respectively). These findings also coincide with those of Orme et al. (1960). The percents separable muscle of these same wholesale cuts were as high, or more highly related to percent total carcass separable muscle within the two weight groups as in the combined weight groups (range, 0.83 to 0.93).

The highest correlation in the combined weight groups between percent wholesale cut separable muscle and percent carcass muscle was the rib (0.95). In order of decreasing magnitude of correlation coefficients between the above two traits were the loin (0.93), chuck (0.92) and flank (0.91). All correlations between percent wholesale cut separable muscle and percent carcass separable muscle were highly significant (P < .01) for the combined weight groups except for that of the brisket.

Within weight group correlation coefficients between percent wholesale cut separable muscle and percent carcass separable muscle showed

percent rib, loin, chuck and flank muscle being most closely related to

percent total carcass muscle (range, 0.91 to 0.96). All of these correlations have excellent predictive possibilities.

High negative correlations were found between the percent separable muscle of the rib and flank with percent separable fat and fat trim yields

in the combined and within weight groups. Within weight groups, separable muscle of the rib and flank also showed the highest relationships to percent separable retail yield from the round, loin, rib and chuck and percent total retail yield (range, 0.79 to 0.82). Hedrick et al. (1963) and Miller et al. (1965) also reported very high correlation coefficients between percent retail yield of the flank and retail yield measures. In their studies, however, neither the flank nor the rib showed the highest correlation to retail yield but instead percent retail yield of the round was highest.

The high relationships between percent separable flank muscle and percents of the seven carcass yields compared in this study, suggest the use of flank muscle for predicting these carcass yields. These relationships were high for both weights and percents in the combined weight groups as well as within weight groups (range, 0.16 to 0.94).

Correlation coefficients between pounds of wholesale cut separable muscle and carcass separable fat and fat trim yields were positive but lower than for pounds of separable muscle and retail yields (range, 0.00 to 0.47) on a combined weight group basis, except for correlations between pounds of rump muscle and fat trim yields (-.02 and -.10). However, correlations between percent wholesale cut separable muscle in the combined weight groups as well as weights and percents within weight groups became negative but were highly significant (P < .01) in most instances.

Relationships Between Wholesale Cut Separable Fat and Carcass Separable

Components, Retail and Fat Trim Yields

Simple correlation coefficients between pounds and percent wholesale cut separable fat and pounds and percent of carcass separable components, retail and fat trim yields for the combined weight groups appear in table 31 and those within weight groups appear in tables 32 and 33.

In the combined weight groups, pounds of wholesale cut separable fat showed the lowest relationships to pounds of separable carcass bone (range, 0.18 to 0.36) and low to moderate relationships to pounds of carcass separable muscle and retail yields (range, 0.22 to 0.55). Highly significant (P < .01) positive correlations were observed between pounds and percent wholesale cut separable fat and carcass separable fat and fat trim yields (range, 0.61 to 0.96), except for kidney knob (range, 0.32 to 0.62). When percent wholesale cut separable fat was correlated with percents carcass separable muscle, bone and retail yields on a combined weight group basis, these correlations (range, 0.54 to 0.94) were highly significantly (P < .01) negative.

On a combined weight group basis, the percent wholesale cut separable fat most closely related to percent carcass separable fat was that from the rib (0.96), followed in order by the loin (0.94), chuck (0.94), plate (0.92) and the round and flank (0.91). Within weight group correlations were also highest for these same wholesale cut percents separable fat and carcass separable fat though not in the same order of magnitude (tables 32 and 33). Percents separable fat from the kidney knob and foreshank showed the lowest relationship to carcass separable fat on a within weight

Wholesale cut separable fat				•	•		
separable fat				Boned,	Boned,		
	Separable muscle	Separable fat	Separable bone	$ ext{trimmed}$ $ ext{retail}$ $ ext{RLRC}^1$	re	Total fat trim	External fat trim from KLRC
	1b. %	1b. %	1b. %	1b. %	` [	1b. %	1b. %
Rumpless round	0.3786	0.87 0.87	0.3056	0.4768	0.4563	0.79 0.67	0.79 0.72
Rump	0.3382	0.87 0.85	0.2458	0.4680	0.4680	0.74 0.82	0.73 0.82
Intact round	0.3790	0.90 0.91	0.2962	0.4875	0.4871	0.80 0.75	0.79 0.78
Loin	0.2993	0.93 0.94	0.1871	0.4073	0.3869	0.87 0.73	0.84 0.73
Rib	0.2294	0.92 0.96	0.0879	0.3280	0.3177	0.86 0.81	0.82 0.78
Chuck	0.4592	0.94 0.94	0.3167	0.5374	0.5269	0.84 0.75	0.77 0.73
Foreshank	0.2460	0.65 0.61	0.1947	0.2954	0.3048	0.61 0.53	0.61 0.60
Brisket	0.3079	0.84 0.80	0.2046	0.3365	0.3264	0.80 0.66	0.74 0.62
Plate	06'- 95'0	0.93 0.92	0.3678	0.5576	0.5376	0.85 0.78	0.72 0.74
Flank	06 98.0	0.94 0.91	0.2630	0.4678	0.4578	0.86 0.79	0.76 0.73
Kidney knob	0.3157	0.72 0.58	0.2547	0.3554	0.3453	0.62 0.42	0.50 0.32

Correlations > 0.220 are significant (P < .05). Correlations > 0.287 are significant (P < .01).

RLRC = Round, loin, rib and chuck.

Table 32. Simple correlation coefficients between wholesale cut separable retail and fat trim yields for the 500 to 550 lb. weight group.	le correlation il and fat to	Simple correlation coefficients bet retail and fat trim yields for the	nts between w or the 500 to	ween wholesale cut se 500 to 550 lb. weight	cut separable fat and weight group.	carcass sep	fat and carcass separable components,
Wholesale cut				Boned,	Boned,		
separable	Separable	Separable	Separable	trimmed	trimmed	Total fat	External fat
fat	muscle 1b. %	fat 1b. %	bone 1b. %	retail RLRC <sup>1</sup> 1b. %	retail yield lb. %	trim 1b. %	trim from RLRC 1b. %
Rumpless round	8188	0.67 0.87	5266	6477	6376	0.58 0.71	0.60 0.69
Rump	7186	0.65 0.88	5273	5187	98 67	0.50 0.84	0.54 0.84
Intact round	8591	0.73 0.91	5871	-,6583	6382	0.61 0.78	0.64 0.76
Lofn	8595	0.93 0.95	7576	6679	6676	0.81 0.75	0.76 0.71
Rib	7695	0.95 0.97	8181	6284	6482	0.83 0.81	0.77 0.76
Chuck	6894	0.90 0.96	6778	5281	5176	0.72 0.76	0.68 0.73
Foreshank	4566	0.30 0.65	1551	3471	2663	0.28 0.60	0.34 0.67
Brisket	5871	0.77 0.73	5858	5361	5762	0.70 0.58	0.66 0.54
Plate	69 89	0.92 0.92	69 78	4978	5078	0.74 0.77	0.61 0.71
Flank	7090	0.90 0.92	7880	5078	4978	0.71 0.77	0.61 0.69
Kidney knob	5471	0.77 0.74	5764	4263	4261	0.59 0.55	0.46 0.44

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01). 1 RLRC = Round, loin, rib and chuck.

Table 33. Simple retail	le correlati	on coefficien rim yields fo	nts between w or the 700 to	Simple correlation coefficients between wholesale cut separable retail and fat trim yields for the 700 to 750 lb, weight group.		fat and carcass sep	separable components,
Wholesale cut				Boned,	Boned,		
separable	Separable	Separable	Separable	trimmed	trimmed	Total fat	External fat
Tar	1b. %	16. %	16. %	1b. %	1b. %	1b. %	1b. %
Rumpless round	7388	0.80 0.90	5982	3662	2954	0.72 0.70	0.80 0.80
Rump	7275	0.84 0.80	6377	2766	2365	0.63 0.77	0.67 0.76
Intact round	7789	0.86 0.93	64 85	3566	2959	0.74 0.75	0.80 0.83
Loin	6992	0.84 0.94	5583	3266	2763	0.73 0.76	0.78 0.81
Rib	7792	0.85 0.94	65 79	3672	3069	0.73 0.83	0.71 0.83
Chuck	6388	0.81 0.92	62 85	3366	2660	0.69 0.77	0.68 0.77
Foreshank	6265	0.58 0.69	5366	5156	4653	0.57 0.66	0.59 0.70
Brisket	69 89	0.70 0.89	5469	59 69	5269	0.67 0.82	0.60 0.76
Plate	6691	0.78 0.90	4368	3568	3369	0.69 0.81	0.58 0.76
Flank	7590	0.88 0.89	4662	4175	3675	0.83 0.83	0.70 0.77
Kidney knob	2536	0.37 0.32	0718	1942	1540	0.21 0.16	0.16 0.13

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01). 1 RLRC = Round, loin, rib and chuck.

group basis (0.58 and 0.61, respectively). As was shown for percent carcass separable fat, percent kidney knob separable fat had the lowest relationships with the fat trim yields in the combined weight groups and within weight groups. These data indicate kidney fat deposition is not entirely consistent with total carcass fat deposition.

Comparing the correlation coefficients within weight groups, the relationships between percents wholesale cut separable fat and percents of the seven carcass yield characteristics were higher in the 500 to 550 lb. weight group (range, 0.44 to 0.94) than in the 700 to 750 lb. weight group (range, -.13 to 0.94). However, the differences were not great and most correlations were highly significant (P < .01) and sufficiently high to be of predictive value.

Relationships Between Wholesale Cut Separable Bone and Carcass Separable

Components, Retail and Fat Trim Yields

Correlation coefficients between pounds and percent wholesale cut separable bone and pounds and percent carcass separable components, retail and fat trim yields for the combined weight groups and for each weight group appear in tables 34, 35 and 36. Highly significant (P < .01) positive correlations were observed for the combined weight groups between pounds and percent of carcass separable muscle, bone and retail yields (range, 0.36 to 0.98) except for pounds and percent flank bone and percent rump bone (0.12 to 0.27). Also highly significant (P < .01) negative relationships were found between percent wholesale cut separable bone and percent carcass separable fat and fat trim yields for the combined weight groups (range -.41 to -.80) except for percent flank and rump bone (range,

Simple correlation coefficients between wholesale cut separable bone and carcass separable components, RL RC External fat -.16 -.53 -.34 -.48 -.58 -.49 -. 65 -.53 -.54 -.41 trim from 0.10 -.01 90.0 0.13 0.04 0.04 0.04 -.13 0.08 0.02 <u>1</u>p Total fat 0.24 -.52 0.12 -.33 -.49 0.18 -.56 -.41 -.55 -.59 -.17 0.28 -.51 0.18 -.71 trim 0.21 0.00 0.25 0.21 0.04 1b. retail yield 0.45 0.36 0.46 0.16 0.21 0.42 0.50 0.63 0.32 0.50 trimmed 0.79 0.80 0.82 0.80 0.80 0.45 0.08 0.82 0.82 0.77 1b. retail and fat trim yields for the combined weight groups. retail RLRC 0.45 0.23 0.39 0.49 0.61 0.48 0.36 0.46 0.54 0.12 trimmed Boned, 0.81 0.81 0.84 0.82 0.82 0.82 0.43 0.79 0.83 0.07 1b. Separable 0.95 0.78 0.93 0.74 0.98 0.83 0.94 0.85 0.89 0.91 0.97 0.92 0.92 0.60 0.53 0.55 0.93 0.88 0.15 0.32 bone 1b. Separable -.80 -.74 -.79 -.80 0.29 -.71 0.24 -.74 0.28 -.76 0.28 -.54 0.10 -.62 -.01 -.73 fat 0.29 0.24 0.31 0.31 1b. Separable 0.85 0.46 0.70 0.81 0.52 0.87 0.53 0.87 0.55 0.45 0.38 0.86 0.63 0.88 0.72 0.80 0.41 0.12 0.27 muscle 0.84 1b. Rumpless round cut Intact round separable Wholesale Table 34. Foreshank Brisket Chuck Plate Flank Loin Rump Rib

Correlations > 0.220 are significant (P < .05). Correlations > 0.287 are significant (P < .01). IRLRC = Round, loin, rib and chuck.

iable 55. Simple retail	il and	and fat trim yields for the 500	in coer	lds fo	r the	500 to	to 550 lb.	e cut ser . weight	group.	חסוום שוי	ות כשורני	മോട ഉപ്പ	arante d	retail and fat trim yields for the 500 to 550 lb, weight group.
Wholesale cut							Boned		Boned,	đ,				
separable	Sepai	Separable	Separable	able	Separable	able	trimmed	ed m no1	trimmed	ed	Total fat	fat	External	al fat
bone	muscle 1b.	% \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	1b.	ار ا	Done 1b.	w   %	retail 1b.	KLRC.	1b.	retall yleld 1b. %	1b.	#I	trim from 1b. %	N KLKC
Rumpless round	0.59	0.59 0.4972	72	59	0.92	0.79	0.58	0.56	0.57	0.56	65	60	60	57
Rump	0.47	0.47 0.39 4.70	4.70	49	0.87	0.74	0.54	0.28	0.57	0.30	71	37	61	32
Intact round	0.58	0.58 0.5276	76	63	0.95	0.87	0.60	0.53	09.0	0.55	71	60	64	55
Loin	0.55	0.67	63	76	08.0	0.87	0.45	0.61	0.38	0.56	42	56	-, 39	50
Rib	0.62	0.62 0.72	60	82	0.74	0.93	0.54	0.65	0.57	0.68	60	72	-,55	64
Chuck	0.67	0.67 0.6078	78	72	0.94	0.93	0.57	0.57	0.55	0.57	99	62	62	57
Foreshank	0.48	0.48 0.4868	68	54	06.0	0.67	0.50	0.45	0.52	0.44	64	51	-, 60	45
Brisket	0.04	0.04 0.28	23	-, 38	0.32	0.52	0.16	0.38	0.22	0.44	33	48	37	52
Plate	0.48	0.48 0.68	55	76	0.78	06.0	0.40	0.56	0.39	0.54	49	58	51	51
Flank	0.03	0.03 0.21	08	25	0.12	0.32	10	0.01	05	0.05	0.05	07	0.04	90

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01). 1 KLRC = Round, loin, rib and chuck.

retail	11 and	and fat trim yields for	im yie	lds fo	r the 700	700 to	to 750 lb.	. weight	BI	-				
separable	Separ	Separable	Separable	able	Separable		trimmed	ed prod	trimmed	a, ed 13	Total fat	fat	External fat	al fat
ропе	muscie 1b.	%	1b.	<b>"</b>	1b.	%	1b. %	%	1b.	lb. %	1b.		1b. %	%
Rumpless round	0.57	0.57 0.4263	63	55	0.85	08.0	0.32	0.34	0.22	0.23	49	46	52	54
Rump	0.67	0.67 0.7552	52	81	0.86	0.86	0.62	0,40	0.54	0.32	60	53	62	60
Intact round	0.66	0.66 0.5965	65	69	0.93	0.86	0.46	0.37	0.35	0.28	57	51	60	59
Loin	0.70	0.64	55	74	06.0	06.0	0.57	0.51	0.51	0.45	52	-, 66	54	69
Rib	0.64	0.64 0.66	44	74	0.78	0.87	0.64	0.47	0.62	0.44	52	-, 65	54	62
Chuck	0.72	0.72 0.46	57	58	0.94	0.89	0.55	0.30	0.46	0.22	51	46	52	48
Foreshank	0.54	0,40	48	48	0.76	0.62	0.44	0.38	0.37	0.32	44	45	56	49
Brisket	0.53	0.66	-, 33	70	0.51	0.71	0.35	09.0	0.37	09.0	40	70	42	68
Plate	0.82	0.80	66	85	0.86	0.87	0.62	09.0	0.57	0.56	64	72	64	. 68
Flank	0.35	0.35 0.4515		44	0.26	0,33	0.26	0.30	0.30	0.35	16	40	16	35

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01). IRLRC = Round, loin, rib and chuck.

-.16 to -.34) with carcass fat yields. Within weight groups, high negative correlations were found between pounds and percents of wholesale cut separable bone and pounds and percents of carcass separable fat and fat trim yields (range, -.39 to -.78) except pounds and percents brisket and flank bone and percent rump bone (range, -.08 to -.49).

These correlations indicate that carcasses with heavier bone and/or higher percent separable bone also had more pounds and/or percent total carcass separable muscle and retail yields. Wythe et al. (1961) and Good et al. (1961) also reported this positive relationship between size and/or quantity of bone and amount of carcass muscle.

Highest relationships were found between pounds of round separable bone and pounds of carcass separable bone for the combined weight groups as well as within each weight group (0.98, 0.95 and 0.93, respectively). The next highest correlations were between pounds of separable chuck bone and pounds of carcass separable bone both within weight groups and on a combined weight group basis.

Percent separable bone in the chuck showed the highest correlation with percent carcass separable bone in the combined weight group and in the 500 to 550 lb. weight group (0.92 and 0.93, respectively). Following the chuck were percent separable bone in the rib (0.91 and 0.93) and plate (0.88 and 0.90) for the combined and 500 to 550 lb. weight groups, respectively. In the 700 to 750 lb. weight group the percent separable bone of the loin showed the highest correlation to percent carcass separable bone (0.90) followed by the chuck (0.89), rib (0.87) and the plate (0.87). Although correlations were not consistently as high as percent rib or chuck bone, percent wholesale round bone was highly significantly

(P < .01) correlated with percent carcass separable bone in the combined weight groups and the 500 to 550 and 700 to 750 lb. weight groups (0.83, 0.87 and 0.86, respectively). Due to the ease of separation of total round bone compared to the other wholesale cuts which were highly correlated to total carcass bone, it is logical to suggest its use for predictive purposes.

Relationships Between Percent Separable Components of Some Easily Separated
Wholesale Cuts and Percent Separable Carcass Components

The 9-10-11 rib section, the flank, intact round and the rump were selected to study their possible predictive value of carcass composition, due to their ease of physical separation. Simple correlation coefficients were calculated between the percent separable components of these cuts and the percents of these same components of the entire carcass for the combined and individual weight groups as well as on a within weight and fat thickness group basis. These correlations appear in table 37.

The 9-10-11 rib section was choosen not only because of ease of separation, but also to determine the reliability of the regression equations reported by Hankins and Howe (1946) between percent separable components and separable carcass components within a group of carcasses of limited weight and fat thickness ranges. The relationship between percent muscle of the 9-10-11 rib section and percent separable carcass muscle was very high in the combined weight groups and within weight groups (0.92, 0.94 and 0.90, respectively). Crown and Damon (1960) reported correlation coefficients of similar magnitude between percent separable muscle of the 9-10-11 rib section and percent carcass separable

Simple correlation coefficients between percent separable components of some wholesale cuts ments of the cerces Table 37.

and	and separable com	ponents or	the carcass.								
Cut and %	Combined										
separable	weight	500-550 1b.	700-750 1b.				Group				
component	group	weight group	weight group	н	ij	III	IV	Δ	VI	VII	VIII
•											
Muscle:											
9-10-11 rfb											
section	0.92	0.94	0.00			0.94	0.92	0.83			
Flank	0.91	0.91	0.91			0.78	0.76	0.95			
Round	0.83	0.89	0.83	0.87	0.52	0.80	0.91	0.30	0.89	0.82	98.0
Rump	99.0	0.76	0.49			0.62	0.51	0.31			
Fat:											
9-10-11 rib											
section	0.94	0.95	0.91			0.95	0.90	•			0.75
Flank	0.91	0.92	0.89	0.93	0.89	0.77	0.71	0.92	0.87	0.80	0.73
Round	0.91	0.91	0.93			0.83	0.86		•		0.87
Rump	0.85	0.88	08.0			0.74	0.68				0.79
Bone:											
9-10-11 rib											
section	0.76	0.78	0.75	0.87	0.67	0.45	0.64	0.09	0.79	0.57	0.74
Flank	0.32	0.32	0.33	0.74	0.68	06		_			0.45
Round	0.83	0.87	98.0	0.74	0.82	0.76		_			0.79
Rump	0.74	0.74	98.0	0.76	0.80	0.20		_		-	0.74

Correlations > 0.220 are significant (P < .05) for the combined weight groups. Correlations > 0.287 are significant (P < .01) for the combined weight groups. Correlations > 0.312 are significant (P < .05) for the individual weight groups. Correlations > 0.403 are significant (P < .01) for the individual weight groups.

muscle. The correlations on a within weight and fat group basis were slightly lower, especially in group VI (range, 0.71 to 0.94).

Percent separable flank muscle showed an equally good relationship to percent separable carcass muscle (range, 0.76 to 0.95). Percent separable muscle of the wholesale round and rump, especially the rump, did not show quite as high or as consistent a relationship to percent carcass muscle (range, 0.30 to 0.91 and 0.11 to 0.76 for round and rump, respectively) as either the flank or 9-10-11 rib sections.

Similar results were observed between the percent separable fat of these cuts and percent separable carcass fat. The percent separable fat of the 9-10-11 rib section, flank, and wholesale round were highly correlated with percent carcass separable fat (range, 0.71 to 0.95). When comparing these relationships within weight and fat groups, the percent separable fat of the 9-10-11 rib sections and the flank were more consistently correlated to separable carcass fat (range, 0.71 to 0.95) than either the wholesale round or rump (range, 0.30 to 0.93).

When comparing correlations between percent separable bone of these four cuts with percent separable carcass bone, the percent round bone showed the highest and most consistent relationships, followed by those of the rump. Correlation coefficients for these characteristics within the eight fat thickness and weight groups ranged from 0.55 to 0.94 and 0.20 to 0.95 for the round and rump, respectively. The percent bone in the flank showed the poorest relationship with percent carcass bone which might be expected due to the small amount of bone present in this wholesale cut (range, -.06 to 0.74). The correlations between percent

9-10-11 rib bone and carcass separable bone were also comparitively low and not consistent within the eight fat thickness and weight groups (range, 0.09 to 0.87).

These data support the accuracy of Hankins and Howe (1946) equations and these observations and those of Crown and Damon (1960) indicate that the percent separable components of the 9-10-11 rib sections were good predictors of carcass components, particularly muscle and fat. However, these data also indicate that the percent separable muscle and fat from the flank were essentially equally as good as the 9-10-11 rib section for estimating percent carcass muscle and fat. Hankins and Howe: (1946) reported that the percent ether extract of the flank showed the highest correlation (0.95) to percent ether extract of the edible portion of the dressed beef carcass. Because of the ease of physical separation of the flank as well as for economic reasons, this cut could be used with essentially the same degree of accuracy as the 9-10-11 rib section separation which today is the most widely used method for estimating total carcass muscle and fat.

Multiple Regression Analyses of Separable Carcass Components, Retail and Fat Trim Yields on Objective Carcass Measurements and Wholesale Cut

## Separable Components

Multiple Regression Analyses of Total Carcass Separable Muscle on Objective Carcass Measurements and Weight of Wholesale Cut Separable Components. In the combined weight groups, 98% of the variation in pounds of separable carcass muscle could be accounted for using the six independent variables presented in table 38. Of these six independent variables, carcass weight,

weight of separable flank fat and weight of separable flank muscle appeared to have the greatest effect on separable carcass muscle. These data support the findings of Cole et al. (1960) who reported a simple correlation between carcass weight and pounds of separable carcass muscle of 0.77. These authors also reported multiple correlations (ranging from 0.85 to 0.97) using carcass weight and weight of wholesale cut separable muscle. The prediction equation shown in table 38 had a standard error of estimate of 3.91 lb. which was less than the most accurate prediction equation of Cole et al. (1960). The most accurate equation of these authors involved separable round muscle as the independent variable for predicting pounds of carcass separable muscle. This equation had a standard error of estimate of 4.38 lb.

Table 38. Coefficient of determination between weight of carcass separable muscle and a combination of weights of wholesale cut separable components and objective carcass measurements, and the regression coefficient for each of the variables (combined weight group).

weight group)				1
	Regression	Bet <b>a</b>	Level of	RZT
Independent variables	coefficients	weights	sig.	deletes
Constant	1.634		0.68	
Carcass wt.	0.282	1.052	0.00	0.79
Probe-Av. 3rd Sacral2	718	140	0.00	0.97
Wt. sep. flank muscle3	3.302	0.215	0.00	0.96
Wt. sep. flank fat	-2.412	401	0.00	0.94
Wt. kidney knob-L.S.4	643	093	0.00	0.97
L.dorsi muscle area	0.779	0.052	0.03	0.98

 $R^2 = 0.98$  Standard error of estimate = 3.91

Value of coefficient of determination if that variable was deleted from the analysis.

 $<sup>^2</sup>$ Av. of 3 probes 4, 8 and 12 in. off the carcass dorsal midline.

 $<sup>^{3}</sup>$ Sep. = separable.

<sup>4</sup>L.S. = left side.

The six independent variables listed in table 39 were associated with 90% of the variation in percent separable carcass muscle in the combined weight groups. The prediction equation given in table 39 showed a standard error of estimate of 1.33%. This compares favorably with the standard error of estimate of 2.51% for the prediction equation of Hankins and Howe (1946) who used percent separable muscle from the 9-10-11 rib section to predict percent carcass muscle.

Table 39. Coefficient of determination between percent carcass separable muscle and a combination of weights of wholesale cut separable components and objective carcass measurements, and the regression coefficient for each of the variables (combined weight groups).

Independent variable	Regression coefficients	Beta weights	Level of sig.	R21 deletes
Constant	54.012		0.00	
Carcass wt.	0.014	0.315	0.00	0.89
Probe-5th Thoracic-4 in.	143	141	0.01	0.89
Probe-Av. 3rd Sacral <sup>2</sup>	229	274	0.00	0.87
Wt. sep. flank muscle <sup>3</sup>	0.883	0.352	0.00	0.87
Wt. sep. flank fat	690	703	0.00	0.80
Kidney knob wtL.S.4	266	234	0.00	0.87

 $R^2 = 0.90$  Standard error of estimate = 1.33

In the 500 to 550 lb. weight group, 89% of the variation in pounds of separable carcass muscle could be accounted for by the following five variables: 1) carcass weight, 2) average of three fat probes at the third sacral vertebra, 3) weight of separable flank muscle, 4) weight of separable flank fat and 5) weight of the kidney knob from the left side of the carcass. The multiple regression analysis of separable carcass muscle weight on these five variables is given in the supplement, p. 177.

<sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Av. of 3 probes 4, 8 and 12 in. off the carcass dorsal midline.

<sup>&</sup>lt;sup>3</sup>Sep. = separable.

<sup>4</sup>L.S. = left side.

Using these same five variables, 92% of the variation in percent separable carcass muscle in the 500 to 550 lb. weight group can be accounted for (table 40). The effect of carcass weight on percent total carcass separable muscle was relatively unimportant in these regression analyses.

Table 40. Coefficient of determination between percent carcass separable muscle and a combination of weights of wholesale cut separable components and objective carcass measurements, and the regression coefficient for each of the variables (500 to 550 lb. weight group).

group).				
Independent variable	Regression coefficients	Beta weights	Level of sig.	R21 deletes
Constant	58.876		0.00	
Carcass wt.	0.007	0.025	0.72	0.92
Probe-Av. 3rd sacra12	289	324	0.00	0.86
Wt. sep. flank muscle <sup>3</sup>	0.942	0.201	0.00	0.89
Wt. sep. flank fat	806	516	0.00	0.82
Kidney knob wtL.S.4	<b></b> 351	228	0.00	0.89

 $R^2 = 0.92$  Standard error of estimate = 1.39

In the 700 to 750 lb. weight group, the independent variables given in table 41 were responsible for 93% of the variation in pounds of separable carcass muscle. The prediction equation presented in this table showed a high degree of accuracy with a standard error of estimate of 3.74 lb. These same six variables were responsible for 91% of the variation in percent separable carcass muscle in the 700 to 750 lb. weight group (suppl., p.177.). This equation showed a standard error of estimate of 1.18%.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

 $<sup>^{2}</sup>$ Av. of 3 probes 4, 8 and 12 in. off the carcass dorsal midline.

 $<sup>^{3}</sup>$ Sep. = separable.

<sup>4</sup>L.S. = left side.

Table 41. Coefficient of determination between weight of carcass separable muscle and a combination of weights of wholesale cut separable components and objective carcass measurements, and the regression coefficient for each of the variables (700 to 750 lb. weight

Regression	Beta	Level of	R21
coefficients	weights	sig.	deletes
80.178		0.00	
0.177	0.267	0.00	0.88
444	126	0.04	0.92
<b></b> 775	279	0.00	0.90
3.918	0.336	0.00	0.85
-2.079	507	0.00	0.83
635	151	0.00	0.91
	80.178 0.177 444 775 3.918 -2.079	80.178 0.177 0.267444126775 3.918 0.336 -2.079507	80.178         0.00           0.177         0.267         0.00          444        126         0.04          775        279         0.00           3.918         0.336         0.00           -2.079        507         0.00

 $R^2 = 0.93$  Standard error of estimate = 3.74

Ninety-six percent of the variation in weight of total carcass separable muscle in the combined weight groups could be accounted for using the three variables presented in table 42. In the 500 to 550 lb. weight group, only 79% of the weight of total carcass separable muscle could be accounted for using these same variables (suppl., p. 177) and 86% in the 700 to 750 lb. weight group (suppl., p 178). The standard error of estimates associated with these equations were 5.19, 5.29 and 4.96 lb., respectively. It is of interest to note that, in contrast with other equations reported in this dissertation, the equation predicting weight of total carcass separable muscle in the 700 to 750 lb. weight group was more accurate than that for the prediction of this same trait in the 500 to 550 lb. weight group.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>2</sup>Av. of 3 probes 4, 8 and 12 in. off the carcass dorsal midline.

 $<sup>^{3}</sup>$ Sep. = separable.

<sup>&</sup>lt;sup>4</sup>L.S. = left side.

Table 42. Coefficient of determination between weight of carcass separable muscle and a combination of weights of wholesale cut separable components, and the regression coefficient for each variable (combined weight group)

(Companed wea	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	-3.193		0.44	
Carcass wt.	0.284	1.060	0.00	0.75
Wt. sep. flank muscle <sup>2</sup>	4.409	0.287	0.00	0.93
Wt. sep. flank fat	-3.394	564	0.00	0.83

 $R^2 = 0.96$  Standard error of estimate = 5.19

Using carcass weight, weight of separable flank muscle and flank fat, 81% of the variation in percent of total separable carcass muscle could be accounted for with a standard error of estimate of 1.81% in the combined weight group (table 43). These same variables accounted for 83% of the variation in this trait in the 500 to 550 lb. weight group (suppl., p. 178) and 86% of the variation in the 700 to 750 lb. weight group (suppl., p. 178). These equations had standard errors of estimate of 1.97 and 1.42%, respectively. Thus, greater accuracy was also

attained in the 700 to 750 lb. carcasses than in the 500 to 550 lb.

carcasses.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>2</sup>Sep. = separable.

Table 43. Coefficient of determination between percent carcass separable muscle and a combination of weights of wholesale cut separable components, and the regression coefficient for each of the variables (combined weight groums)

	Regression	Beta	Level of	R21
Independent variable	coefficients	weights	sig.	deletes
Constant	50.345		0.00	
Carcass wt.	0.012	0.267	0.02	0.80
Wt. sep. flank muscle <sup>2</sup>	1.266	0.504	0.00	0.73
Wt. sep. flank fat	-1.100	-1.121	0.00	0.31

 $R^2 = 0.81$ Standard error of estimate = 1.81

Multiple Regression Analyses of Total Carcass Separable Muscle on Objective Carcass Measurements and Percent Wholesale Cut Separable Muscle. The 9-10-11 rib section is used rather extensively for predicting carcass composition because of the high relationships to total carcass composition and the ease of separation of this section. Prediction equations for estimating carcass composition from the wholesale flank and round were calculated and were compared to the 9-10-11 rib section.

In the combined weight groups, using carcass weight, length of round and the percent separable muscle of the respective cuts, pounds and percent total carcass separable muscle could be most accurately predicted using the 9-10-11 rib section separable muscle, followed by flank separable muscle and then round separable muscle (table 44). Differences were small between equations 1, 2 and 3 for predicting pounds of total carcass separable muscle. It should be noted that in using equations 4, 5 and 6 to predict percent carcass separable muscle using 9-10-11 rib

 $<sup>^{</sup>m l}$ Value of coefficient of determination if that variable was deleted from 2the analysis. Sep. = separable.

section or flank separable muscle gave similar results, but using round separable muscle in the equation decreased accuracy.

Table 44. Multiple regression equations for estimating pounds and percent total carcass separable muscle using objective carcass measurements and percent wholesale cut separable muscle as predictors, and the coefficient of determination for each equation (combined weight groups).

Equation No.	Estimating equation	Coeffic determi		Standard error of estimate
	$-103.986 + 0.239(X_1) + 0.470(X_2) + 2.1$		0.96	5.28
2 🏠 =	$-78.232 + 0.252(X_1) + 0.438(X_2) + 1.88$	30(X4)	0.95	5.77
3 <b>Ŷ</b> =	$-200.864 + 0.222(X_1) + 1.978(X_2) + 2.66$	60(X <sub>5</sub> )	0.94	6.32
4 <b>1</b> =	$16.767006(x_1) + 0.212(x_2) + 0.7056$	(X3)	0.86	1.57
5 <b>1</b> =	$25.052002(X_1) + 0.198(X_2) + 0.6060$	(X <sub>4</sub> )	0.83	1.74
6 Ŷ <sub>1</sub> =	$-12.531013(X_1) + 0.744(X_2) + 0.814$	4(X <sub>5</sub> )	0.74	2.14
_				

 $<sup>\</sup>hat{Y}_1$  = Estimated percent total carcass separable muscle.

 $X_1 = Carcass weight (1b.).$ 

 $X_2$  = Length of round (in.)

 $X_3$  = Percent 9-10-11 rib section separable muscle.

 $X_4$  = Percent flank separable muscle.

 $X_5$  = Percent round separable muscle.

In contrast to this, round separable muscle used in the equations shown in table 45 was the most accurate predictor of these three cuts in the 500 to 550 lb. weight group, while flank separable muscle was least accurate. However, differences among these equations were not great. In the 700 to 750 lb. weight group (table 46), these equations showed

Table 45. Multiple regression equations for estimating pounds and percent total carcass separable muscle using objective carcass measurements and percent wholesale cut separable muscle as predictors, and the coefficient of determination for each equation (500 to 550 lb. weight group).

	equation (500 to 550 lb. weight a	group).		
E-mati		0 454		Standard
Equation No.	Estimating equation		cient of nination	error of estimate
				ESCIMACE
1 🕎 =	$-42.462 + 0.180(X_1)375(X_2) + 2$	2.033(X <sub>3</sub> )	0.85	4.50
2 🕎 =	$16.929 + 0.167(X_1) - 1.138(X_2) + 1$	. 691(X <sub>4</sub> )	0.80	5.17
3 <b>Ŷ</b> =	$-125.147 + 0.063(X_1) + 1.303(X_2) +$	- 3.094(X <sub>5</sub> )	0.86	4.25
4 $\hat{Y}_1 =$	$35.418028(X_1)096(X_2) + 0.7$	'36(X <sub>3</sub> )	0.89	1.59
5 <b>?</b> 1=	$57.036034(x_1)356(x_2) + 0.6$	506(X <sub>4</sub> )	0.84	1.90
6 Ŷ <sub>1</sub> =	$5.896071(X_1) + 0.516(X_2) + 1.1$	.13(X <sub>5</sub> )	0.89	1.55
$\mathbf{\hat{Y}} = \mathbf{Est}$	timated pounds of total carcass sep	arable mus	cle.	
$\hat{Y}_1 = E_1$	stimated percent total carcass sepa	rable musc	le.	
$X_1 = C_1$	arcass weight (lb.)			
$x_2 = Le$	ength of round (in.).			
$x_3 = Pe$	ercent 9-10-11 rib section separabl	e muscle.		
$X_4 = Pe$	ercent flank separable muscle.			
$x_5 = Pe$	ercent round separable muscle.			

flank separable muscle as the most accurate predictor of weight or percent total carcass separable muscle. In this weight group, round separable muscle showed relatively low prediction accuracy for either pounds or percent total carcass separable lean.

Table 46. Multiple regression equations for estimating pounds and percent total carcass separable muscle using objective carcass measurements and percent wholesale cut separable muscle as predictors, and the coefficient of determination for each equation (700 to 750 lb. weight group).

	equation (700 to 750 lb. weight group	цр).		
Equation		Coeffic	ient of	Standard error of
No.	Estimating equation	determi		estimate
	$87.434 + 0.178(x_1) + 1.183(x_2) + 2.3$	330(X <sub>3</sub> )	0.81	5.79
$2 \hat{Y} = -$	$33.168 + 0.116(x_1) + 1.763(x_2) + 2.2$	236(X <b>4)</b>	0.87	4.77
3 Ŷ = -	$153.699 + 0.210(X_1) + 1.170(X_2) + 2.$	434(X <sub>5</sub> )	0.70	7.34
4 $\hat{Y}_1 = 1$	$9.892014(x_1) + 0.388(x_2) + 0.646$	5(X3)	0.82	1.59
$5  \stackrel{\wedge}{\mathbf{Y}}_{1} = 3$	$5.084032(x_1) + 0.562(x_2) + 0.613$	3(x <sub>4</sub> )	0.87	1.38
6 $\Re_{1}^{2} = 2$	.305006( $x_1$ ) + 0.407( $x_2$ ) + 0.663	(X <sub>5</sub> )	0.70	2.08
🕏 = Esti	mated pounds of total carcass separa	able musc	1e.	
$\hat{Y}_1 = Est$	imated percent total carcass separat	le muscl	e.	
$X_1 = Car$	cass weight (1b.).			
$X_2 = Len$	gth of round (in.).			
$X_3 = Pere$	cent 9-10-11 rib section separable m	nuscle.		
$X_4 = Pere$	cent flank separable muscle.			
X <sub>5</sub> = Per	cent round separable muscle.			

Multiple Regression Analyses of Separable Carcass Muscle on Combinations of Objective Carcass Measurements and Flank Separable Components. Due to the ease of separation of the wholesale flank and the economy both in time and economic loss of product, regression equations using the flank separable components for prediction of total carcass separable muscle were calculated and appear in tables 47, 48 and 49 for the combined weight groups, the 500 to 550 lb. weight group, and the 700 to 750 lb. weight group, respectively. These equations show a high degree of accuracy for

Multiple regression equations for estimating pounds and percent total carcass separable muscle using objective carcass measurements and flank separable components as predictors, and the coefficient of determination for each equation (combined weight groups). Table 47.

	בכבביייי דוו בשבון בלתשרדונון (במוחדוובת אבדעור פוסחקיי	7-0 664 - 40-4	6.00000
70000		COEILICIENC	Scandard
Equation		OI	error or
INO.	ה בארדוואורדווא המתשרדונות ביים ווארדוואורדונה המתשרדונות ביים ווארדוואורדונה המתשרדונות ביים ווארדונה המתשרדו	nere iminarion	езгтшаге
-1	$\hat{\mathbf{Y}} = 12.715 + 0.259(\mathbf{x}_1)276(\mathbf{x}_2)119(\mathbf{x}_3)393(\mathbf{x}_4) + 0.893(\mathbf{x}_5)595(\mathbf{x}_6)$	96.0	5.16
7	$\hat{Y} = -45.567 + 0.259(x_1)281(x_2)123(x_3)379(x_4) + 1.480(x_5)$	96.0	5.15
က	$\sqrt{Y} = -48.794 + 0.255(x_1)270(x_2)387(x_4) + 1.566(x_5)$	96.0	5.15
4	$\hat{Y}_1$ = 55.174 + 0.001( $X_1$ )137( $X_2$ )031( $X_3$ )120( $X_4$ ) + 0.289( $X_5$ )185( $X_6$ )	0.88	1.51
5	$\sqrt{Y_1}$ = 37.029 + 0.001( $X_1$ )138( $X_2$ )032( $X_3$ )116( $X_4$ ) + 0.472( $X_5$ )	0.87	1.51
9	$\sqrt{Y_1}$ = 36.190 + 0.0003( $X_1$ )136( $X_2$ )118( $X_4$ ) + 0.494( $X_5$ )	0.87	1.51
	ightarrow  m Y = Estimated pounds of total carcass separable muscle.		
	$\overline{Y_1}$ = Estimated percent total carcass separable muscle.		
	$X_1$ = Carcass weight (1b.).		
	$\rm X_2$ = Probe - 5th thoracic vertebra, 4 in. off the carcass dorsal midline (mm).		
	$x_3$ = Probe - 1st lumbar vertebra, 12 in. off the carcass dorsal midline (mm).		
	${ m X_4}$ = Probe - Av. of 3 at the 3rd sacral vertebra, 4, 8 and 12 in. off the carcass dorsal midline (mm).	ss dorsal midl	ine (mm).
	$X_5$ = Percent flank separable muscle.		

 $X_6$  = Percent flank separable fat.

Multiple regression equations for estimating pounds and percent total carcass separable muscle using objective carcass measurements and flank separable components as predictors, and the coefficient of determination for each equation (500 to 550 lb. weight group) Table 48.

To:ret		Coefficient	Standard
No.	Estimating equation	de termination	estimate
1	$ \frac{\lambda}{Y} = 73.831 + 0.202(x_1)324(x_2) + 0.027(x_3)356(x_4) + 0.415(x_5)848(x_6) $	0.86	4.59
2	$\dot{X} = -15.535 + 0.210(x_1)320(x_2) + 0.048(x_3)333(x_4) + 1.280(x_5)$	0.85	4.64
က	$\hat{Y} = -14.168 + 0.210(x_1)298(x_2)331(x_4) + 1.258(x_5)$	0.85	4.58
4	$\hat{Y}_1 = 78.524 = .020(x_1)151(x_2)0005(x_3)120(x_4) + 0.144(x_5)296(x_6)$	0.88	1.70
5	$\hat{Y}_1 = 47.351017(x_1)149(x_2) + 0.007(x_3)112(x_4) + 0.445(x_5)$	0.88	1.71
9	$\hat{Y}_1 = 47.545017(x_1)146(x_2)112(x_4) + 0.442(x_5)$	0.88	1.68
	$\Lambda = { m Estimated \ pounds \ of \ total \ carcass \ separable \ muscle.}$		
	V = Dottmoted norecant total caroace constable misself		

 $\mathbf{Y}_1$  = Estimated percent total carcass separable muscle.

 $X_1$  = Carcass weight (1b.).

 $X_2$  = Probe - 5th thoracic vertebra, 4 in. off the carcass dorsal midline.

 $x_3$  = Probe - 1st lumbar vertebra, 12 in. off the carcass dorsal midline.

 $X_4$  = Probe - Av. of 3 probes at 3rd sacral vertebra, 4, 8 and 12 in. off the carcass dorsal midline.

 $X_5$  = Percent flank separable muscle.

 $X_6$  = Percent flank separable fat.

Multiple regression equations for estimating pounds and percent total carcass separable muscle using objective carcass measurements and flank separable components as predictors, and the coefficient of determination for each equation (700 to 750 lb. weight group) Table 49.

1 4 6 1		Coefficient	Standard
No.	Estimating equation	or determination	estimate
H	$A = -191.412 + 0.177(x_1)555(x_2)252(x_3)466(x_4) + 3.795(x_5) + 2.062(x_6) 0.91$	0.91	4.12
2	$\vec{Y} = 14.891 + 0.163(x_1)469(x_2)154(x_3)483(x_4) + 1.848(x_5)$	0.91	4.17
က		0.91	4.17
4	$\dot{X}_1 = 5.209014(X_1)193(X_2)073(X_3)102(X_4) + 0.931(X_5) + 0.451(X_6)$	06.0	1.25
5	$\hat{Y}_1 = 50.334018(X_1)174(X_2)052(X_3)106(X_4) + 0.505(X_5)$	06.0	1.25
9	$\hat{Y}_1 = 50.232022(x_1)152(x_2)116(x_4) + 0.548(x_5)$	0.89	1.26
	$\Lambda$ = Estimated pounds of total carcass separable muscle.		
	$\stackrel{ extstyle \wedge}{Y_1}$ = Estimated percent total carcass separable muscle.		
	$X_1$ = Carcass weight (1b.).		

 ${\rm X}_2$  = Probe - 5th thoracic vertebra, 4 in. off the carcass dorsal midline.

 $x_3$  = Probe - 1st lumbar vertebra, 12 in. off the carcass dorsal midline.

 $X_4$  = Probe - Av. of 3 probes at the 3rd sacral vertebra, 4, 8 and 12 in. off the carcass dorsal midline.

X<sub>5</sub> = Percent flank separable muscle.

 $X_6$  = Percent flank separable fat.

predicting either pounds or percent carcass separable muscle (coefficient of determination range, 0.88 to 0.96). Slightly greater accuracy of prediction was obtained in the 700 to 750 lb. weight group, which is in contrast with most other prediction equations presented. These data suggest that the separable components of the flank are of greater predictive value in heavier carcasses than in lighter carcasses.

Multiple Regression Analyses of Total Carcass Separable Fat on Objective Carcass Measurements and Percent Wholesale Cut Separable Fat. Carcass weight, fat probe at the 5th thoracic vertebra (4 in. off the carcass dorsal midline) and separable fat from the 9-10-11 rib section, flank and round were used in multiple regression analyses as shown in tables 50, 51 and 52.

In the combined weight groups (table 50), 9-10-11 rib section separable fat included in the equation was the best predictor of carcass separable fat (pounds and percent) followed closely by flank separable fat and round separable fat (coefficient of determination range, 0.85 to 0.95). This same order was observed in the 500 to 550 lb. weight group (table 51), but in the 700 to 750 lb. weight group, flank separable fat was essentially comparable to 9-10-11 rib and round separable fat for predicting percent total carcass separable fat (coefficient of determination range, 0.85 to 0.87)(table 52). The coefficients of determination were very similar within weight groups, but it can be seen from the equations shown in tables 51 and 52 that greater accuracy can be obtained in predicting either pounds or percent carcass separable fat in 500 to 550 lb. weight carcasses as opposed to 700 to 750 lb. weight carcasses.

Multiple regression equations for estimating pounds and percent total carcass separable fat using objective carcass measurements and percent wholesale cut separable fat as predictors, and the coefficient of determination for each equation (combined weight groups) Table 50.

Equation		Coefficient of	Standard error of
No.	Estimating equation	determination	estimate
H	$\Lambda = -82.782 + 0.179(x_1) + 0.896(x_2) + 1.850(x_3)$	0.95	5.64
2	$^{\wedge}_{Y} = -122.346 + 0.173(x_1) + 1.174(x_2) + 1.821(x_4)$	0.94	6.55
က	$^{\wedge}_{Y}$ = -73.978 + 0.176( $x_1$ ) + 0.807( $x_2$ ) + 2.886( $x_5$ )	0.94	6.29
7	$^{\wedge}_{Y_1}$ = 7.9340005( $x_1$ ) + 0.303( $x_2$ ) + 0.629( $x_3$ )	0.91	1.58
5	$\hat{Y}_1 = -6.402003(x_1) + 0.380(x_2) + 0.637(x_4)$	0.88	1.83
9	$\dot{Y}_1 = 11.634001(x_1) + 0.308(x_2) + 0.932(x_5)$	0.85	2.05
<> =	= Estimated noimds of carcass senarable fat.		

I = ESTIMATED pounds or carcass separable rat.

 $\Lambda_1$  = Estimated percent carcass separable fat.

 $X_1$  = Carcass weight (1b.)

 $X_2$  = Probe - 5th thoracic, 4 in. off carcass dorsal midline (mm.).

 $X_3$  = Percent 9-10-11 rib section separable fat.

 $X_4$  = Percent flank separable fat.

 $X_5$  = Percent round separable fat.

Multiple regression equations for estimating pounds and percent total carcass separable fat using objective carcass measurements and percent wholesale cut separable fat as predictors, and the coefficient of determination for each equation (500 to 550 lb. weight group). Table 51.

		Coefficient	Standard
uation	• • • • • • • • • • • • • • • • • • • •	of	error of
Į.	Estimating equation	determination	estimate
<b>~</b> 1	$\sqrt[4]{}$ = -126.461 + 0.284(x <sub>1</sub> ) + 0.791(x <sub>2</sub> ) + 1.559(x <sub>3</sub> )	0.95	3.82
7	$\hat{Y} = -164.739 + 0.299(x_1) + 1.074(x_2) + 1.430(x_4)$	0.92	5.07
ന	$^{\wedge}_{Y}$ = -173.335 + 0.380( $x_1$ ) + 0.868( $x_2$ ) + 2.453( $x_5$ )	0.92	5.01
4	$\hat{Y}_{1}$ = -10.727 + 0.036( $x_1$ ) + 0.300( $x_2$ ) + 0.606( $x_3$ )	0.94	1.46
5	$\hat{Y}_{1}$ = -26.506 + 0.042( $x_{1}$ ) + 0.382( $x_{3}$ ) + 0.580( $x_{4}$ )	0.91	1.81
9	$\Lambda_{1}$ = -29.546 + 0.074( $X_1$ ) + 0.312( $X_2$ ) + 0.977( $X_5$ )	0.91	1.84
$\hat{\mathbf{Y}} = \mathbf{Est}$	stimated pounds of total carcass separable fat.		

 $X_1$  = Estimated percent total carcass separable fat.

 $X_1$  = Carcass weight (1b.).

 $\rm X_2$  = Probe - 5th thoracic vertebra, 4 in. off the carcass dorsal midline (mm.).

 $x_3$  = Percent 9-10-11 rib section separable fat.

 $X_4$  = Percent flank separable fat.

 $X_5$  = Percent round separable fat.

Multiple regression equations for estimating pounds and percent total carcass separable fat using objective carcass measurements and percent wholesale cut separable fat as predictors, and the coefficient of determination for each equation (700 to 750 lb. weight group). Table 52.

		Coefficient	Standard
Equation No.	Estimating equation	ot determination	error of estimate
1	$^{\wedge}_{Y}$ = -96.248 + 0.176( $x_1$ ) + 1.005( $x_2$ ) + 2.194( $x_3$ )	0.86	6.47
2	$\hat{Y} = -198.084 + 0.228(x_1) + 1.310(x_2) + 2.360(x_4)$	0.85	92.9
က	$\hat{Y} = -71.687 + 0.168(x_1) + 0.366(x_2) + 3.272(x_5)$	0.85	69.9
7	$\hat{Y}_1 = 5.846 + 0.003(x_1) + 0.266(x_2) + 0.622(x_3)$	0.87	1.67
5	$\hat{Y}_1$ = -23.084 + 0.018( $x_1$ ) + 0.352( $x_2$ ) + 0.670( $x_4$ )	0.85	1.76
9	$\hat{Y}_1$ = 12.925 + 0.0005( $x_1$ ) + 0.076( $x_2$ ) + 0.940( $x_5$ )	0.87	1.68

 $\Lambda$  = Estimated pounds of total carcass separable fat.

 $\gamma_1$  = Estimated percent total carcass separable fat.

 $X_1$  = Carcass weight (1b.).

 $\rm X_2$  = Probe - 5th thoracic vertebra, 4 in. off the carcass dorsal midline.

 $X_3$  = Percent 9-10-11 rib section separable fat.

 $X_4$  = Percent flank separable fat.

 $X_5$  = Percent round separable fat.

Multiple Regression Analyses of Total Carcass Separable Bone on Objective Carcass Measurements and Percent Wholesale Cut Separable Components. Percent 9-10-11 rib section bone, percent round bone and percent flank separable muscle were used in combination with carcass weight and length of round to calculate multiple regression equations for the prediction of total carcass separable bone. Flank separable muscle was used rather than flank separable bone since the percent of flank separable bone was low and showed a very low relationship to total carcass separable bone.

In the combined weight groups (table 53) there was little difference in the predictive value of any of the equations for either pounds or percent separable carcass bone (coefficient of determination range, 0.91 to 0.92 and 0.79 to 0.83 for pounds and percents, respectively).

In the 500 to 550 lb. weight group (table 54), the equations for predicting pounds and percent carcass separable bone using flank separable muscle showed the most accurate predictive values (0.87 and 0.89 for pounds and percents, respectively). The coefficients of determinations and standard error of estimates for the 700 to 750 lb. weight group were similar to those of the combined weight groups. These data indicate that pounds and percent carcass separable bone can most accurately be predicted in lighter weight carcasses.

Multiple regression equations for estimating pounds and percent total carcass separable bone using objective carcass measurements and percent wholesale cut separable components as predictors, and the coefficient of determination for each equation (combined weight groups). Table 53.

Farretton		Coefficient	Standard
No.	Estimating equation	determination	estimate
1	$^{A}_{1} = -48.402 + 0.026(x_{1}) + 1.909(x_{2}) + 0.942(x_{3})$	0.91	1.91
7	$\hat{Y} = -47.633 + 0.025(x_1) + 1.946(x_2) + 0.309(x_4)$	0.92	1.79
က	$\hat{Y} = -44.629 + 0.031(x_1) + 1.200(x_2) + 1.887(x_5)$	0.92	1.81
4	$\hat{Y}_1$ = -4.833010(x <sub>1</sub> ) + 0.634(x <sub>2</sub> ) + 0.325(x <sub>3</sub> )	0.81	0.61
53	$\hat{Y}_1 = -4.569010(x_1) + 0.650(x_2) + 0.105(x_4)$	0.83	0.57
9	$\hat{Y}_1 = -3.705010(x_1) + 0.466(x_2) + 0.554(x_5)$	0.79	0.64

 $\dot{\mathbf{Y}} = \mathbf{Estimated}$  pounds of carcass separable bone.

 $Y_1$ = Estimated percent carcass separable bone.

 $X_1$  = Carcass weight (1b.).

 $X_2$  = Length of round (in.).

 $x_3$  = Percent 9-10-11 rib section separable bone.

 $X_4$  = Percent flank separable muscle.

 $X_5$  = Percent round separable bone.

Multiple regression equations for estimating pounds and percent total carcass separable bone using objective carcass measurements and percent wholesale cut separable components as predictors, and the coefficient of determination for weight group) each equation (500 to 550 lb. Table 54.

Fanation		Coefficient	Standard error of
No.	Estimating equation	determination	estimate
н	$^{\wedge}_{Y}$ = -25.202 + 0.001(x <sub>1</sub> ) + 1.600(x <sub>2</sub> ) + 0.841(x <sub>3</sub> )	0.79	1.81
7	$\hat{Y} = -37.887 + 0.024(x_1) + 1.576(x_2) + 0.334(x_4)$	0.87	1.42
ო	$\sqrt{Y} = -11.152014(X_1) + 0.646(X_2) + 2.291(X_5)$	0.83	1.60
4	$\vec{Y}_1$ = 1.435020 ( $x_1$ ) + 0.613( $x_2$ ) + 0.308( $x_3$ )	0.83	0.67
'n	$\hat{Y}_1 = -3.160012(x_1) + 0.611(x_2) + 0.120(x_4)$	0.89	0.56
9	$ \sqrt{Y_1} = 6.493026(X_1) + 0.272(X_2) + 0.829(X_5) $	0.86	0.61
X = Est	$\stackrel{\textstyle \wedge}{\rm Y}$ = Estimated pounds of total carcass separable bone.		

 $\vec{\mathbf{Y}}_1$  = Estimated percent total carcass separable bone.

 $X_1$  = Carcass weight (1b.).

 $X_2$  = Length of round (in.).

 $X_3$  = Percent 9-10-11 rib section separable bone.

 $X_4$  = Percent flank separable muscle.

 $X_5$  = Percent round separable bone.

Multiple regression equations for estimating pounds and percent total carcass separable bone using objective carcass measurements and percent wholesale cut separable components as predictors, and the coefficient of determination for each equation (700 to 750 lb. weight group) Table 55.

		Coefficient	Standard
Equation		Jo	error of
No.	Estimating equation	determination	estimate
1	$\hat{Y} = -63.479 + 0.039(x_1) + 1.994(x_2) + 1.209(x_3)$	08.0	1.96
2	$\sqrt{Y} = -71.717 + 0.044(X_1) + 2.300(X_2) + 0.302(X_4)$	0.79	2.04
ო	$\hat{Y} = -64.258 + 0.065(x_1) + 1.019(x_2) + 1.906(x_5)$	0.83	1.82
7	$\hat{\mathbf{Y}}_1$ = -7.921003( $\mathbf{x}_1$ ) + 0.564( $\mathbf{x}_2$ ) + 0.344( $\mathbf{x}_3$ )	0.80	0.53
۳	$\hat{Y}_1 = -10.200002(x_1) + 0.657(x_2) + 0.082(x_4)$	0.77	0.57
9	$\hat{Y}_1$ = -8.164 + 0.004(x <sub>1</sub> ) + 0.306(x <sub>2</sub> ) + 0.522(x <sub>5</sub> )	0.81	0.51
$\stackrel{\wedge}{Y} = Es($	= Estimated pounds of total carcass separable bone.		

 $\stackrel{\wedge}{\gamma}_1$  = Estimated percent total carcass separable bone.

 $X_1$  = Carcass weight (1b.).

 $X_2$  = Length of round (in.).

 $x_3$  = Percent 9-10-11 rib section separable bone.

= Percent flank separable muscle.

 $X_5$  = Percent round separable bone.

Relationships Between Weight of Certain Entire Muscles and Muscle Groups and Weight of Total Separable Carcass Muscle and Bone

Weight of the <u>1</u>. <u>dorsi</u> muscle showed the highest relationship to pounds of total carcass muscle in the combined weight groups and in the 500 to 550 lb. weight group (0.90 and 0.70, respectively) (table 56). In the 700 to 750 lb. weight group, however, the highest relationship to total carcass muscle was shown by the <u>semimembranosus</u> plus <u>adductor</u> (0.71), followed by the biceps femoris (0.63) and the l. dorsi muscle (0.61).

Correlation coefficients between muscle weights and total carcass muscle weight were markedly lower when calculated on a within weight group basis, particularly for the <u>quadriceps</u> muscles (range, 0.20 to 0.40). All other correlations between individual and muscle group weights and total carcass muscle were highly significant (P < .01)(range, 0.45 to 0.90). Although correlations were slightly lower than for individual and muscle group weights, essentially the same relationships were observed between these muscle weights and total bone weight.

Relationships Between Weights of Certain Entire Bones and Bone Groups and
Weight of Total Separable Carcass Muscle and Bone

Highly significant correlation coefficients (P < .01) were found between weight of the radius plus ulna, femur, tibia plus fibula, scapula and humerus and total carcass bone weight for the combined weight groups and within weight groups and are shown in table 57 (range, 0.62 to 0.91). Correlations between individual and bone group weights and total carcass muscle ranged from 0.24 to 0.85 and with the exception of the radius plus ulna and tibia plus fibula in the 500 to 550 lb. weight group all were highly significant (P < .01). These correlation coefficients were similar to those reported by Orme (1963) in lambs.

Simple correlation coefficients between some individual muscle and muscle group weights and carcass separable muscle and bone. Table 56.

	Combined w	Combined weight groups	500 to 550 lb.	550 1b.	700 to 750 lb.	750 1b.
			weight	weight group	weight	weight group
	Total	Total	Total	Total	Tota1	Total
Muscles:	separable muscle	separable bone	separable muscle	separable bone	separable muscle	separable bone
Semitendinosus	0.86	0.76	0.57	0.28	0.72	0.54
Quadriceps	0.64	0.63	0.20	0.21	0.34	0.40
Semimembranosus and Adductor	0.87	0.84	99.0	0.53	0.71	0.68
Biceps femoris	0.87	0.84	09.0	0.56	0.63	0.63
Psoas major	0.75	0.64	0.45	0.12	0.52	0.40
L. dorsi	06.0	0.78	0.70	0.16	0.61	0.54
Correlations > 0.2	220 are signi	220 are significant (P < .05) for the combined weight groups	5) for the	combined weig	ht groups.	

for the combined weight groups. for the combined weight groups. for the individual weight groups. for the individual weight groups. (P < .01) (P < .01) (P < .05) Correlations > 0.220 are significant Correlations > 0.287 are significant Correlations > 0.312 are significant Correlations > 0.403 are significant

Simple correlation coefficients between individual bone and bone group weights and carcass separable muscle and bone. Table 57.

and card	cass separao	CERSS SEPAIRDIE MUSCIE AND DONE	500 to 550 lb.	550 1b.	700 to 750 lb.	750 lb.
	Combined w	Combined weight groups	weight	weight group	weight group	group
	Total	Total	Total	Total	Total	Total
Bones:	separable muscle	separable bone	separable muscle	separable bone	separable muscle	separable bone
Radius + Ulna	0.79	0.85	0.24	0.63	0.42	0.62
Femur	0.79	06.0	0.41	0.87	0.52	0.72
Tibia + Fibula	0.77	06.0	0.39	0.78	0.52	0.81
Scapula	0.85	0.91	99.0	0.88	0.58	0.71
Humerus	0.78	0.84	0.50	0.74	0.46	0.62

for the combined weight groups. for the individual weight groups. for the individual weight groups. for the combined weight groups. Correlations > 0.220 are significant (P < .05) Correlations > 0.287 are significant (P < .01) Correlations > 0.312 are significant (P < .05) Correlations > 0.403 are significant (P < .05) The highest relationship in the combined weight group and the 500 to 550 lb. weight group was observed between scapula weight and total bone weight (0.91 and 0.88, respectively). In the 700 to 750 lb. weight group, the highest relationship was observed between tibia plus fibula weight and total bone weight (0.81). Relationships between weights of these bones and total carcass bone weight were sufficiently high for use in prediction equations. These data support the findings of Butterfield (1963b) who developed several prediction equations for separable carcass bone using the weights of various bones as the independent variables in his equations.

Multiple Regression Analyses of Pounds and Percent Separable Carcass Muscle and Bone on Individual and Groups of Muscle and Bone Weights

Multiple Regression Analyses of Pounds and Percent Separable Carcass Muscle on Muscle and Bone Weights. In the combined weight groups, 94% of the variation in pounds of separable carcass muscle could be accounted for using the independent variables shown in table 58. The same five variables only accounted for 78% of the variation in percent separable carcass muscle in the combined weight groups with a standard error of estimate of 1.98% (suppl., p. 179).

Eighty-eight percent of the variation was accounted for in pounds of separable carcass muscle in the 500 to 550 lb. weight group using the same variables that were included in the combined weight group analysis plus weight of the femur, tibia and fibula and the humerus (suppl., p. 179). However, in this weight group, when the five variables used in

Table 58. Coefficient of determination between weight of carcass separable muscle and a combination of muscle and bone weights, and the regression coefficient for each of the variables (combined weight groups)

groups).				
	Regression	Beta	Level of	R <sup>21</sup>
Independent variables	coefficients	weights	sig.	deletes
Constant	4.978		0.31	
Carcass wt.	0.049	0.184	0.00	0.93
Semitendinosus wt.	10.055	0.259	0.00	0.92
Biceps femoris wt.	5.934	0.247	0.00	0.92
L. dorsi wt.	2.899	0.202	0.00	0.94
Scapula wt.	12.652	0.189	0.00	0.93

 $R^2 = 0.94$  Standard error of estimate = 6.21

the combined weight group analysis were included, 88% of the variation in percent separable carcass muscle was accounted for (table 59). Using the multiple regression equation presented in this table to estimate percent separable carcass muscle resulted in a standard error of estimate of 1.65%. Weight of the <u>1</u>. dorsi muscle had the greatest effect upon percent separable carcass muscle of any single variable in this prediction equation. This is probably attributable to the part to whole relationship.

Estimation of pounds and percent separable carcass muscle from muscle and bone weights in the 700 to 750 lb. weight groups resulted in reduced accuracy. Carcass weight and those variables with a significant (P < .05) effect on pounds and percent separable carcass muscle resulted in coefficients of determination of 0.73 for pounds of separable carcass muscle

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

Table 59. Coefficient of determination between percent of carcass separable muscle and a combination of muscle and bone weights, and the regression coefficient for each of the variables (500 to

550 lb. wei	ght group).			
	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	51.336		0.00	
Carcass wt.	<b>0</b> 86	312	0.00	0.80
Semitendinosus wt.	2.748	0.229	0.00	0.84
Biceps femoris wt.	2.154	<b>0.2</b> 56	0.00	0.83
L. dorsi wt.	1.870	0.360	0.00	0.78
Scapula wt.	6.236	0.335	0.00	0.80

 $R^2 = 0.88$  Standard error of estimate = 1.65

(suppl., p. 180) and 0.77 for percent of separable carcass muscle (suppl., p. 180).

These results support those of Orme et al. (1960) who presented simple linear regression equations for the prediction of pounds of separable carcass muscle from muscle weights. The standard error of estimate of the multiple regression equations presented here were less in most instances than those presented by Orme et al. (1960).

Multiple Regression Analyses of Pounds and Percent Separable Carcass Bone on Muscle and Bone Weights. In the combined weight groups, accurate prediction equations for either pounds or percent separable carcass bone were calculated from muscle and bone weights. Six independent variables, listed in table 60, were responsible for 94% of the variation in pounds of separable carcass bone.

<sup>1</sup> Value of coefficient of determination if that variable was deleted from the analysis.

Table 60. Coefficient of determination between weight of separable carcass bone and a combination of muscle and bone weights, and the regression coefficient for each of the variables (combined weight groups)

groups).				
	Regression	Bet <b>a</b>	Level of	R <sup>21</sup>
Independent variables	coefficients	weights	sig.	deletes
Constant	814		0.53	
Carcass wt.	0.0002	0.002	0.91	0.94
<u>Semitendinosus</u> wt.	0.961	0.098	0.04	0.94
Biceps femoris wt.	0.806	0.133	0.01	0.94
Femur wt.	1.779	0.172	0.01	0.94
Tibia & Fibula wt.	3.843	0.302	0.00	0.92
Scapula wt.	6.036	0.359	0.00	0.91

 $R^2 = 0.94$  Standard error of estimate = 1.58

With one less variable (Semitendinosus weight), 88% of the variation in percent separable carcass bone was accounted for (standard error of estimate of 0.50%)(table 61). In contrast to the analysis for pounds of separable carcass bone, carcass weight has a highly significant (P < .01) effect upon percent separable carcass bone in the combined weight groups.

Ninety-five and 96% of the variation in pounds and percent of separable carcass bone, respectively, can be accounted for in the 500 to 550 lb. weight group. These prediction equations had a high degree of accuracy with a standard error of estimate of 0.92 lb., using five variables, to estimate pounds of carcass bone (suppl., p. 180) and a standard error of estimate of 0.33%, using six variables, to predict percent separable carcass bone (suppl., p. 181).

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

Table 61. Coefficient of determination between percent carcass separable muscle and a combination of muscle and bone weights, and the regression coefficient for each of the variables (combined weight groups)

weight group	s).			
Independent variables	Regression coefficients	Bet <b>a</b> weights	Level of sig.	R <sup>21</sup> deletes
Constant	11.172		0.00	
Carcass wt.	018	-1.239	0.00	0.31
Biceps femoris wt.	0.224	0.172	0.03	0.87
Femur wt.	0.806	0.362	0.00	0.85
Tibia + Fibula wt.	1.200	0.437	0.00	0.83
Scapula wt.	2.175	0.600	0.00	0.78

 $R^2 = 0.88$  Standard error of estimate = 0.50

In the 700 to 750 lb. group, the prediction equations were less accurate. For pounds of separable carcass bone, 85% of the variation was accounted for using four independent variables (suppl., p. 181).with a standard error of estimate of 1.73 lb. For percent separable carcass bone, 88% of the variation was accounted for using five independent variables (suppl., p. 181) with a standard error of estimate of 0.42%.

## Relationships Between Weight and Percent

## Total Muscle, Fat and Bone

Correlation coefficients between weights and percents of total carcass separable muscle, fat and bone for the combined and individual weight
groups appear in table 62. Correlations between percent muscle and percent fat were very high (-.98) on a combined and within weight group basis.
The correlations between percent fat and percent bone were also high (-.84).

<sup>1</sup> Value of coefficient of determination if that variable was deleted from the analysis.

Positive correlations were observed between percent muscle and bone (0.74, 0.73 and 0.75 for the combined, 500 to 550 lb. and 700 to 750 lb. weight groups, respectively).

Table 62. Simple correlation coefficients between weights and percents of total carcass separable muscle, fat and bone for the combined and individual weight groups.

Separ	able	Separ	able	Separ	able
musc	1e	fa	t	<u>bon</u>	.e
1b.	%	1b.	%	1b.	%
1.00	1.00	0.40	98	0.90	0.74
		1.00	1.00	0.28	84
				1.00	1.00
1.00	1.00	82	98	0.65	0.73
		1.00	1.00	79	84
				1.00	1.00
1.00	1.00	85	<b></b> 98	0.78	0.75
		1.00	1.00	65	84
				1.00	1.00
	1.00	1.00 1.00	muscle     fa       1b.     %       1.00     1.00       1.00     1.00       1.00     1.00       1.00     1.00      82     1.00       1.00     1.00      85	muscle     fat       1b.     %       1.00     1.00       1.00     1.00       1.00     1.00       1.00     1.00       1.00     1.00	muscle         fat         bon           1b.         %         1b.           1.00         1.00         0.40        98         0.90           1.00         1.00         0.28         1.00           1.00         1.00        82        98         0.65           1.00         1.00        79         1.00           1.00         1.00        65        65

Correlations > 0.220 are significant (P < .05) for combined weight groups. Correlations > 0.287 are significant (P < .01) for combined weight groups. Correlations > 0.312 are significant (P < .05) for individual weight groups. Correlations > 0.403 are significant (P < .01) for individual weight groups.

Correlations involving weight of separable components were lower than for percents; however, all, except that between weight of total carcass fat and bone for the combined weight groups, were highly significant (P < .01). These relationships are similar to the results reported by Brown et al. (1962), Hedrick et al. (1963) and Miller et al. (1965). These authors reported high negative relationships between carcass fat and muscle and positive relationships between carcass bone and muscle.

Relationships Between Wholesale Cut Retail Yield and Total Carcass Separable

Components, Retail and Fat Trim Yields

Simple correlation coefficients between pounds and percent wholesale cut retail yields and pounds and percent separable components, retail and fat trim yields on a combined and within weight group basis appear in tables 63, 64 and 65.

The highest relationships between pounds of wholesale cut retail yields and pounds of carcass separable muscle, boned, trimmed round, loin, rib and chuck yield and total retail yield were the chuck and round yields (range, 0.60 to 0.95). These relationships existed for the combined as well as for the within weight groups. In general, pounds of wholesale cut retail yield was positively correlated with pounds and percent carcass muscling and bone and on a within weight group basis, negatively related to pounds and percent separable carcass fat and fat trim yields.

On a percent basis in the combined weight groups, the relationships between wholesale cut retail yield and separable carcass muscle and bone were lower than for weight. Correlations between percent wholesale cut retail yields and carcass separable fat were negative and higher than weight correlations. However, correlation coefficients between percent and weights of retail yield from the round, loin, rib and chuck and total retail yield were quite similar.

Correlations between percent retail yield of the flank and percent round, loin, rib and chuck yield (range, 0.72 to 0.82) and total retail yield (range, 0.84 to 0.89) were higher than those for weight. On a percent basis in all groups, correlations between retail yield from the

Table 63. Simple correlation coeffici	Simple	correl	ation c			etween	wholesa	le cut 1	cetail y	ield and	carcas	38 Sepa	ents between wholesale cut retail yield and carcass separable components,	ponents,
	retail	and fa	retail and fat trim yields		for the	e combi	for the combined weight groups.	ght grou	ъв.			•		•
	Total	al	Total	al	Total	al	Boned,	ed,	Boned	ed,				
Wholesale	separable	able	separable	able	separable	able	trimmed	ned	trimmed	ned			External fat	1 fat
cut retail	muscle	le	fat	t)	bone	اره	retail RLRC1	RLRC1	retail yield	yield	Total fat	fat	trim from RLRC	m RLRC
yield	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%
Round	0.90	0.90 0.55	0.39	57	0.87	0.42	0.94	08.0	0.95	0.85	0.24	82	0.11	<b>-</b> .83
Loin	0.81	0.81 0.66	0.48	69	0.75	0.52	0.88	0.85	0.87	0.86	0.29	86	0.16	87
Rib	0.75	0.75 0.56	0.40	60	0.63	0.50	0.84	0.84	0.83	0.88	0.18	- 88	0.03	85
Chuck	0.92	0.92 0.59	0.52	62	0.82	0.49	0.97	0.91	0.95	0.89	0.38	88	0.24	•. 82
Foreshank	0.40	0.40 0.29	03	-,35	0.51	0.34	0.45	0.52	0.52	0.64	19	63	20	i 69
Brisket	0.59	0.59 0.61	0.20	62	0.55	0.44	09.0	0.76	0.67	0.81	90.0	80	0.02	. 48
Plate	0.65	0.65 0.42	0.47	42	0.55	0.28	0.73	0.57	0.77	0.65	0.28	60	0.20	49
Flank	0.56	0.56 0.73	0.34	75	0.46	0.57	0.67	0.81	0.73	0.88	0.09	87	90.0	73

Correlations > 0.220 are significant (P < .05). Correlations > 0.287 are significant (P < .01). IRLRC = Round, loin, rib and chuck.

Table 64. Simple correlation coefficients between wholesale cut retail yield and carcass separable components, retail and fat trim yields for the 500 to 550 lb. weight group.	Simple retail	correla	ation c t trim	oeffici yields	ents b for th	etween e 500 t	wholesa	le cut 1 b. weigh	Simple correlation coefficients between wholesale cut retail yiretail and fat trim yields for the 500 to 550 lb. weight group.	ield and	carcat	ss sepa	rable com	nponents,
Wholesale	Total	al able	Total separable	al	Total separable	al	Boned,	ed,	Boned,	ed,			External fat	al fat
cut retail	muscle	le .1	fat	الد	bone	اله	retail RLRC1	RLRC1	retail yield	yield	Total fat	fat	trim from RLRC	M RLRC
yield	Ib.	%	Ib.	%	Ib.	%	Ib.	%	Ib.	%	Ib.	%	1b.	%
Round	0.73	0.73 0.57	75	59	0.74 0.43	0.43	0.88	0.83	0.88	0.88	82	84	83	86
Loin	0.40	0.40 0.72	36	3674	0.37	0.55	0.67	0.89	0.72	0.91	60	90	63	888.
RID	0.46	0.46 0.62	21	65	0.10	0.54	0.71	0.86	0.69	0.92	50	92	54	06
Chuck	0.71	0.71 0.64	46	<b>-</b> . 68	0.44	0.57	0.81	06.0	0.71	0.92	56	92	57	87
Foreshank	0.36	0.36 0.40	55	48	0.63	0.54	97.0	0.55	0.58	0.67	-, 64	71	58	99
Brisket	0.28	0.28 0.63	36	65	0.32	0.53	0.36	0.76	0.50	0.84	48	84	39	73
Plate	0.24	0.24 0.52	15	52	0.09	0.40	0.53	0.70	09.0	0.74	-,45	71	50	65
Flank	0.32	0.32 0.78	34	81	0.25	0.67	0.57	0.82	0.70	0.89	64	. 88	57	77

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01). IRLRC = Round, loin, rib and chuck.

1
!
١
- (
)
1
Ì
1

Table 65. Simple correlation coefficients between wholesale cut retail yield and carcass separable components, retail and fat trim yields for the 700 to 750 lb. weight group.	Simple retail	correl	ation cut trim	oeffici yields	ents b	etween e 700 t	wholesa to 750 11	le cut 1 b. weigh	Simple correlation coefficients between wholesale cut retail yiretail and fat trim yields for the 700 to 750 lb. weight group.	ield and .	carca	ss seps	rable co	mponents,
	Tot	Total	Total	a1	Total	al	Boned	ed,	Boned	ed,				
Wholesale	sepaı	separable	separable	able	separable	able	trimmed	ned	trimmed	med	Total fat	fat	External fat	al fat
cut retail	muscle	:1e	fat	ايد	bone		retail RLRC1	RLRC1	retail yield	yield	trim		trim from RLRC	om RLRC
yield	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%
Round	09.0	0.60 0.50	51	51	0.56 0.39	0.39	0.74	99.0	0.76	0.76	65	73	51	73
Loin	0.35	0.35 0.58	22	60	0.35	0.48	09.0	0.71	0.50	0.70	43	76	30	81
Rib	0.44	0.44 0.45	22	48	0.35	0.43	0.71	0.74	0.68	0.71	47	72	45	67
Chuck	0.62	0.49	37	48	0.49	0.33	06.0	0.82	0.86	0.81	60	74	50	52
Foreshank	0.26	0.26 0.02	20	20 0.04	0.22	17	0,39	0.22	0.52	0.41	37	28	20	03
Brisket	0.38	0.38 0.57	27	51	0.26 0.20	0.20	0.39	0.61	0.54	0.73	41	<b>-</b> . 68	28	46
Plate	0.12	0.12 0.22	0.03	18	90.0	0.02	0.28	0.14	97.0	0.31	13	22	0.03	01
Flank	0.13	0.13 0.65	0.11	-, 61	06	0.34	0.41	0.72	0.57	0.84	15	79	03	57

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01). 1 RLRC = Round, loin, rib and chuck.

flank and the seven carcass compositional components were essentially as high or higher than those from any other wholesale cut. These findings are similar to those reported by Hedrick et al. (1963) and Miller et al. (1965) who found the flank retail yield and carcass retail yield relationship quite high and included retail flank yield as an independent variable in some of their prediction equations. In contrast to these data, Brungardt and Bray (1963) reported that percent trimmed round and round retail yields were most highly related to carcass retail yields. However, correlations between separable muscle from the round and carcass retail yields differed greatly when comparing them on a combined and within weight group basis (tables 28, 29 and 30) while those between separable muscle of the flank and carcass retail yields were much more consistent in all groups.

Thus, because of the ease in obtaining flank retail yield and/or separable components and the economics involved, the wholesale flank (Wellington, 1952, cutting procedure) should be considered for use in estimating retail and/or fat trim yields. Percent retail yields of the wholesale cuts in the 500 to 550 lb. weight group showed higher relationships to percent carcass retail and fat trim yields than those in the 700 to 750 lb. weight group.

Relationships Between Pounds and Percent Wholesale Cut Fat Trim and Total

Carcass Separable Components, Retail and Fat Trim Yields

Correlations between pounds and percent wholesale cut fat trim yields and pounds and percent carcass separable components, retail and fat trim yields appear in tables 66, 67 and 68. Pounds and percents of separable

Simple correlation coefficients between wholesale cut fat trim and carcass separable components, retail and fat trim yields for the combined weight groups, Table 66.

	Total	Total Total	Total	1	Total	11	Total Boned,	d,	1	d,				
Wholesale	separable	able 1e	separable fat	able ⊤	separable hone	able	trimmed retail RLRC1	$_{ m RL,RC}^{ m l}$	trimmed retail vield	ned vield	Total fat	fat	External fat	fat RI.RC
cut fat trim 1b.	1 1b.	%	15.	%	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%
Round	0.26	0.2662	0.41 0.67	0.67	0.16	59	0.16	80	0.12	85	0.55	0.88	97.0	0.89
Loin	0.08	70	0.30 0.74	0.74	0.03	64	0.00	82	04	84	0.45	0.89	0.32	0.87
Rib	0.11	0.1166	0.58 0.71	0.71	02	64	0.04	86	01	89	0.76	0.94	0.66	0.88
Chuck	0.35	67	0.76 0.72	0.72	0.21	65	0.29	06	0.26	89	0.87	0.93	0.72	0.82
Foreshank	0.21	-,33	0.41	0.37	0.13	39	0.08	57	0.03	64	0.64	0.70	0.58	0.54
Brisket	0.23	70	0.71	0.73	0.20	62	0.22	74	0.20	86	0.81	0.88	0.72	0.70
Plate	0.55	0.5554	08.0	0.55	0.47	45	0.58	99	0.55	73	0.78	0.70	0.61	0.58
Flank	0.30	73	0.91	0.75	0.21	58	0.36	81	0.34	88	06.0	0.87	0.80	0.74
Kidney knob 0.2857	0.28	57	0.70 0.56	0.56	0.23	47	0.36	46	0.36	97	0.56	0.32	0.43	0.21

Correlations > 0.220 are significant (P < .05). Correlations > 0.289 are significant (P < .01). IRLRC = Round, loin, rib and chuck.

Table 67. Simple correlation coefficients between wholesale cut fat trim and carcass separable components	Simple	correl	ation c	oeffici	ents b	etween	wholesa	le cut 1	Eat trim	and cal	rcass s	eparabl	e compone	ints,
	retail	and fa	retail and fat trim yields	yields	for th	for the 500 t	to 550 lb. weight group.	b. weigh	it group	•				
	Total	al	Total	a1	Total		Boned	ed,	Boned	ed,				
Wholesale	separable	able	separable	able	separable	able	trimmed	ned	trimmed	med	Total fat	fat	External fat	1 fat
cut fat	muscle	:le	fa	fat	bone	- 1	retail	귐	retail yield	yield	trim		trim from RLRC	m RLRC
trim	Ib.	%	iè.	%	ie.	%	Ib.	%	lb.	%	1b.	%	Ib.	%
Round	26	2662		0.41 0.67	20	57	48	84	55	90	0.59	06.0	0.53	0.92
Loin	15	69		0.31 0.73	17	62	28	87	34	90	0.47	0.92	0.26	0.88
RIB	4	4467	0.58	0.58 0.72	52	65	60	87	70	93	0.83	96.0	0.72	0.91
Chuck	43	4367	0.66	0.66 0.73	60	68	62	88	70	91	0.88	0.95	0.75	0.89
Foreshank	31	45	0.41	0.41 0.47	38	48	55	60	64	70	0.73	0.77	0.72	0.74
Brisket	71	72	0.66	0.66 0.77	60	69	77	82	79	89	0.82	06.0	0.75	0.79
Plate	52	5259	0.67	0.67 0.61	56	54	55	76	55	78	0.74	0.77	0.62	0.70
Flank	74	7478		0.86 0.81	71	67	66	83	68	89	0.82	0.88	0.72	0.77
Kidney knob5570 0.74 0.71	-,55	70	0.74	0.71	57	64	29	-,53	26	47	0.47	0.42	0.32	0.29

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01).

RLRC = Round, loin, rib and chuck.

Table 68.	Simple	correl	Simple correlation coeffication to the state of the state	oeffici	ents b	ents between	wholesale cut fat trim	le cut i	fat trim	and car	cass s	eparabl	Simple correlation coefficients between wholesale cut fat trim and carcass separable components, retail and fat trim wields for the 700 to 750 lb weight grown	nts,
	Total	all al	Total	al	Total	al	Boned	ed,	Boned	ed,				
Wholesale	separable	able	separable	able	separable	able	trimmed	ned .	trimmed	ned	Total fat	fat	External fat	1 fat
cut fat	muscle	:le	fat	إب	bone	اه	retail RLRC	RLRC1	retail yield	yield	trim		trim from RLRC	m RLRC
trim	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%
Round	07	61		0.06 0.66	21	64	32	68	36	72	0.30	08.0	0.19	0.83
Loin	14	1469		0.24 0.74	15	68	33	71	40	70	0.47	0.84	0.32	0.87
Rib	48	4864		0.41 0.69	52	65	67	81	65	76	0.61	0.85	0.42	0.77
Chuck	54	5468	0.51	0.70	44	60	66	85	63	83	0.67	0.86	0.42	0.68
Foreshank	10	1013		0.06 0.14	08	15	44	40	48	51	0.37	0.51	0.23	0.31
Brisket	48	67		0.44 0.65	18	44	48	67	41	76	0.55	0.79	0.42	0.59
Plate	33	43	0.42 0.41	0.41	10	27	10	34	15	67	0.35	0.44	0.13	0.19
Flank	71	65	0.83 0.61	0.61	47	35	51	73	45	84	0.85	0.80	0.69	09.0
Kidney knob7842	78	45		0.82 0.38	65	25	60	42	52	45	0.86 0.20	0.20	0.15	0.12

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01). RLRC = Round, loin, rib and chuck.

carcass muscle and bone were negatively related to pounds and percents wholesale cut fat trim within weight groups (range, -.07 to -.78); whereas, in the combined weight groups the relationships between pounds of these characteristics were positive but low (range, 0.08 to 0.55). Percent wholesale cut fat trim was highly significantly (P < .01) negatively related to percent carcass muscle and bone both in the combined and within weight groups except those between percent fat trim from the foreshank and separable muscle and bone and percent plate, flank and kidney fat and total carcass bone in the 700 to 750 lb. weight group.

Among the wholesale cuts, pounds of fat trim from the kidney knob and flank showed the highest relationships to pounds of total separable carcass fat for the combined and within weight groups (range, 0.70 to 0.91). When compared on a percent basis, correlations for kidney knob and flank with total carcass separable fat were lower, with the greatest reduction in the 700 to 750 lb. weight group. The relationships between percent fat trim of the other wholesale cuts and total carcass separable fat were highly significant (P < .01) with the exception of the percent foreshank fat in the 700 to 750 lb. weight group (0.14). These data indicate that foreshank fat deposition is not consistent with total carcass fat deposition.

Highly significant (P < .01) correlation coefficients were found between percent wholesale cut fat trim yields and percent boned, trimmed round, loin, rib and chuck retail yield and total retail yield in the combined and within weight groups, except for plate fat trim in the 700 to 750 lb. weight group (range, -.40 to -.93). The lowest relationships

for all wholesale cuts with retail yields were noted in the 700 to 750 lb. weight group (range, -10 to -.85). In the 500 to 550 lb. weight group, percent fat trim from all wholesale cuts except the kidney knob and foreshank showed correlations with retail yields of -.76 or higher. Percent flank fat trim was also highly related to retail yields (range, -.73 to -.89) and would again suggest use of the flank for predictive purposes.

Percents wholesale cut fat trim, except that from the kidney knob, were positively and highly significantly (P < .01) related to percent total fat trim yields for the combined weight groups (range, 0.54 to 0.93). With the exception of the kidney knob, foreshank and plate, correlations between percent fat trim yields and percent carcass fat trim yields were highly significant (P < .01) within the two weight groups (range, 0.59 to 0.96). Correlations between pounds and percent wholesale cut fat trim yields and pounds and percent carcass separable components, retail and fat trim yields were generally higher in the 500 to 550 lb. weight group than in the 700 to 750 lb. weight group.

Relationships Between Wholesale Cut Bone Yield and Total Carcass Separable

Components, Retail and Fat Trim Yields

Simple correlation coefficients between pounds and percent wholesale cut bone yields and pounds and percent carcass separable components, retail and fat trim yields for the combined and within weight groups appear in tables 69, 70 and 71. These relationships were similar to those previously discussed between separable bone and the seven carcass compositional components. Both pounds and percents wholesale cut bone yields were generally

Table 69. Simple correlation coefficients between wholesale cut bone yield and carcass separable components retail and fat trim yields for the combined weight groups.	Simple retail	corre and f	lation at trim	Simple correlation coefficeretail and fat trim yields	ients for t	betweer he comb	ents between wholesale cut bonders the combined weight groups.	ale cut [ght gro	bone yi	eld and	carcas	s sepa1	rable co	mponents,
	Total	a1	Total	al	1	al	Boned,	, pg	Boned	ed,				
Wholesale	separable	able	separable	able	separable	able	trimmed	ned	trimmed	ned	Total fat	fat	Exter	External fat
cut pone	muscle	:le	fat	íŤ	bone	as	retail RLRC	RLRC1	retail yield	yield	trim	g	trim from	rom RLRC
yield	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%
Round	0.84	0.84 0.38	0.32	48	0.92	0.65	0.83	0.30	0.82	0.31	0.22	48	0.09	48
Loin	0.74	0.74 0.35	0.37	40	0.80	0.51	0.74	0.29	0.74	0.32	0.23	46	0.14	40
Rib	0.76	0.76 0.52	0.31	56	0.73	09.0	0.70	0.38	69.0	0.40	0.24	52	0.14	44
Chuck	0.90	0.90 0.59	0.31	99	0.92	0.76	0.84	0.44	0.83	0.48	0.24	62	0.08	57
Foreshank	0.80	0.80 0.04	0.32	03	0.88	0.10	08.0	0.07	0.81	04	0.20	06	90.0	15
Brisket	0.48	0.48 0.63	07	71	0.62	0.79	0.44	0.57	97.0	0.58	10	66	16	59
Plate	0.79	0.79 0.67	0.24	72	0.83	0.77	0.78	0.61	0.73	0.58	0.12	66	0.00	59
Flank	0.10	0.10 0.14	0.13	22	0.21	0.39	0.21	0.28	0.25	0.36	03	38	01	29

Correlations > 0.220 are significant (P < .05). Correlations > 0.287 are significant (P < .01). 1 KLRC = Round, loin, rib and chuck.

Table 70.	Simple retail	correl and fa	Simple correlation coeffic: retail and fat trim yields	oeffici yields	lents betweer for the 500	etween e 500 t	wholesa to 550 l	le cut l b. weigh	n wholesale cut bone yiel to 550 lb, weight group.	ld and .	carcass	separ	Simple correlation coefficients between wholesale cut bone yield and carcass separable components retail and fat trim yields for the 500 to 550 lb, weight group.	onents,
	Total	a1	Total	al	Total	a1	Boned	ed,	Boned	ed,				
Wholesale	sepai	separable	separable	able	separable	able	trimmed	med	trimmed	ped	Total fat	fat	External	al fat
cut pone	muscle	:le	fa	fat	bone	as	retail	$retail$ $RLRC^1$	retail yield	yield	trim	g	trim fr	trim from RLRC
yteld	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%	1b.	%
Round	0.45	0.45 0.34	70	45	0.76	0.76 0.59	0.47	0.37	0.51	0.43	71	55	69	54
Loin	0.02	0.02 0.21	21	25	0.30	0.38	0.16	0.30	0.26	0.38	45	47	37	-, 38
Rib	0.26	0.26 0.43	20	48	0.23	0.55	0.20	0.41	0.22	0.47	31	56	22	-,45
Chuck	0.58	0.58 0.54	66	62	0.72	0.72	0.51	0.52	0.50	0.57	-, 65	67	<b>-</b> . 60	60
Foreshank	0.39	0.3904	60	60 0.09	0.68	12	0.42	0.02	0.53	0.01	70	06	65	10
Brisket	0.26	0.26 0.64	57	73	0.56	0.80	0.23	09.0	0.31	0.63	48	67	40	59
Plate	0.31	0.31 0.58	43	66	0.58	0.77	0.41	09.0	97.0	0.59	57	65	58	59
Flank	11	11 0.14		0423	0.23	97.0	90.0	0.21	0.19	0.32	21	-,35	17	29

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01).

IRLRC = Round, loin, rib and chuck.

Table 71. Simple correlation coefficients between wholesale cut bone yield and carcass separable components retail and fat trim vields for the 700 to 750 lb. weight prome	Simple retail	correl	Simple correlation coefficeretail and fat trim vields	oeffici	lents by	etween 700 t	ents between wholesale cut bone yiel for the 700 to 750 lb, weight group.	le cut l	one yie	ld and c	arcass	separa	ble comp	onents,
	Total	al:	Total	al	Total	al	Boned	ad,	Boned	ed,				
Wholesale	sepaı	separable	separable	able	separable	able	trimmed	ned	trimmed	med	Total fat	fat	External fat	al fat
cut bone yield	muscle 1b.	%	fat 1b.	it %	bone 1b.	% %	retail KLRC* 1b. %	KLRC*	retail yield 1b. %	yield %	trim 1b.	# %	trim from RLRC 1b. %	om RLRC %
Round	0.58	0.58 0.46	58	57	0.83 0.78	0.78	0.48	0.32	0.41	0.22	56	50	50	54
Loin	0.62	0.62 0.66	53	73	0.77	08.0	0.50	0.44	0.42	0.41	53	64	42	-, 63
Rib	0.63	0.63 0.71	68	74	0.61	0.72	0.32	0.39	0.26	0.32	55	54	49	50
Chuck	0.80	0.80 0.68	67	76	0.86	0.84	0.56	0.36	0.52	0.32	60	57	63	59
Foreshank	0.52	0.52 0.27	48	36	0.76	0.54	97.0	0.30	0.44	0.18	50	35	50	41
Brisket	0.56	0.56 0.62	44	68	0.68	0.76	0.43	0.44	0.47	0.42	42	59	42	58
Plate	0.71	0.71 0.79	57	82	0.73	0.78	0.69	0.67	99.0	0.64	62	75	53	62
Flank	0.00	0.00 0.10	0.19	12	0.07	0.16	0.41	0.37	0.44	0.37	09	-,38	0.04	19

Correlations > 0.312 are significant (P < .05). Correlations > 0.403 are significant (P < .01).

IRLRC = Round, loin, rib and chuck.

positively related to measures of carcass muscling and bone and negatively related to carcass fatness.

As would be expected, in the combined weight groups the highest relationships were noted between pounds and percent wholesale cut bone yields (except pounds and percent flank bone yield and percent foreshank bone yield) and pounds and percent total carcass bone (range, 0.51 to 0.92). In the combined weight groups, the highest relationships were observed between pounds of bone from the round (0.92), chuck (0.92) and foreshank (0.88) and pounds of separable carcass bone. On a percent basis, these same relationships were markedly reduced (range, 0.10 to 0.76).

Multiple Regression Analyses of Pounds and Percent Retail Yield Measures
on a Combination of Wholesale Cut Yields and

Carcass Scores and Measurements

Multiple Regression Analyses of Pounds and Percent Boned, Trimmed Retail Yield from the Round, Loin, Rib and Chuck on a Combination of Wholesale Cut Yields and Objective Carcass Measurements. In the combined weight groups, 95% of the variation in pounds of boned, trimmed retail yield from the round, loin, rib and chuck (RLRC) could be accounted for by the six independent variables listed in table 72. The prediction of weight of retail yield from the RLRC using this equation resulted in a standard error of estimate of 5.28 lb. In the 500 to 550 lb. weight group, 85% of the variation in retail yield from the RLRC was accounted for by seven independent variables (suppl., p. 182). In the 700 to 750 lb. weight group, only five variables had a significant effect on weight of boned, trimmed RLRC yield and they were associated with 76% of the variation present in this variable (suppl., p. 182).

Table 72. Coefficient of determination between weight of boned, trimmed round, loin, rib and chuck retail yield and a combination of weights of wholesale cut yields and objective carcass measurements, and the regression coefficient for each of the variables (combined weight groups).

	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	-75.841		0.00	
Carcass wt.	0.217	0.906	0.00	0.84
Round score <sup>2</sup>	1.535	0.104	0.00	0.94
Length of round	4.028	0.347	0.00	0.93
Wt. foreshank bone-R.S.3	-4.945	157	0.02	0.94
Wt. flank yield-R.S.	2.198	0.201	0.00	0.92
Wt. flank fat-R.S.	-2.197	407	0.00	0.89

 $R^2 = 0.95$  Standard error of estimate = 5.28

When the wholesale cut yields were expressed in percentages, 94% of the variation in pounds of boned, trimmed retail RLRC yield was accounted for combined weight groups using six independent variables (suppl., p. 183). In the combined weight groups, 79% of the variation in percent retail yield of the RLRC was accounted for by the six variables listed in table 73.

In the 500 to 550 lb. weight group, 79% of the variation in weight of boned, trimmed RLRC yield could be accounted for using carcass weight, length of round, 1. dorsi muscle area and percent retail yield of the flank (suppl., p. 183). Eighty-three percent of the variation in percent

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>2</sup>See table 2 for scoring system used.

 $<sup>^{3}</sup>$ R.S. = Right side.

Table 73. Coefficient of determination between percent boned, trimmed round, loin, rib and chuck retail yield and a combination of percent wholesale cut yields and carcass scores and measurements, and the regression coefficient for each of the variables (combined weight groups)

(combined weigh	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	10.953		0.09	
Carcass wt.	023	600	0.00	0.72
Round score <sup>2</sup>	0.417	0.179	0.02	0.77
Length of round	1.439	0.786	0.00	0.71
Wt. foreshank bone-R.S. <sup>3</sup>	-1.670	336	0.02	0.77
L. dorsi muscle area	0.394	<b>0.1</b> 85	0.01	0.77
% flank yield-R.S.	0.301	0.696	0.00	0.52

 $R^2 = 0.79$ Standard error of estimate = 1.70

retail yield of the RLRC could be accounted for using these same four variables plus weight of the foreshank bone (suppl., p. 183).

In the 700 to 750 lb. weight group, only 76% of the variation in pounds of retail yield of the RLRC (suppl., p. 184) and 67% of the variation in percent of the same trait (suppl., p. 184) could be accounted for using carcass weight, round score, length of round and percent retail yield of the flank.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

2 See table 2 for scoring system used.

 $<sup>^{3}</sup>$ R.S. = Right side.

Multiple Regression Analyses of Pounds and Percent Boned, Trimmed Retail Round, Loin, Rib and Chuck Yield on Objective Carcass Measurements and Percent Flank Retail Yield. In the combined weight groups, 94% of the variation in pounds of RLRC yield and 76% of the variation in percent RLRC yield can be accounted for using the variables listed in the equations shown in table 74. In the 500 to 550 lb. weight group (table 75), somewhat greater accuracy in predicting percent of RLRC yield was attained. The coefficients of determination for equations 1 (0.78) and 2 (0.81) in the 500 to 550 lb. weight group were lower compared to equations 1 (0.94) and 2 (0.94) in the combined weight groups. The coefficients of determination for equations 3 (0.78 and 4 (0.81) in the 500 to 550 lb. weight group were greater than the corresponding equations for the combined weight groups (0.76 and 0.76, respectively).

In the 700 to 750 lb. weight group (table 76), less of the total variation in either pounds or percent RLRC yield could be accounted for using these multiple regression analyses than in either the combined or the 500 to 550 lb. weight groups (coefficient of determination range 0.67 to 0.75). These data indicate that as beef carcasses get heavier (fatter), more difficulty in accurately estimating pounds or percent of RLRC yield is encountered.

These regression analyses were slightly less accurate with somewhat larger standard errors of estimate than those reported by Murphy et al. (1960) and Brungardt and Bray (1963).

chuck retail yield using objective carcass measurements and percent flank retail yield as predictors, Multiple regression equations for estimating pounds and percent boned, trimmed round, loin, rib and and the coefficient of determination for each equation (combined weight groups) Table 74.

Equation	τ.	Coefficient of	Standard error of
No.	Estimating equations	determination	estimate
1	$ \frac{4}{3} = -44.501 + 0.204(x_1) + 0.024(x_2)314(x_3) + 1.610(x_5) + 0.797(x_6) $	0.94	5.86
2	$\hat{Y} = -41.795 + 0.206(x_1)307(x_3)034(x_4) + 1.510(x_5) + 0.786(x_6)$	0.94	5.86
ო	$\hat{\Lambda}_1$ = 40.969014(x <sub>1</sub> )049(x <sub>2</sub> )128(x <sub>3</sub> ) + 0.407(x <sub>5</sub> ) + 0.236(x <sub>6</sub> )	0.76	1.80
4	$\hat{Y}_1$ = 40.614014(X <sub>1</sub> )129(X <sub>3</sub> )031(X <sub>4</sub> ) + 0.429(X <sub>5</sub> ) + 0.235(X <sub>6</sub> )	0.76	1.79
<\r>□	$\Lambda$ = Estimated pounds of boned, trimmed retail round, loin, rib and chuck yield.		
√1 √1	$\hat{Y}_1$ = Estimated percent boned, trimmed retail round, loin, rib and chuck yield.		

 $X_2$  = Probe - 5th thoracic vertebra, 4 in. off the carcass dorsal midline (mm.).

 $X_1$  = Carcass weight (1b.).

 $X_3$  = Probe - 3rd sacral vertebra, 12 in. off the carcass dorsal midline (mm.).

 $X_4$  = Fat measurement A (mm.).

 $X_5$  = Length of round (in.).

chuck retail yield using objective carcass measurements and percent flank retail yield as predictors, Multiple regression equations for estimating pounds and percent boned, trimmed round, loin, rib and and the coefficient of determination for each equation (500 to 550 lb, weight group). Table 75.

Equation No.	Estimating equation	Coefficient of determination	Standard error of estimate
1	$^{\wedge}_{Y}$ = -100.107 + 0.295(x <sub>1</sub> )200(x <sub>2</sub> )418(x <sub>3</sub> ) + 2.391(x <sub>5</sub> ) + 0.543(x <sub>6</sub> )	0.78	5.14
2	$\hat{A} = -61.465 + 0.257(x_1)437(x_3)284(x_4) + 1.935(x_5) + 0.493(x_6)$	0.81	4.84
٣	$A_1 = 18.648 + 0.019(x_1)112(x_2)153(x_3) + 0.697(x_5) + 0.190(x_6)$	0.78	2.01
4	$\sqrt{4} = 32.241 + 0.003(x_1)162(x_3)110(x_4) + 0.566(x_5) + 0.177(x_6)$	0.81	1.91
$\hat{\mathbf{Y}} = \mathbf{E}\mathbf{s}$	$\hat{\mathbf{Y}} = \mathrm{Estimated}$ pounds of boned, trimmed retail round, loin, rib and chuck yield.		

 $\hat{Y}_1$  = Estimated percent boned, trimmed retail round, loin, rib and chuck yield.

 $X_1$  = Carcass weight (1b.).

 $\rm X_2$  = Probe - 5th thoracic vertebra, 4 in. off the carcass dorsal midline (mm.).

 $X_3$  = Probe - 3rd sacral vertebra, 12 in. off the carcass dorsal midline (mm.).

 $X_4$  = Fat measurement A (mm.).

 $X_5$  = Length of round (in.).

Table 76. Multiple regression equations for estimating pounds and/or percent boned, trimmed round, loin, rib	and chuck retail yield using objective carcass measurements and percent flank retail yield as pre-	dictors, and the coefficient of determination for each equation (700 to 750 1b. weight group).	Standard
Table 76.			

Equation No.	Estimating equations	Coefficient of determination	error of
-	$A = -71.018 + 0.211(x_1) + 0.038(x_2)326(x_3) + 1.714(x_5) + 1.268(x_6)$	0.74	5.94
7	$\hat{A} = -59.662 + 0.183(x_1)422(x_3) + 0.168(x_4) + 1.846(x_5) + 1.333(x_6)$	0.75	5.74
က	$\hat{Y}_1 = 41.033016(x_1)034(x_2)124(x_3) + 0.378(x_5) + 0.299(x_6)$	0.67	1.59
4	$\hat{Y}_1 = 40.567020(x_1)146(x_3) + 0.021(x_4) + 0.450(x_5) + 0.309(x_6)$	0.68	1.58

 $\hat{Y}$  = Estimated pounds of boned, trimmed retail round, loin, rib and chuck yield.

 $\hat{Y}_1$  = Estimated percent boned, trimmed retail round, loin, rib and chuck yield.

 $X_1$  = Carcass weight (1b.).

 $X_2$  = Probe - 5th thoracic vertebra, 4 in. off the carcass dorsal midline (mm.).

 $X_3$  = Probe - 3rd sacral vertebra, 12 in. off the carcass dorsal midline (mm.).

 $X_4$  = Fat measurement A (mm.).

 $X_5$  = Length of round (in.).

Multiple Regression Analyses of Pounds and Percent Boned, Trimmed Total

Carcass Retail Yield on a Combination of Weights of Wholesale Cut Yields

and Carcass Scores and Measurements. In the combined weight groups, 96%

of the variation in pounds of total carcass boned, trimmed retail yield

was accounted for using six independent variables (table 77).

Table 77. Coefficient of determination between pounds of boned, trimmed total carcass retail yield and a combination of weights of wholesale cut yields and objective carcass measurements, and the regression coefficient for each of the variables (combined weight groups).

weight groups).				
T-11	Regression	Beta	Level of	R <sup>21</sup>
Independent variables	coefficients	weights	sig.	deletes
Constant	-25.066		0.08	
Carcass wt.	0.189	0.651	0.00	0.93
Depth of 10th rib	1.992	0.061	0.02	0.96
Length of round	1.493	0.106	0.03	0.96
Wt. untrimmed round-R.S. <sup>2</sup>	0.542	0.204	0.01	<b>0.</b> 96
Wt. retail flank yield-R.S.	3.692	<b>0.27</b> 8	0.00	0.91
Wt. retail flank fat-R.S.	-2.089	319	0.00	0.93

 $R^2 = 0.96$  Standard error of estimate = 5.70

Estimating percent total carcass boned, trimmed retail yield in the combined weight groups, 84% of the variation could be accounted for using the five independent variables shown in table 78.

Greater accuracy was found in the 500 to 550 lb. weight group as 87% of the variation in pounds of boned, trimmed retail yield was related

<sup>1</sup> Value of coefficient of determination if that variable was deleted from the analysis.

 $<sup>^{2}</sup>$ R.S. = Right side.

Table 78. Coefficient of determination between percent boned, trimmed total carcass retail yield and a combination of weights of wholesale cut yields and carcass scores and measurements, and the regression coefficient for each of the variables (combined weight groups).

weight groups).	Regression	Beta	Level of	
Independent variables	coefficients	weights	sig.	deletes_
Constant	41.717		0.00	
Carcass wt.	024	494	0.00	0.80
Round score <sup>2</sup>	0.602	0.201	0.00	0.81
Length of round	1.118	0.474	0.00	0.79
Wt. retail flank yield-R.S.3	1.116	0.502	0.00	0.67
Wt. retail flank fat-R.S.	788	718	0.00	0.64

 $R^2 = 0.84$  Standard error of estimate = 1.91

to the five independent variables shown in table 79. This prediction equation, involving five variables, would be useful for the prediction of pounds of boned, trimmed retail yield in 500 to 550 lb. carcasses. These same five variables accounted for 90% of the variation in percent boned, trimmed carcass retail yield in the 500 to 550 lb. weight group (table 80).

In the 700 to 750 lb. weight group, 80% of the variation in weight of boned, trimmed carcass retail yield could be accounted by carcass weight and weights of retail flank muscle and fat (suppl., p. 185). Seventy-six percent of the variation in percent of this same variable could be accounted for using these same three independent variables plus the weight of the kidney knob (suppl., p. 185).

<sup>1</sup> Value of coefficient of determination if that variable was deleted from the analysis.

<sup>2</sup>See table 2 for scoring system used.

 $<sup>^{3}</sup>$ R.S. = Right side

Table 79. Coefficient of determination between weight of boned, trimmed total carcass retail yield and a combination of weights of wholesale cut yields and objective carcass measurements, and the regression coefficient for each of the variables (500 to

550 lb. weight group).

Independent variables	Regression coefficients	Beta weights	Level of sig.	R21 deletes
Constant	-130.465		0.01	
Carcass wt.	0.312	0.376	0.00	0.77
Length of round	3.905	0.344	0.00	0.80
Wt. retail flank yield-R.S. <sup>2</sup>	3.557	0.446	0.00	0.70
Wt. retail flank fat-R.S.	-2.142	518	0.00	0.72
L. dorsi muscle area	2.085	0.203	0.01	0.83

 $R^2 = 0.87$  Standard error of estimate = 5.38

Table 80. Coefficient of determination between percent boned, trimmed total carcass retail yield and a combination of weights of wholesale cut yields and objective carcass measurements, and the regression coefficient for each of the variables (500 to 550 lb. weight group).

Independent variables	Regression coefficients	Beta weights	Level of sig.	R21
		Weights		deretes
Constant	15.513		0.33	
Carcass wt.	0.005	0.014	0.81	0.90
Length of round	1.327	0.299	0.00	0.85
Wt. retail flank yield-R.S. <sup>2</sup>	1.230	0.394	0.00	0.77
Wt. retail flank fat-R.S.	808	500	0.00	0.76
L. dorsi muscle area	0.907	0.226	0.00	0.86

 $R^2 = 0.90$  Standard error of estimate = 1.83

<sup>1</sup> Value of coefficient of determination if that variable was deleted from the analysis.

 $<sup>^{2}</sup>$ R.S. = Right side.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

 $<sup>^{2}</sup>$ R.S. = Right side.

The preceding equations compare favorably with prediction equations of Murphy et al. (1960), Breidenstein (1962) and Brungardt and Bray (1963) for accuracy of prediction. The equations of Murphy et al. (1960) and Brungardt and Bray (1963) predict only the boned, trimmed retail yield of the RLRC; whereas, those presented here also estimate total carcass boned, trimmed retail yield. Greater accuracy was noted in the 500 to 550 lb. weight group than either the 700 to 750 lb. weight group or the combined weight groups.

Multiple Regression Analyses of Pounds and Percent Boned, Trimmed Total Carcass Retail Yield on a Combination of Wholesale Cut Yield Percentages and Carcass Scores and Measurements. When percent wholesale cut retail and fat trim yields were used in the prediction equations, the results were similar to those for wholesale cut retail weights in the combined weight groups. Ninety-five percent of the variation in pounds of total carcass boned, trimmed retail yield could be accounted for (combined weight groups) using the following four variables: 1) carcass weight; 2) length of round; 3) 1. dorsi muscle area; and 4) % retail flank yield (suppl., p. 186).

The same variables plus round score, accounted for 85% of the variation in percent boned, trimmed carcass retail yield in the combined weight groups (suppl., p. 186).

In the 500 to 550 lb. weight group, 84 and 88% of the variation could be accounted for in pounds and percent, respectively, of boned, trimmed retail carcass yield (suppl., p. 187). In the 700 to 750 lb. weight group,

79 and 74% of the variation in the same two variables could be accounted for using the regression equations presented in the suppl., p. 188.

These equations, particularly on a combined weight group basis and in the 500 to 550 lb. weight group, compare favorably in accuracy with the prediction equations of Murphy et al. (1960), Breidenstein (1962) and Brungardt and Bray (1963).

Multiple Regression Analyses of Boned, Trimmed Total Carcass Retail Yield on Objective Carcass Measurements and Percent Flank Retail Yield. Ninety-four percent of the variation in pounds of boned, trimmed carcass retail yield (combined weight groups) could be accounted for using either equation 1 or 2 shown in table 81. Eighty-three and 84% of the variation in percent boned, trimmed carcass retail yield could be accounted for using equations 3 and 4 (table 81), respectively.

Greater accuracy of prediction of percent boned, trimmed carcass retail yield was accomplished in the 500 to 550 lb. weight group (table 82) using either equation 3 or 4. In the 700 to 750 lb. weight group, however, only 74% of the variation in percent boned, trimmed carcass retail yield could be accounted for using the corresponding equations (equations 3 or 4, table 83).

These data indicate that total carcass boned, trimmed retail yield could be quite accurately estimated using a combination of objective carcass measurements and percent flank retail yield. However, it would appear that greater accuracy of prediction can be obtained in lighter weight carcasses than those of heavier weights.

Multiple regression equations for estimating pounds and percent boned, trimmed total carcass metail yield using objective carcass measurements and percent flank retail yield as predictors, and the coefficient of determination for each equation (combined weight groups). Table 81.

			Standard
Equation		Coefficient of	error of
No.	Estimating equations	determination	estimate
F.	$^{\wedge}_{Y}$ = -67.173 + 0.255( $x_1$ ) + 0.144( $x_2$ )208( $x_3$ ) + 1.707( $x_5$ ) + 1.283( $x_6$ )	0.94	6.59
2	$\sqrt{Y} = -60.751 + 0.260(x_1)193(x_3)016(x_4) + 1.454(x_5) + 1.261(x_6)$	0.94	6.60
m	$\hat{Y}_1 = 44.627015(x_1)019(x_2)102(x_3) + 0.460(x_5) + 0.392(x_6)$	0.83	1.92
4	$\hat{Y}_1 = 45.402014(x_1)100(x_3)030(x_4) + 0.436(x_5) + 0.388(x_6)$	0.84	1.91
<			

 $\dot{Y}$  = Estimated pounds of boned, trimmed total carcass retail yield.

 $\vec{Y}_1$  = Estimated percent boned, trimmed total carcass retail yield.

 $X_1$  = Carcass weight (1b.).

 $\rm X_2$  = Probe - 5th thoracic vertebra, 4 in. off the carcass dorsal midline (mm.).

 $X_3$  = Probe - 3rd sacral vertebra, 12 in. off the carcass dorsal midline (mm.).

 $X_4$  = Fat measurement A (mm.).

 $X_5$  = Length of round (in.).

Multiple regression equations for estimating pounds and percent boned, trimmed total carcass retail yield using objective carcass measurements and percent retail flank yield as predictors, and the coefficient of determination for each equation (500 to 550 lb. weight group) Table 82.

Equation No.	Estimating equation	Coefficient of determination	standard error of estimate
1	$^{\wedge}_{Y}$ = -153.246 + 0.357( $x_1$ ) + 0.044( $x_2$ )438( $x_3$ ) + 3.547( $x_5$ ) + 0.933( $x_6$ )	0.84	5.79
7	$\hat{Y} = -92.587 + 0.317(x_1)452(x_3)361(x_4) + 2.581(x_5) + 0.810(x_6)$	0.87	5.32
ന	$\hat{Y}_1 = 10.981 + 0.022(X_1)026(X_2)158(X_3) + 1.085(X_5) + 0.336(X_6)$	0.87	2.10
7	$\hat{Y}_1 = 32.862 + 0.005(x_1)165(x_3)140(x_4) + 0.766(x_5) + 0.297(x_6)$	0.89	1.90
•			

 $\stackrel{ extstyle op}{ extstyle op}$  = Estimated pounds of boned, trimmed total carcass retail yield.

 $\hat{\mathrm{Y}}_1$  = Estimated percent boned, trimmed total carcass retail yield.

 $X_1$  = Carcass weight (1b.).

 $X_2$  = Probe - 5th thoracic vertebra, 4 in. off the carcass dorsal midline (mm.).

 $X_3$  = Probe - 3rd sacral vertebra, 12 in. off the carcass dorsal midline (mm.).

 $X_4$  = Fat measurement A (mm.).

 $X_5$  = Length of round (in.).

Multiple regression equations for estimating pounds and percent boned, trimmed total carcass retail yield using objective carcass measurements and percent flank retail yield as predictors, and coefficient of determination for each equation (700 to 750 lb. weight group). Table 83.

	· /dary	.75.	
7 4 4 5 5 T		Coefficiant of	Standard
No.	Estimating equation	determination	estimate
Н	$\hat{\mathbf{Y}} = -104.379 + 0.300(x_1)069(x_2)019(x_3) + 1.229(x_5) + 1.820(x_6)$	08.0	6.33
2	$\hat{Y} = -96.877 + 0.266(x_1)154(x_3) + 0.190(x_4) + 1.540(x_5) + 1.899(x_6)$	0.82	60.9
က	$\hat{Y}_1 = 44.587008(x_1)075(x_2)043(x_3) + 0.220(x_5) + 0.443(x_6)$	0.74	1,65
4	$\hat{Y}_1 = 42.242012(x_1)074(x_3) + 0.022(x_4) + 0.352(x_5) + 0.455(x_6)$	0.74	1.65
<b>♦</b>	V = Detimoted sounds of board trimmed total serves retail wield		

Y = Estimated pounds of boned, trimmed total carcass retail yield.

 $oldsymbol{\hat{Y}}_1$  = Estimated percent boned, trimmed total carcass retail yield.

 $X_1$  = Carcass weight (1b.).

 $X_2$  = Probe - 5th thoracic vertebra, 4 in. off the carcass dorsal midline (mm.).

 $X_3$  = Probe - 3rd sacral vertebra, 12 in. off the carcass dorsal midline (mm.).

 $X_4$  = Fat measurement A (mm.).

 $X_5$  = Length of round (in.).

Relationships Between <u>L</u>. <u>dorsi</u> Muscle Potassium and Sodium and <u>L</u>. <u>dorsi</u>

Weight, Area and Total Carcass Separable Muscle

Simple correlation coefficients, were calculated between 1. dorsi muscle potassium and sodium and 1.dorsi muscle weight, area and total carcass separable muscle on a fresh basis (total ppm of chemically determined element present in the muscle sample) and on a protein basis (ppm of the element/percent protein in the sample). These correlations appear in table 84.

Table 84. Simple correlation coefficients between muscle potassium and sodium and 1. dorsi muscle weight, area, and total carcass separable muscle.

	Potassium		Sodium	
	Fresh basis	Protein basis <sup>1</sup>	Fresh basis	Protein basis <sup>1</sup>
Combined weight groups				
L. dorsi muscle wt.	0.16	0.09	0.24	0.20
L. dorsi muscle area Total separable muscle	0.20	0.19	0.08	0.07
of carcasses	0.17	0.01	0.25	0.16
500 to 550 lb. weight group				
L. dorsi muscle wt.	0.04	0.30	20	02
L. dorsi muscle area Total separable muscle	0.11	0.24	24	14
of carcasses	0.18	0.06	29	32
700 to 750 lb. weight group				
L. dorsi muscle wt.	0.14	0.06	09	12
L. dorsi muscle area Total separable muscle	0/19	0.26	08	06
of carcasses	0.11	0.08	0.00	02

Correlations > 0.220 are significant (p < .05) for the combined weight groups.

Correlations > 0.287 are significant (P < .01) for the combined weight groups.

Correlations > 0.312 are significant (P < .05) for the individual weight groups.

Correlations > 0.403 are significant (P < .01) for the individual weight groups.

Expressed as ppm/percent protein.

It is apparent upon examining these data that <u>1</u>. <u>dorsi</u> muscle potassium and sodium content had little relationship to <u>1</u>. <u>dorsi</u> muscle weight, area or total carcass separable muscle. Correlations reported here showed little if any predictive value for total carcass separable muscle. This is in contrast to the results of Kirton and Pearson (1963a) and (1963b) who found correlations, between potassium content of a sample of entire lamb carcasses and percent protein of the sample, which were high enough to be of limited predictive use. It should be noted however that these workers determined potassium content of a tissue sample of the entire carcass whereas, in this study a sample of the <u>1</u>. <u>dorsi</u> muscle was studied. Gillett <u>et al</u>. (1965) found considerable variation in potassium content between various entire porcine muscles.

Means and Standard Deviations of Some Subjective Carcass Scores and Some
Chemical and Organoleptic Characteristics

The means and standard deviations for the subjective carcass scores, chemical and organoleptic characteristics discussed in this section for the combined and individual weight groups and fat thickness ranges within weight groups are presented in table 85.

On a within weight and fat thickness group basis, the means for carcass grade and marbling score show a positive relationship to 12th rib fat thickness and total carcass fat content. As fat thickness (12th rib) increased, there was a marked increase in carcass grade and marbling score particularly in groups I through IV (5.20 to 8.00 for carcass grade and 11.70 to 18.30 for marbling score). Maturity score means show a negative relationship to measures of carcass fatness.

Means and standard deviations of several subjective carcass scores and some chemical and organoleptic characteristics for the combined and individual weight groups and fat thickness ranges within weight groups. Table 85.

groups.	S.											
		Combined	500- 550 1b.	700 <b>-</b> 750 1b.				Group				
Trait		weight group	weight group	weight group	I 0.26"- 0.50"	II 0.51"- 0.75"	111 0.76"- 1.00"	IV 1.01"- 1.25"	V 0.26"- 0.50"	VI 0.51"- 0.75"	VII 0.76"- 1.00"	VIII 1.01"- 1.25"
Carcass grade	Mean S.D.	7.20	7.02	7.38	5.20	7.00	7.90	8.00 1.05	6.90 1.20	7.40	7.60	7.60
Marbling score	Mean S.D.	16.99 3.82	16.05 3.88	17.92 3.56	11.70	16.70 3.37	17.50 2.84	18.30 3.20	17.20 2.94	17.80	18.30 4.78	18.40 3.84
Maturity	Mean S.D.	3.42 1.08	2.82 0.71	4.02	3.10	2.70 0.82	2.70	2.80	4.70 1.64	3.90	3.50	4.00
% ether extract of L. D. $\frac{2}{3}$	Mean S.D.	8.56 3.50	8.34 3.89	8.79	4.09	7.85	11.13 3.78	10.28 3.00	7.60	7.76	10.82 3.46	2.58
% protein of L. D.	Mean S.D.	21.18 0.91	21.04 0.90	21.33 0.92	21.56	21.21 0.30	20.64	20.73 0.92	21.39	21.24 1.06	21.13 1.03	21.55
% moisture of L. D.	Mean S.D.	69.50	69.74 3.14	69.26 2.66	73.24	69.94 2.34	67.55 3.18	68.26	70.47	70.48	67.27 2.72	68.84 2.26
Tenderness Warner-Bratzler	Mean S.D.	10.08 1.36	9.76	10.39 1.47	10.25 1.09	10.14 1.01	9.02	9.64 0.81	10.85 1.66	10.31 1.68	10.29	10.12 1.67
Tenderness Pane1	Mean S.D.	6.60	6.66 0.66	6.53 0.69	6.19	6.72	6.83	6.92	6.20	6.44	6.46	7.03
Juiciness Panel	Mean S.D.	6.48	6.52 0.57	6.45	6.38	6.77	6.38	6.56	6.57	6.23	6.45	6.53 0.25
Flavor Panel IS.D. = Standard	Mean 6 S.D. 0 deviation.	6.30 0.50 Ion.	6.29 0.48 <sup>Z</sup> L.D	6.31 0.52 = 1.	6.00 6 0.56 0 dorsi muscle	6.54 0.31 cle.	6.25 0.44 3nf	5 6.38 4 0.48 3Nine point	6.46 0.55 hedonic	5.98 0.46 scale.	6.31 0.54	6.47
				•				ı				

Means for percent protein, ether extract and moisture of the 1. dorsi muscle showed the expected relationships to fat thickness except for groups III and VII which had greater mean percents ether extract and percents protein with less moisture than groups IV and VIII.

Mean tenderness scores, both Warner-Bratzler and taste panel, indicated that as fat thickness, carcass grade and marbling score increased, tenderness increased. There were no marked relationships between panel flavor and juiciness scores and any of the characteristics studied. Within weight and fat thickness group correlations were determined for organoleptic data but are not presented in this dissertation.

## SUMMARY

Ten steer carcasses were selected within each of four fat thickness ranges (average of three measurements at the 12th rib) (0.26 to 0.50 in., 0.51 to 0.75 in., 0.76 to 1.00 in. and 1.01 to 1.25 in.) within each of two weight groups (500 to 550 lb. and 700 to 750 lb.). The carcasses were subjectively scored for each grade factor and some linear measurements of fat and muscle were recorded. External fat thickness probes were made 4, 8 and 12 in. off the dorsal midline of the left side of each carcass, perpendicular to the anterior edge of the 5th, 8th and 11th thoracic vertebrae, the 1st, 4th and 6th lumbar vertebrae and the 3rd and 5th sacral vertebrae.

The left side of each carcass was cut into wholesale cuts and physically separated into muscle, fat and bone with the exception of the round and rib. The rump was removed from the wholesale round and then the two portions individually physically separated. The wholesale rib was cut into the 6-7-8, 9-10-11 and 12th rib sections and each section was individually physically separated. The right side of each carcass was cut into boneless, closely trimmed (approximately 0.3 in.) retail cuts by wholesale cut.

Analysis of variance showed highly significant (P < .01) differences due to fat thickness for pounds and percent of the seven carcass compositional and yield characteristics studied. It also showed highly significant differences (P < .01) due to carcass weight for these same compositional and yield characteristics except percents carcass separable muscle, fat and bone. Highly significant (P < .01) interaction effects (inter-

action of carcass weight and fat thickness) were noted in percents round, loin, rib and chuck retail yield, total carcass retail yield, total fat trim and external fat trim from the round, loin, rib and chuck.

Subjective carcass scores showed little promise for estimation of carcass yield or composition. Despite a deliberate attempt to evaluate conformation independent of fat, subjective conformation scores were more highly correlated with measures of carcass fatness than muscling.

The fat probes were highly positively correlated with measures of carcass fatness and highly negatively related to measures of muscling and bone. The most promising fat thickness probe sites for predicting carcass composition, retail and fat trim yields were the 5th thoracic vertebra, 4 in. off the dorsal carcass midline and the average of the 3 probes at the 3rd sacral vertebra.

Regression equations for estimating carcass separable components, retail and fat trim yields, from the independent variables fat probes and carcass weight were more accurate for the 500 to 550 lb. weight group than the 700 to 750 lb. weight group. In general, pounds of the carcass separable components, retail or fat trim yields were more accurately predicted in the combined weight groups than percents.

Correlations between pounds and percents separable wholesale cut muscle and pounds and percent total carcass separable muscle, bone and retail yields were highly significant (P < .01) for the combined weight groups (range, 0.48 to 0.98) except for pounds and percent brisket muscle and percent foreshank muscle (range, 0.02 to 0.61). Pounds of separable chuck muscle showed the highest relationship to pounds of total carcass separable muscle for the combined weight groups (0.98) as well as within

weight groups (0.94 and 0.92 for the 500 to 550 lb. and 700 to 750 lb. weight groups, respectively).

Percent rib separable muscle showed the highest correlation with percent carcass separable muscle (0.95) followed by the loin (0.93), chuck (0.92) and the flank (0.91) (combined weight groups). All correlations between percent wholesale cut separable muscle and percent carcass separable muscle were highly significant (P < .01) for the combined weight groups except for that of the brisket.

In the combined weight groups, the percent rib separable fat was most highly related to percent total carcass separable fat (0.96) followed in order by the loin and chuck (0.94), plate (0.92) and the round and flank (0.91). These same cuts were also most highly related to carcass separable fat within weight groups although not in the same order.

Percent separable bone in the chuck had the highest correlation with percent carcass separable bone in the combined and 500 to 550 lb. weight groups (0.92 and 0.93, respectively). In the 700 to 750 lb. weight group, percent loin separable bone showed the highest relationship to percent carcass separable bone (0.90) followed by the chuck (0.89).

Percent 9-10-11 rib section and wholesale flank separable muscle and fat had the highest and most consistent correlations with percent total carcass separable muscle and fat for all weight and fat thickness groups (range, 0.71 to 0.95). Slightly lower correlations were observed between the percent wholesale round separable components and total carcass separable components for the individual and combined weight groups.

Ninety-four to 96% of the variation in pounds and 74 to 86% of the variation in percent of carcass separable muscle could be accounted for by

percents separable muscle from the 9-10-11 rib, flank or round, carcass weight and length of round in the combined weight groups. Regression equations including percent round separable muscle were less accurate than the 9-10-11 rib section or the flank for estimating total carcass muscle.

From 85 to 95% of the variation in pounds and percent carcass separable fat could be accounted for in the combined weight groups by carcass weight, 5th thoracic vertebra fat probe 4 in. off the midline and percent separable fat of the 9-10-11 rib section, flank or round.

Seventy-nine to 91% of the variation in percent total carcass separable bone could be accounted for in the combined weight groups by carcass weight, length of round and percent separable 9-10-11 rib section bone, round bone or separable flank muscle. Regression equations including flank muscle most accurately estimated either pounds or percent carcass separable bone followed in order by the percent 9-10-11 rib section bone and percent round bone.

These results indicate that the wholesale flank separable components could be utilized to accurately estimate carcass separable components.

This accuracy, the ease and small amount of time for separation of the flank and small economic loss of product compared to the rib, round or chuck strongly suggest the use of the wholesale flank separable components for estimation of total carcass composition.

Of the muscles studied, weight of the <u>1</u>. <u>dorsi</u> muscle had the highest relationship to pounds of total carcass muscle in the combined and 500 to 550 lb. weight groups (0.90 and 0.70, respectively). In the 700 to

750 lb. weight group, the highest correlation with pounds of carcass separable muscle was observed for the <u>semimembranosus</u> plus <u>adductor</u> muscles (0.71).

Highly significant correlation coefficients (P < .01) were found between weight of the radius plus ulna, femur, tibia plus fibula, scapula and humerus and total carcass bone weight for the combined and individual weight groups (range, 0.62 to 0.91).

Even though regression equations involving muscle and bone weights plus carcass weight to predict pounds or percent carcass separable muscle and bone were accurate, the laborious physically dissection techniques and economic loss of product limit their usefulness.

Correlations between percent muscle and fat were high (-.98) for the combined and individual weight groups. Correlations between percent carcass fat and bone were also highly significant (P < .01) (-.84) for the combined and individual weight groups. Positive correlations were observed between percent muscle and bone (0.74, 0.73 and 0.75 for the combined, 500 to 550 lb. and 700 to 750 lb. weight groups, respectively).

In most instances, percent retail and fat trim yields of the flank showed the highest relationships to total carcass retail and fat trim yields for the combined and individual weight groups (range, 0.57 to 0.89). These retail yield data also suggest the use of the wholesale flank for estimating round, loin, rib and chuck (RLRC) and total carcass retail yield.

Flank separable components plus combinations of objective measurements accounted for 95% of the variation in pounds of RLRC retail yield and 96% of the variation in carcass retail yield in the combined weight groups. Only 79 and 84% of the variation in percents of these retail yields could be accounted for in the combined weight groups by these same independent variables. Greater accuracy for estimating retail yields was observed in the 500 to 550 lb. weight group than in the 700 to 750 lb. weight group. Also, greater accuracy was obtained in the prediction of total carcass retail yield than for RLRC retail yield.

Low, non-significant correlation coefficients were found between 1.

dorsi muscle potassium and sodium and 1. dorsi muscle weight, area and total carcass separable muscle (range, 0.00 to 0.30). Means of the organoleptic analyses were not consistent with compositional data.

## LITERATURE CITED

- Allen, Deloran, M. 1963. Estimation of carcass quality and composition in the beef steer using live animal and carcass scores and measurements. M.S. Thesis, University of Idaho, Moscow, Idaho.
- Benne, E. J., N. H. Van Hall and A. M. Pearson. 1956. Analysis of fresh meat. J. Assn. Official Agr. Chem. 39:937.
- Black, W. H. 1938. A comparison of several methods of measuring performance in beef cattle. Proc. Am. Soc. An. Prod. p. 103.
- Black, W. H. and Bradford Knapp, Jr. 1936. A method of measuring performance in beef cattle. Proc. Am. Soc. An. Prod. p. 72.
- Branaman, G. A., A. M. Pearson, W. T. Magee, Ruth M. Griswold and G. A. Brown. 1962. Comparison of the cutability and eatability of beefand dairy-type cattle. J. Animal Sci. 21:321.
- Breidenstein, B. C. 1962. More definite system is needed to predict retail yield of beef carcass. The National Provisioner. 174, No. 14: 77.
- Briskey, E. J. and R. W. Bray. 1964. A special study of the beef grade standards for the American National Cattlemen's Association.
  University of Wisconsin, Madison, Wisconsin.
- Brodie, B. B., J. Axelrod, R. Soberman and B. B. Levy. 1949. The estimation of antipyrine in biological materials. J. Biol. Chem. 179: 25.
- Brody, S. and A. C. Ragsdale. 1924. The changes of form with age in the dairy cow. Mo. Agr. Exp. Sta. Res. Bul. 67.
- Brown, C. J., J. C. Hillier and J. A. Whatley. 1951. Specific gravity as a measure of the fat content of the pork carcass. J. Animal Sci. 10:97.
- Brown, Delos D., A. M. Mullins, R. S. Temple, R. F. Boulware and J. S. Sullivan, Jr. 1962. Relationship between certain carcass characteristics of purebred and crossbred cattle. J. Animal Sci. 21:388. (Abstr.).
- Brungardt, V. H. and R. W. Bray. 1963. Estimate of retail yield of the four major cuts in the beef carcass. J. Animal Sci. 22:177.
- Butler, O. D. 1957. The relation of conformation to carcass traits. J. Animal Sci. 16:227.

- Butler, O. D., L. E. Kunkle, F. A. Orts, G. T. King and T. C. Cartwright. 1961. Important beef carcass traits easily obtained. J. Animal Sci. 20:914. (Abstr.).
- Butterfield, R. M. 1963a. Estimation of carcass composition: The anatomical approach. Proc. Symp. Carcase Composition and Appraisal of Meat Animals: Australia. 4:1. (D. E. Tribe, editor).
- Butterfield, R. M. 1963b. Relative growth of the musculature of the ox. Proc. Symp. Carcase Composition and Appraisal of Meat Animals: Australia. 7:1. (D. E. Tribe, editor).
- Butterfield, R. M. 1965. Practical implications of anatomical research in beef cattle. Proc. New Zealand Soc. An. Prod. 25:152.
- Cahill, V. R., L. E. Kunkle, R. F. Plimpton, Jr. and Earle W. Klosterman. 1961. Edible portion and beef carcass evaluation. J. Animal Sci. 20:914. (Abstr.).
- Callow, E. H. 1947. Comparative studies of meat. I. The chemical composition of fatty and muscular tissue in relation to growth and fattening. J. Agr. Sci. 37:113.
- Carpenter, Z. L., G. T. King and Maurice Shelton. 1965. Determinants of lamb carcass cutability. J. Animal Sci. 24:861. (Abstr.).
- Carter, R. C. and C. M. Kincaid. 1959. Estimates of genetic and phenotypic parameters in beef cattle. III. Genetic and phenotypic correlations among economic characters. J. Animal Sci. 18:331.
- Clark, R. T. 1954. Heritability of beef carcass characteristics. Proc. Recip. Meat Conf. 7:206.
- Cole, J. W., L. E. Orme and C. M. Kincaid. 1960. Relationship of loin eye area, separable lean of various beef cuts and carcass measurements to total carcass lean in beef. J. Animal Sci. 19:89.
- Cole, J. W., C. B. Ramsey and R. H. Epley. 1962. Simplified method for predicting pounds of lean in beef carcasses. J. Animal Sci. 21:355.
- Cole. J. W., C. B. Ramsey, C. S. Hobbs and R. S. Temple. 1964. Effects of type and breed of British, Zebu and dairy cattle on production, palatability and composition. III. Percent wholesale cuts and yield of edible portion as determined by physical and chemical analysis. J. Animal Sci. 23:71.
- Cook, A. C., H. L. Kohli and W. H. Dawson. 1951. Relationship of five body measurements to slaughter grade, carcass grade and dressing percentage in Milking Shorthorn steers. J. Animal Sci. 10:386.

- Crown, R. M. and R. A. Damon, Jr. 1960. The value of the 12th rib cut for measuring beef carcass yields and meat quality. J. Animal Sci. 19:109.
- Dawson, W. T., R. L. Hiner and L. L. Madsen. 1955. Evaluating beef and dual purpose cattle by the proportion of weight carried on the front and hind legs. J. Animal Sci. 14:1179.
- Duncan, D. B. 1955. Multiple range and multiple F tests. Biometrics 11:1.
- Dumont, B. L., P. Le Guelte and J. Arnoux. 1961. Biometric study of beef cattle. II. Estimation of the weight of the musculature in Charolais cattle. Ann. de Zootechn. 10:321.
- Ensminger, M. E. 1955. Animal Science. The Interstate Co., Dansville, Ill. p. 361.
- Field, R. A., C. O. Schoonover, G. E. Nelms and C. J. Kercher. 1963.

  Evaluations of some old and new beef carcass measurements. J. Animal Sci. 22:826. (Abstr.).
- Gillett, T. A., A. M. Pearson and A. H. Kirton. 1965. Variation in potassium and sodium in muscles of the pig. J. Animal Sci. 24:177.
- Gnaedinger, R. H., E. P. Reineke, A. M. Pearson, W. D. Van Huss, J. A. Wessel and H. J. Montoye. 1963. Determination of body density by air displacement, helium dilution, and underwater weighing. Ann. N.Y. Acad. Sci. 110:96.
- Good, D. L., G. M. Dahl, S. Wearden and D. J. Weseli. 1961. Relationships among live and carcass characteristics of selected slaughter steers.

  J. Animal Sci. 20:698.
- Gottsch, A. Harold, R. A. Merkel and D. L. Mackintosh. 1961. Relationship of muscle, fat and bone and some physical measurements to beef cutability. J. Animal Sci. 20:917. (Abstr.).
- Green, W. W., G. L. Jessup, Jr. and F. E. White. 1955. Intercorrelations of weights of wholesale cuts of beef carcasses. J. Animal Sci. 14:1059.
- Gregory, Keith E., R. M. Koch, L. N. Hazel and Doyle Chambers. 1961.

  Principles of record of performance in beef cattle. Nebr. Agr. Exp.

  Circ. 106.
- Hammond, J. and A. B. Appleton. 1932. Growth and development of mutton qualities in the sheep. Oliver and Boyd, Edinburgh, Scotland.
- Hankins, O. G. and P. E. Howe. 1946. Estimation of the composition of beef carcasses and cuts. U.S.D.A. Tech. Bul. 926.

- Hankins, O. G. and H. W. Titus. 1939. Growth, fattening and meat production. U.S.D.A. Yearbook of Agr. p. 450.
- Hedrick, H. B., W. E. Meyer, M. A. Alexander, J. F. Lasley, J. E. Comfort, A. J. Dyer and H. D. Naumann. 1963. Indices of meatiness in beef. Mo. Agr. Exp. Sta. Res. Bul. 820.
- Henderson, D. W., D. E. Goll and E. A. Kline. 1966. Relationships of muscling and finish measurements from three different groups of beef carcasses with carcass yield. J. Animal Sci. 25:323.
- Henry, W. A., and J. W. Sanborn. 1883-1890. Pig feeding experiments and feeding for lean meat. Wis. and Mo. Agr. Exp. Sta. An. Rep. 4.
- Kirton, A. H. 1964. How important is conformation. Proc. Ruakura Farmers' Conference Week. New Zealand Dept. of Agr.
- Kirton, A. H., and A. M. Pearson. 1963a. Comparison of methods of measuring potassium in pork and lamb and prediction of their composition from sodium and potassium. J. Animal Sci. 22:125.
- Kirton, A. H. and A. M. Pearson. 1963b. Relationships between potassium content and body composition. Ann. N.Y. Acad. Sci. 110:221.
- Kirton, A. H., A. M. Pearson, R. H. Nelson, E. C. Anderson and R. L. Schuch. 1961. Use of naturally occurring potassium-40 to determine the carcass composition of live sheep. J. Animal Sci. 20:635.
- Knapp, Bradford, Jr., A. L. Baker, J. R. Quesenberry, and R. T. Clark. 1941. Record of performance in Hereford cattle. Mont. Agr. Exp. Sta. Bul. 397.
- Knapp, Bradford, Jr. and W. H. Black. 1942. A discussion of length of feeding period in beef cattle record of performance tests. J. Animal Sci. 1:65. (Abstr.).
- Knapp, Bradford, Jr. and R. T. Clark. 1950. Revised estimates of heritability of economic characteristics in beef cattle. J. Animal Sci. 9:582.
- Knapp, Bradford, Jr. and A. W. Nordskog. 1946a. Heritability of growth and efficiency in beef cattle. J. Animal Sci. 5:62.
- Knapp, Bradford, Jr. and A. W. Nordskog. 1946b. Heritability of live animal scores, grades and certain carcass characteristics in beef cattle. J. Animal Sci. 5:194.
- Knox, J. H. and Marvin Koger. 1946. A comparison of gains and carcasses produced by three types of feeder steers. J. Animal Sci. 5:331.
- Kropf, D. H. and R. L. Graf. 1959. The effect of carcass grade, weight and classification upon boneless beef yield. J. Animal Sci. 18:95.

- Lawes, J. B. and J. H. Gilbert. 1860. On the composition of oxen, sheep and pigs and their increase whilst fattening. J. Roy. Agr. Soc. 21.
- Lewis, R. W., V. H. Brungardt and R. W. Bray. 1964. Influence of subcutaneous fat contours in estimating trimmable fat and retail yield of heifer carcasses. J. Animal Sci. 23:1203. (Abstr.).
- Luitingh, H. C. 1962. Developmental changes in beef steers as influenced by fattening, age and type of ration. J. Agr. Sci. 58:1.
- Lush, J. L. 1928. Changes in body measurements of steers during intensive fattening. Tex. Agr. Exp. Sta. Bul. 385.
- MacLeod, John. 1964. The relationship of some linear and area measurements to muscling in pork carcasses. M.S. Thesis, Michigan State University, East Lansing, Michigan.
- Mc Meekan, C. P. 1941. Growth and development in the pig, with special reference to carcass quality characters. J. Agr. Sci. 31:1.
- Meek, A. 1901. Growth of farm Ungulates I. Approach from a study of the external characters. The Veterinarian. 74:335.
- Miller, J. C., H. B. Hedrick, G. B. Thompson, R. R. Freitag, W. E. Meyer, A. J. Dyer and H. D. Naumann. 1965. Factors affecting <u>longissimus</u> dorsi and subcutaneous fat measurements and indices of beef carcass cut-out. Mo. Agr. Exp. Sta. Bul. 880.
- Moore, F. D. 1946. Determination of total body water and solids with isotopes. Science 104:157.
- Morales, M. F., E. N. Rathbun, R. E. Smith and N. Pace. 1945. Studies on body composition II. Theoretical considerations regarding the major body tissue components, with suggestions for application to man. J. Biol. Chem. 158:677.
- Mounib, M. S. and J. V. Evans. 1957. Comparison between three methods used for the preparation of tissues for determinations of potassium and sodium. Analyst. 92:522.
- Murphy, C. E., D. K. Hallet, W. E. Tyler and J. C. Pierce, Jr. 1960. Estimating yields of retail cuts from beef carcasses. J. Animal Sci. 19: 1240. (Abstr.).
- Naumann, H. D. 1952. A recommended procedure for measuring and grading beef for carcass evaluation. Proc. Ann. Recip. Meat Conf. 5:108.
- Orme. L. E. 1958. Methods of estimating carcass characteristics in beef. Ph.D. Thesis. Michigan State University, East Lansing, Michigan.
- Orme. L. E. 1963. Estimating composition from linear measurements, live probe and body weight. Ann. N.Y. Acad. Sci. 110:307.

- Orme. L. E., J. W. Cole, C. M. Kincaid and R. J. Cooper. 1960. Predicting total carcass lean in mature beef from weights of certain entire muscles. J. Animal Sci. 19:726.
- Orme, L. E., A. M. Pearson, L. J. Bratzler, W. T. Magee and A. C. Wheeler. 1959. The muscle-bone relationship in beef. J. Animal Sci. 18:1271.
- Pace, N., L. Kline, H. K. Schachman and M. Harfenist. 1947. Studies on body composition. IV. Use of radioactive hydrogen for measurement in vivo of total body water. J. Biol. Chem. 168:459.
- Palmer, A. Z., J. W. Carpenter, R. L. Reddish, C. E. Murphey and D. K. Hallet. 1961. Estimated and actual yields of boneless retail cuts from Brahman crossbred cattle and carcasses. J. Animal Sci. 20:919. (Abstr.).
- Palsson, H. 1939. Meat qualities in sheep with special reference to Scottish breeds and crosses. J. Agr. Sci. 29:544.
- Palsson, H. and J. B. Verges. 1952. Effects of the plane of nutrition on growth and the development of carcass quality in lambs. Part I. The effects of high and low planes of nutrition at different ages. J. Agr. Sci. 42:1.
- Pearson, A. M. 1965. Body composition. Newer Methods of Nutritional Biochemistry. Academic Press, New York (A. A. Albanese, editor).
- Ramsey, C. B., J. W. Cole and C. S. Hobbs. 1962. Relation of beef carcass grades, proposed yield grades and fat thickness to separable lean, fat and bone. J. Animal Sci. 21:193.
- Shelby, C. E., R. T. Clark, J. R. Quesenberry and R. R. Woodward. 1960. Heritability of some economic characteristics in record of performance bulls. J. Animal Sci. 19:450.
- Snedecor, G. W. 1956. Statistical Methods (5th ed.) Iowa State College Press, Ames, Iowa.
- Swiger, L. A., K. E. Gregory, L. J. Sumption and B. C. Breidenstein. 1964.

  The importance of measuring cut-out in cattle. J. Animal Sci. 23:854.

  (Abstr.).
- Tallis, G. M., Earle W. Klosterman and V. R. Cahill. 1957. Body measurements in relation to beef carcass characteristics. J. Animal Sci. 16:1027.
- Trowbridge, P. F., C. R. Moulton and L. D. Haigh. 1918. Composition of the beef animal and energy costs of fattening. Mo. Agr. Exp. Sta. Res. Bul. 30.

- Wallace, L. R. 1948. Growth of lambs before and after birth in relation to the level of nutrition. Part I. J. Agr. Sci. 38:93.
- Warner, K. F., N. R. Ellis and P. E. Howe. 1934. Cutting yields of hogs an index of fatness. J. Agr. Res. 48:241.
- Warwick, E. J. 1958. Fifty years of progress in breeding beef cattle. J. Animal Sci. 17:922.
- Wellington, G. H. 1953. Recommended procedure for cutting beef. Proc. Ann. Recip. Meat Conf. 6:73.
- White, F. E. and W. W. Green. 1952. Relationships of measurements of live animals to weights of wholesale cuts of beef. J. Animal Sci. 11:370.
- Whiteman, J. V., J. C. Hillier and J. A. Whatley. 1951. Carcass studies on hogs of different breeding. J. Animal Sci. 10:638.
- Wilson, James and C. F. Curtis. 1893. Steer feeding. Iowa Agr. Exp. Sta. Bul. 20. p. 639.
- Wythe, L. D., Jr., F. A. Orts and G. T. King. 1961. Bone-muscle relationships in beef carcasses. J. Animal Sci. 20:3.
- Zinn, Dale W., R. M. Durham and Robert Stovall. 1963. Muscle growth and development in the beef animal during the feeding period. J. Animal Sci. 22:829. (Abstr.).
- Zinn, D. W., H. Elliott, D. Burnett and R. M. Durham. 1961. Evaluation of U.S.D.A. beef grading methods. J. Animal Sci. 20:922.




Table A. Coefficient of determination between weight of carcass separable fat and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (combined region)

weight groups).				
	Regression	Beta		R21
Independent variable	coefficients	weights	Sig.	deletes
Constant	-41.671		0.00	
Carcass wt.	0.176	0.655	0.00	0.68
Probe-5th thoracic-4 in.	1.578	0.254	0.00	0.90
Probe-8th thoracic-12 in.	931	165	0.00	0.93
Probe-1st lumbar-12 in.	0.642	0.187	0.00	0.92
Probe-6th lumbar-12 in.	346	113	0.01	0.93
Probe-Av. 3rd sacral <sup>2</sup>	1.050	0.204	0.00	0.92
Fat measurement B	0.367	0.140	0.00	0.93
Fat measurement D	0.796	0.180	0.00	0.92

 $R^2 = 0.94$  Standard error of estimate - 6.67

Table B. Coefficient of determination between percent carcass separable fat and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (combined weight groups).

	Regression	Beta		R <sup>21</sup>
Independent variable	coefficients	weights	Sig.	deletes
Constant	23.515		0.00	
		074	- <del>-</del>	0.05
Carcass wt.	004	074	0.20	0.85
Probe-5th thoracic-4 in.	0.573	0.447	0.00	0.76
Probe-8th thoracic-8 in.	274	254	0.00	0.83
Probe-1st lumbar-12 in.	0.215	0.304	0.00	0.82
Probe-6th lumbar-12 in.	092	146	0.02	0.84
Probe-Av. 3rd sacra12	0.358	0.338	0.00	0.80
Fat measurement B	0.100	0.186	0.00	0.83
Fat measurement D	0.249	0.273	0.00	0.82

 $R^2 = 0.86$  Standard error of estimate = 2.10

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>2</sup>Average of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Average of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

Table C. Coefficient of determination between weight of carcass separable fat and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to

750 lb. weight group).

	Regression	Beta		RZI
Independent variables	coefficients	weights	Sig.	deletes
Constant	-55.228		0.22	
Carcass wt.	0.098	0.115	0.10	0.87
Probe-5th thoracic-4 in.	2.191	0.482	0.00	0.72
Probe-8th thoracic-8 in.	-1.470	434	0.00	0.83
Probe-1st lumbar-8 in.	702	327	0.00	0.84
Probe-1st lumbar-12 in.	1.496	0.559	0.00	0.74
Probe-4th lumbar-8 in.	0.464	0.241	0.01	0.85
Probe-3rd sacral-8 in.	0.839	0.291	0.00	0.84
Circumference of round	1.649	0.176	0.02	0.86
Fat measurement D	0.941	0.334	0.00	0.84

 $R^2 = 0.88$  Standard error of estimate = 6.47

Table D. Coefficient of determination between percent carcass separable fat and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to 750 lb weight group)

<u>ъ).                                    </u>		<u> </u>	
Regression	Beta		R <sup>21</sup>
coefficients	weights	Sig.	deletes
17.367		0.13	
018	078	0.24	0.89
0.573	0.474	0.00	0.74
371	411	0.00	0.84
174	305	0.00	0.85
0.390	0.548	0.00	0.76
0.120	0.236	0.01	0.86
0.240	0.314	0.00	0.84
0.446	0.178	0.02	0.88
0.274	0.366	0.00	0.84
	Regression coefficients 17.367 018 0.573 371 174 0.390 0.120 0.240 0.446	Regression beta weights  17.367018078 0.573 0.474371411174305 0.390 0.548 0.120 0.236 0.240 0.314 0.446 0.178	Regression coefficients         Beta weights         Sig.           17.367         0.13          018        078         0.24           0.573         0.474         0.00          371        411         0.00           0.390         0.548         0.00           0.120         0.236         0.01           0.240         0.314         0.00           0.446         0.178         0.02

 $R^2 = 0.89$  Standard error of estimate = 1.66

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

Table E. Coefficient of determination between percent carcass separable fat and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (500 to 550 lb weight group)

	Regression	Beta	Level of	R <sup>21</sup>
Independent variables	coefficients	weights	sig.	deletes
Constant	21.047		0.00	
Probe-5th thoracic-4 in.	0.852	0.618	0.00	0.67
Probe-8th thoracic-8 in.	248	190	0.09	0.83
Probe-1st lumbar-12 in.	0.036	0.041	0.72	0.85
Probe-Av. 3rd sacral <sup>2</sup>	0.353	0.308	0.00	0.79
Fat measurement D	0.316	0.293	0.00	0.80

 $R^2 = 0.85$  Standard error of estimate = 2.46

Table F. Coefficient of determination between percent carcass separable fat and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to 750 lb weight group)

/50 lb. weight gi	roup).			1
	Regression	Beta	Level of	R21
Independent variables	coefficient	weights	sig.	deletes
Constant	19.984		0.00	
Probe-5th thoracic-4 in.	0.524	0.434	0.00	0.71
Probe-8th thoracic-8 in.	294	327	0.01	0.80
Probe-1st lumbar-12 in.	0.267	0.374	0.00	0.76
Probe-Av. 3rd sacral <sup>2</sup>	0.411	0.431	0.00	0.76
Fat measurement D	0.212	0.283	0.01	0.80

 $R^2 = 0.83$  Standard error of estimate = 1.93

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Average of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

Average of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

Table G. Coefficient of determination between percent external fat trim from the round, loin, rib and chuck and a combination of objective carcass measurements, and the regression coefficient for each

of the measurements (combined weight groups). Level of Regression Beta Independent variables delet<u>es</u> coefficients weights sig. 8.430 0.00 Constant 0.011 0.00 0.63 Carcass wt. 0.524 Probe-4th lumbar-12 in. 0.104 0.355 0.00 0.64 Probe-Av. 3rd sacral<sup>2</sup> 0.00 0.138 0.334 0.64 Length of round -.543 -.518 0.00 0.62

 $R^2 = 0.71$  Standard error of estimate = 1.12

Table H. Coefficient of determination between percent external fat trim from the round, loin, rib and chuck and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (500 to 550 lb. weight group).

the measurements	(300 60 330 10.	METRIC BI	.oup).	
	Regression	Beta	·Level of	R21
Independent variable	coefficients	weights	sig.	deletes
Constant	4.769		0.43	
Carcass wt.	<b></b> 025	183	0.02	0.91
Probe-5th thoracic-8 in.	081	246	0.00	0.90
Probe-8th thoracic-12 in.	140	254	0.04	0.92
Probe-11th thoracic-4 in.	0.148	0.335	0.01	0.91
Probe-11th thoracic-12 in.	129	250	0.03	0.91
Probe-6th lumbar-4 in.	0.114	0.272	0.01	0.91
Probe-6th lumbar-12 in.	0.103	0.354	0.00	0.90
Probe-3rd sacral-12 in.	0.086	0.332	0.00	0.87
Probe-5th sacral-4 in.	117	395	0.00	0.88
Circumference of round	0.230	0.138	0.03	0.92
Fat measurement A	0.118	0.387	0.00	0.89
Fat measurement B	0.053	0.251	0.02	0.91
Fat measurement C	0.177	0.454	0.00	0.90
Fat measurement E	0.162	0.611	0.00	0.85
Fat measurement F	163	660	0.00	0.88

R<sup>2</sup> = 0.93 Standard error of estimate = 0.77

1 Value of coefficient of determination if that variable was deleted from the analysis.

<sup>1</sup> Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Average of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

Table I. Coefficient of determination between percent external fat trim from the round, loin, rib and chuck and a combination of objective measurements, and the regression coefficient for each of the measurements (700 to 750 lb. weight group).

measurements (700	to 750 lb. wei	ght group).	<u> </u>	
	Regression	Beta	Level of	R21
Independent variables	coefficient	weights	sig.	deletes
Constant	9.229		0.03	
Carcass wt.	019	235	0.01	0.94
Probe-5th thoracic-4 in.	0.096	0.228	0.01	0.94
Probe-5th thoracic-8 in.	0.062	0.277	0.00	0.93
Probe-8th thoracic-4 in.	175	435	0.00	0.93
Probe-8th thoracic-12 in.	281	<b></b> 796	0.00	0.85
Probe-11th thoracic-4 in.	272	717	0.00	0.92
Probe-1st lumbar-4 in.	0.312	0.837	0.00	0.88
Probe-1st lumbar-8 in.	139	699	0.00	0.91
Probe-1st lumbar-12 in.	0.153	0.618	0.00	0.89
Probe-4th lumbar-8 in.	0.120	0.672	0.00	0.92
Probe-4th lumbar-12 in.	0.094	0.430	0.00	0.92
Probe-3rd sacral-8 in.	0.405	1.518	0.00	0.89
Probe-Av. 3rd sacral2	641	-1.931	0.00	0.90
Probe-5th sacral-12 in.	0.104	0.289	0.00	0.92
Fat depth over brisket	-1.043	595	0.00	0.89
Circumference of round	0.216	0.248	0.01	0.94
Fat measurement B	0.051	0.280	0.00	0.93
Fat measurement D	0.265	1.015	0.00	0.85
Fat measurement E	0.109	0.564	0.01	0.94
Fat measurement F	134	750	0.00	0.93
Fat measurement G	0.101	0.531	0.00	0.93

 $R^2 = 0.96$  Standard error of estimate = 0.46

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Average of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

Table J. Coefficient of determination between percent total retail fat trim and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (combined weight groups).

weight groups).				
	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
_				
Constant	23.525		0.00	
Carcass wt.	0.019	0.347	0.01	0.79
Probe-5th thoracic-4 in.	0.241	0.191	0.03	0.80
Probe-8th thoracic-8 in.	294	277	0.00	0.79
Probe-11th thoracic-4 in.	0.212	0.191	0.04	0.80
Probe-1st lumbar-12 in.	0.165	0.237	0.02	0.80
Probe-4th lumbar-12 in.	0.187	0.245	0.00	0.79
Probe-6th lumbar-8 in.	0.096	0.152	0.05	0.80
Probe-3rd sacral-4 in.	212	216	0.03	0.80
Probe-Av. 3rd sacral <sup>2</sup>	0.387	0.372	0.00	0.77
Depth of fat over brisket	1.088	0.184	0.01	0.79
Length of round	-1.049	396	0.00	0.78
Fat measurement G	115	185	0.04	0.80

 $R^2 = 0.81$  Standard error of estimate = 2.41

<sup>2</sup>Average of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

Table K. Coefficient of determination between weight of total retail fat trim and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to 750 lb. weight group).

/50 lb. weight gi	roup).			
Independent variables	Regression coefficient	Beta weights	Level of sig.	R <sup>21</sup> deletes
Constant	32.886		0.41	
Carcass wt.	0.008	0.013	0.85	0.80
Probe-5th thoracic-4 in.	0.962	0.285	0.01	0.75
Probe-6th lumbar-8 in.	0.637	0.368	0.00	0.72
Probe-6th lumbar-12 in.	455	274	0.02	0.76
Probe-3rd sacral-12 in.	0.622	0.252	0.04	0.77
Probe-5th sacral-8 in.	0.714	0.351	0.00	0.72
Fat measurement D	0.782	0.374	0.00	0.73
Fat measurement F	377	264	0.04	0.77

 $R^2 = 0.80$  Standard error of estimate = 6.22

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

Table L. Coefficient of determination between percent total retail fat trim and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to

/50 lb. weight gr	oup).			1
Independent variables	Regression coefficients	Beta weights	Level of sig.	R <sup>Z±</sup> deletes
Constant	42.801		0.00	
Carcass wt.	042	233	0.02	0.68
Probe-4th lumbar-12 in.	0.144	0.286	0.03	0.68
Probe-6th lumbar-8 in.	0.177	0.357	0.01	0.66
Probe-6th lumbar-12 in.	149	314	0.01	0.67
Probe-3rd sacral-12 in.	0.243	0.344	0.00	0.64
Probe-5th sacral-8 in.	0.219	0.375	0.00	0.64

 $R^2 = 0.73$  Standard error of estimate = 2.00

Table M. Coefficient of determination between percent carcass separable muscle and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (combined weight groups).

	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	62.175		0.00	
Carcass wt.	0.0004	008	0.86	0.73
Probe-5th thoracic-4 in.	289	286	0.00	0.69
Probe-Av. 3rd sacral <sup>2</sup>	400	479	0.00	0.59
Fat measurement C	173	234	0.01	0.70

 $R^2 = 0.73$  Standard error of estimate = 2.20

Table N. Coefficient of determination between weight of carcass separable muscle and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (500 to 550 lb. weight group).

.о сру.			
Regression coefficients	Beta weights	Level of	R <sup>21</sup> deletes
		<u> </u>	
96.744		0.01	
0.128	0.190	0.09	0.68
-1.144	438	0.00	0.61
-1,231	566	0.00	0.51
	Regression coefficients 96.744 0.128 -1.144	Regression Beta coefficients weights  96.744 0.128 -1.144438	Regression coefficients         Beta weights         Level of sig.           96.744         0.01           0.128         0.190         0.09           -1.144        438         0.00

 $R^2 = 0.71$  Standard error of estimate = 6.23

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>1</sup> Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Average of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Average of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

Table 0. Coefficient of determination between weight of carcass separable muscle and a combination of objective carcass measurements. and the regression coefficient for each of the measurements

(700 to 750 lb. weight group).

Independent variables	Regression coefficients	Beta weights	Level of sig.	R <sup>21</sup> deletes
Constant Carcass wt.	111.062 0.149	0.225	0.02 0.03	0.63
Probe-5th thoracic-4 in. Probe-Av. 3rd sacra12	828 -1.847	235 664	0.05 0.00	0.64 0.37

 $R^2 = 0.68$ Standard error of estimate - 7.61

Table P. Coefficient of determination between percent carcass separable muscle and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to 750 1h waight grown)

	group).			1
Independent variables	Regression coefficients	Beta weights	Level of sig.	R <sup>2</sup> deletes
Constant Carcass wt. Probe-5th thoracic-4 in. Probe-Av. 3rd sacral2	77.117 022 275 474	121 277 607	0.00 0.23 0.02 0.00	0.64 0.60 0.39

 $R^2 = 0.65$ Standard error of estimate = 2.23

Table 0. Coefficient of determination between weight of carcass separable bone and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (500 to 550 lb. weight group).

	Regression	Beta	Level of	R <sup>21</sup>
Independent variables	coefficients	weights	sig.	deletes
Constant .	-16.544		0.29	
Carcass wt.	0.021	0.090	0.32	0.78
Probe-5th thoracic-4 in.	474	533	0.00	0.64
Length of round	1.501	0.480	0.00	0.66

 $R^2 = 0.78$ Standard error of estimate = 1.83

 $<sup>{}^{</sup> extsf{1}}$ Value of coefficient of determination if that variable was deleted from

<sup>&</sup>lt;sup>2</sup>Average of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

 $<sup>^{\</sup>mathrm{l}}$ Value of coefficient of determination if that variable was deleted from the analysis. <sup>2</sup>Average of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

 $<sup>^{</sup>m l}$ Value of coefficient of determination if that variable was deleted from the analysis.

Table R. Coefficient of determination between percent carcass separable bone and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (500

to 550 lb. weight group). R21 Regression Beta Level of weights deletes Independent variables coefficients sig. Constant 4.603 0.44 0.09 -.013 0.81 Carcass wt. **-.138** -.174 0.00 0.72 Probe-5th thoracic-4 in. -.468 0.72 Length of round 0.577 0.442 0.00

 $R^2 = 0.83$  Standard error of estimate = 0.68

Table S. Coefficient of determination between weight of carcass separable bone and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to 750 lb. weight group).

750 ID. WEIGHT	Regression	Beta	Level of	R <sup>21</sup>
Independent variables	coefficients	weights	sig.	deletes
Constant	-55.665		0.00	
Carcass wt.	0.052	0.238	0.01	0.75
Probe-Av. 3rd sacral <sup>2</sup>	<b></b> 358	390	0.00	0.69
Length of round	2.096	0.581	0.00	0.57

 $R^2 = 0.80$  Standard error of estimate = 1.98

Table T. Coefficient of determination between percent carcass separable bone and a combination of objective carcass measurements, and the regression coefficient for each of the measurements (700 to 750 lb. weight group).

750 ID. WEIGHT	Regression	Beta	Level of	R <sup>21</sup>
Independent variables	coefficients	weights	sig.	deletes
Constant	-5.973		0.11	
Carcass wt.	00004	001	0.94	0.77
Probe-Av. 3rd Sacral <sup>2</sup>	092	380	0.00	0.66
Length of round	0.612	0.640	0.00	0.49

 $R^2 = 0.77$  Standard error of estimate = 0.57

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Average of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Average of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

Table U. Coefficient of determination between weight of carcass separable muscle and a combination of objective carcass measurements and weight of flank separable components, and the regression coefficient for each of the variables (500 to 550 lb. weight group).

CTERT TOT EACH OF	the variables	(300 60 330	ID. WEIGHT	group).
	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	27.15 <b>4</b>		0.30	
Carcass wt.	0.247	0.366	0.00	0.82
Probe-Av. 3rd sacra12	<b></b> 777	<b></b> 357	0.00	0.82
Wt. separable flank muscle	3.317	0.290	0.00	0.82
Wt. separable flank fat	-2.176	<b></b> 570	0.00	0.76
Wt. kidney knob-left side	<b></b> 768	204	0.01	0.87
-				

 $R^2 = 0.89$  Standard error of estimate = 3.96

Table V. Coefficient of determination between percent carcass separable muscle and a combination of objective carcass measurements, flank separable component weights and kidney knob weight, and the regression coefficient for each of the variables (700 to 750 lb. weight group).

Independent variables	Regression coefficients	Beta weights	Level of sig.	R <sup>21</sup> deletes
Constant	66.132		0.00	
Carcass wt.	008	045	0.50	0.91
Probe-5th thoracic-4 in.	157	158	0.02	0.89
Probe-Av. 3rd sacral <sup>2</sup>	154	197	0.02	0.89
Wt. separable flank muscle	0.973	0.296	0.00	0.85
Wt. separable flank fat	658	571	0.00	0.79
Wt. kidney knob-left side	167	142	0.02	0.89

 $R^2 = 0.91$  Standard error of estimate = 1.18

Table W. Coefficient of determination between weight of carcass separable muscle and a combination of carcass weight and weights of flank separable components, and the regression coefficient for each of the variables (500 to 550 lb. weight group).

Independent variables	Regression coefficients	Beta weights	Level of	R21 deletes
Independent Vallables	coefficients	weights	sig.	dereces
Constant	27.984		0.40	
Carcass wt.	0.214	0.317	0.00	0.73
Wt. separable flank muscle	4.986	0.435	0.00	0.62
Wt. separable flank fat	-3.246	850	0.00	0.39

R<sup>2</sup> = 0.79 Standard error of estimate = 5.29

<sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Average of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

<sup>1</sup> Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>2</sup>Average of 3 probes, 4, 8 and 12 in. off the carcass dorsal midline.

Table X. Coefficient of determination between weight of carcass separable muscle and a combination of carcass weight and weights of flank separable components, and the regression coefficient for each of the independent variables (700 to 750 lb weight group)

or the independent	variables (700	E0 /30 II	o. weight gi	toup).
	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	6 <b>0.400</b>		0.06	
Carcass wt.	0.193	0.291	0.00	0.81
Wt. separable flank muscle	4.321	0.370	0.00	0.76
Wt. separable flank fat	-3.231	<b></b> 788	0.00	0.39

 $R^2 = 0.86$  Standard error of estimate = 4.96

Table Y. Coefficient of determination between percent carcass separable muscle and a combination of carcass weight and weights of flank separable components, and the regression coefficient for each of the variables (500 to 550 lb. weight group).

R21 Level of Regression Beta deletes Independent variables coefficients weights sig. 0.00 Constant 60.932 0.70 0.83 Carcass wt. -.010 -.036 0.00 0.72 Wt. separable flank muscle 1.592 0.340 Wt. separable flank fat -1.228 -.786 0.00 0.48

 $R^2 = 0.83$  Standard error of estimate = 1.97

Table Z. Coefficient of determination between percent carcass separable muscle and a combination of carcass weight and weights of flank separable components, and the regression coefficient for each of the variables (700 to 750 lb. weight group).

02 010 102200	100 00 130 251	" U - D - V	<u> </u>	
	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	60.360		0.00	
Carcass wt.	004	020	0.79	0.86
Wt. separable flank muscle	1.059	0.322	0.00	0.78
Wt. separable flank fat	934	810	0.00	0.36
<del>-</del>				

 $R^2 = 0.86$  Standard error of estimate = 1.42

<sup>1</sup> Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

Table AA. Coefficient of determination between percent carcass separable muscle and a combination of muscle and bone weights, and the regression coefficient for each of the variables (combined

weight groups).				
Independent variables	Regression coefficients	Beta weights	Level of sig.	R <sup>21</sup> deletes
Constant	53.115		0.00	
Carcass wt.	066	-1.513	0.00	0.14
Semitendinosus wt.	2.932	0.463	0.00	0.71
Biceps femoris wt.	1.832	0.468	0.00	0.72
Longissimus dorsi wt.	0.883	0.378	0.00	0.75
Scapula wt.	5.036	0.462	0.00	0.71

 $R^2 = 0.78$  Standard error of estimate = 1.98

Table AB. Coefficient of determination between weight of carcass separable muscle and a combination of muscle and bone weights and the regression coefficient for each of the variables (500 to 550 lb weight group)

Ib. Weight group	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
_				
Constant	<b>-</b> 6.525		0.82	
Carcass wt.	0.014	0.021	0.77	0.88
Semitendinosus wt.	6.310	0.215	0.00	0.85
Biceps femoris wt.	<b>6.</b> 183	0.301	0.00	0.81
L. dorsi wt.	5.811	0.458	0.00	0.75
Femur wt.	-8.657	330	0.03	0.86
Tibia plus fibula wt.	9.337	0.305	0.01	0.86
Scapula wt.	12.288	0.270	0.03	0.86
Humerus wt.	8.033	0.204	0.03	0.86

 $R^2 = 0.88$  Standard error of estimate = 4.27

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

Table AC. Coefficient of determination between weight of carcass separable muscle and a combination of muscle weights, and the regression coefficient for each of the variables (700 to 750 lb. weight

Independent variables	Regression coefficient	Beta weights	Level of sig.	R <sup>2</sup> 1 deletes
Constant Carcass wt.	-8.902 0.112	0.168	0.82 0.07	0.70 0.46
<u>Biceps femoris</u> wt.	14.621 8.341	0.558 0.483	0.00 0.00	0.48

 $R^2 = 0.73$  Standard error of estimate = 6.94

Table AD. Coefficient of determination between percent carcass separable muscle and a combination of muscle weights, and the regression coefficient for each of the variables (700 to 750 lb. weight

Independent variables	Regression coefficient	Beta weights	Level of sig.	R <sup>21</sup> deletes
Constant	40.729		0.00	
Carcass wt.	030	161	0.06	0.74
Semitendinosus wt.	3.923	0.532	0.00	0.51
Biceps femoris wt.	2.560	0.527	0.00	0.52

 $R^2 = 0.77$  Standard error of estimate = 1.83

Table AE. Coefficient of determination between weight of carcass separable bone and a combination of muscle and bone weights, and the regression coefficient for each of the variables (500 to 550 lb. weight group)

weight group).	Regression	Beta	Level of	R <sup>21</sup>
Independent variables	coefficients	weights	sig.	deletes
Constant	-1.738		0.77	
Carcass wt.	004	016	0.71	0.95
Biceps femoris wt.	1.494	0.214	0.00	0.91
Tibia + fibula wt.	3.762	0.362	0.00	0.87
Scapula wt.	5.495	0.355	0.00	0.90
Humerus wt.	4.127	0.308	0.00	0.90

 $R^2 = 0.95$  Standard error of estimate = 0.92

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

Table AF. Coefficient of determination between percent carcass separable bone and a combination of muscle and bone weights, and the regression coefficient for each of the variables (500 to 550

lb. weight group)	•			1
	Regression	Beta	Level of	RZI
Independent variables	coefficients	weights	sig.	deletes
Constant	13.223		0.00	
Carcass wt.	025	265	0.00	0.91
Biceps femoris wt.	0.641	0.220	0.00	0.93
Psoas major wt.	465	094	0.02	0.96
Tibia + fibula wt.	1.434	0.330	0.00	0.90
Scapula wt.	2.241	0.347	0.00	0.92
Humerus wt.	1.233	0.220	0.00	0.94

 $R^2 = 0.96$  Standard error of estimate = 0.33

Table AG. Coefficient of determination between weight of carcass separable bone and a combination of muscle and bone weights, and the regression coefficient for each of the variables (700 to 750

ID. Weight group	) ·			
	Regression	Beta	Level of	R <sup>21</sup>
Independent variables	coefficients	weights	sig.	deletes
Constant	-17.789		0.10	
Carcass wt.	0.024	0.110	0.12	0.84
Semitendinosus wt.	2.123	0.245	0.00	0.80
Tibia + fibula wt.	6.853	0.566	0.00	0.61
Scapula wt.	5.060	0.312	0.00	0.79

 $R^2 = 0.85$  Standard error of estimate = 1.73

Table AH. Coefficient of determination between percent carcass separable bone and a combination of muscle and bone weights, and the regression coefficient for each of the variables (700 to 750 lb weight group)

ID. WEIGHT GIO	ш).			
Independent variables	Regression coefficients	Beta weights	Level of sig.	R <sup>21</sup> deletes
Constant	2.226		0.45	
Carcass wt.	004	064	0.37	0.88
Semitensinosus wt.	0.516	0.225	0.00	0.84
Biceps femoris wt.	0.307	0.203	0.03	0.86
Tibia + fibula wt.	1.604	0.500	0.00	0.75
Scapula wt.	1.286	0.299	0.00	0.83

 $R^2 = 0.88$  Standard error of estimate = 0.42

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

Table AI. Coefficient of determination between weight of round, loin, rib and chuck retail yield and a combination of wholesale cut weights and yields, and the regression coefficient for each

of the variables (500 to 550 lb. weight group).

	Regression	Beta	Level of	R <sup>21</sup>
Independent variables	coefficients	weights	sig.	deletes
Constant	-117.856		0.00	
Carcass wt.	0.234	0.376	0.00	0.76
Length of round	3.306	0.388	0.00	0.80
Weight of round-right side	0.747	0.258	0.02	0.82
Weight of shank bone-right				
side	-5.740	275	0.04	0.83
Weight of flank yield-				
right side	2.141	0.357	0.00	0.76
Weight of flank fat-right sid	de <b>-1.</b> 585	510	0.00	0.75
L. dorsi muscle area	1.642	0.213	0.02	0.82

 $R^2 = 0.85$  Standard error of estimate = 4.39

Table AJ. Coefficient of determination between weight of round, loin, rib and chuck retail yield and a combination of objective carcass measurements, carcass scores and flank yield and fat trim weights, and the regression coefficient for each of the variables (700 to 750 lb. weight group).

	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	-101.832		0.04	
Carcass wt.	0.239	0.429	0.00	0.64
Round score	2.387	0.272	0.03	0.72
Langth of round	3.382	0.369	0.01	0.70
Wt. flank yield	2.321	0.379	0.00	0.64
Wt. flank fat	-2.144	619	0.00	0.50

 $R^2 = 0.76$  Standard error of estimate = 5.72

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

Table AK. Coefficient of determination between weight of round, loin, rib and chuck retail yield and a combination of carcass scores, measurements, foreshank bone weight and percent flank yield, and the regression coefficient for each of the variables (combined weight groups)

bined weight groups).				
Regression	Beta	Level of	RZI	
coefficients	weights	sig.	deletes	
-99.978		0.00		
0.210	0.878	0.00	0.81	
1.986	0.135	0.00	0.93	
4.389	0.378	0.00	0.92	
-1.035	105	0.03	0.94	
-4.613	146	0.04	0.94	
0.946	0.346	0.00	0.87	
	Regression coefficients -99.978 0.210 1.986 4.389 -1.035 -4.613	Regression Beta weights  -99.978 0.210 0.878 1.986 0.135 4.389 0.378 -1.035 -1.05 -4.613146	Regression coefficients         Beta weights         Level of sig.           -99.978         0.00           0.210         0.878         0.00           1.986         0.135         0.00           4.389         0.378         0.00           -1.035        105         0.03           -4.613        146         0.04	

R<sup>2</sup> = 0.94 Standard error of estimate - 5.60 <sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

Table AL. Coefficient of determination between weight of round, loin, rib and chuck retail yield and a combination of objective carcass measurements and percent flank fat trim, and the regression coefficient for each of the variables (500 to 550 lb. weight group)

group).	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	-73.816		0.10	
Carcass wt.	0.249	0.399	0.00	0.67
Length of round	3.390	0.397	0.00	0.70
L. dorsi area	2.589	0.335	0.00	0.70
% flank fat trim	689	674	0.00	0.52

 $R^2 = 0.79$  Standard error of estimate = 5.00 Value of coefficient of determination if that variable was deleted from the analysis.

Table AM. Coefficient of determination between percent round, loin, rib and chuck retail yield and a combination of objective carcass measurements, foreshank bone weight and percent flank fat trim, and the regression coefficient for each of the variables (500 to 550 lb. weight group).

	B			<del></del>
	Regression	Bet <b>a</b>	Level of	R <sup>21</sup>
Independent variables	coefficients	weights	sig.	deletes
Constant	21.636		0.17	
Carcass wt.	0005	002	0.93	0.83
Length of round	1.687	0.503	0.00	0.74
Wt. of foreshank bone-				
right side	-2.402	292	0.03	0.81
L. dorsi muscle area	0.957	0.315	0.00	<b>0.</b> 76
% flank fat trim	293	728	0.00	0.60
77 0 00	0. 1. 1	<u> </u>	. 1 77	

 $R^2 = 0.83$  Standard error of estimate = 1.77 Value of coefficient of determination if that variable was deleted from

Table AN. Coefficient of determination between weight of round, loin, rib and chuck retail yield and a combination of carcass measurements, scores and percent yield, and the regression coefficient for each of the variables (700 to 750 lb. weight group).

	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	-150.434		0.00	
Carcass wt.	0.185	0.333	0.00	0.66
Round score	2.431	0.277	0.02	0.72
Length of round	4.097	0.447	0.00	0.66
% flank yield	1.381	0.689	0.00	0.31

 $R^2 = 0.76$  Standard error of estimate - 5.57 lValue of coefficient of determination if that variable was deleted from the analysis.

Table AO. Coefficient of determination between percent round, loin, rib and chuck retail yield and a combination of carcass measurements, scores and percent flank yield, and the regression for each of the variables (700 to 750 lb. weight group).

Independent variables	Regression coefficients	Bet <b>a</b> weights	Level of sig.	R <sup>21</sup> deletes
Carcass wt.	026	199	0.06	0.63
Round score	0.600	<b>0.</b> 286	0.05	0.63
Length of round	1.152	0.524	0.00	0.53
% flank yield	0.338	0.702	0.00	0.20

R<sup>2</sup> = 0.67 Standard error of estimate = 1.57 <sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis. Table AP. Coefficient of determination between weight of carcass retail yield and carcass weight, flank yield and fat trim, and the regression coefficient for each of the variables (700 to 750 lb weight group).

ib. weight group).			
Regression	Beta weights	Level of	R <sup>21</sup> deletes
COCITICIENTS	WCIGITED	3+6.	<u>ucrees</u>
-32.346		0.41	
0.360	0.523	0.00	0.57
3.467	0.457	0.00	0.60
-2.749	642	0.00	0.42
	Regression coefficients  -32.346 0.360 3.467	Regression Beta weights  -32.346 0.360 0.523 3.467 0.457	Regression coefficients         Beta weights         Level of sig.           -32.346         0.41           0.360         0.523         0.00           3.467         0.457         0.00

 $R^2 = 0.80$  Standard error of estimate - 6.30

Table AQ. Coefficient of determination between percent carcass retail yield and a combination of kidney knob weight and flank retail yield and fat trim weights, and the regression coefficient for each of the variables (700 to 750 lb. weight group).

	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	54.413		0.00	
Carcass wt.	0.020	0.130	0.17	0.75
Kidney knob wtright side	210	238	0.01	0.71
Wt. flank yield	0.677	0.393	0.00	0.62
Wt. flank fat trim	693	712	0.00	0.32

 $R^2 = 0.76$  Standard error of estimate = 1.57

<sup>1</sup> Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

Table AR. Coefficient of determination between weight of carcass retail yield and a combination of objective carcass measurements and percent flank yield, and the regression coefficient for each of the variables (combined weight groups).

(combined weigh	nt groups).	·	
Regression	Beta	Level of	R <sup>21</sup>
coefficients	weights	sig.	deletes
-78.712		0.00	
0.243	0.837	0.00	0.78
1.864	0.132	0.01	0.94
1.282	0.078	0.02	0.94
1.277	0.385	0.00	0.84
	Regression coefficients -78.712 0.243 1.864 1.282	Regression beta weights  -78.712 0.243 1.864 1.282 0.078	Regression coefficients         Beta weights         Level of sig.           -78.712         0.00           0.243         0.837         0.00           1.864         0.132         0.01           1.282         0.078         0.02

 $R^2 = 0.95$  Standard error of estimate = 6.42 Value of coefficient of determination if that variable was deleted from the analysis.

Table AS. Coefficient of determination between percent carcass retail yield and a combination of carcass measurements, scores and percent flank yield and the regression coefficient for each of the variables (combined weight groups).

	Regression	Beta	Level of	R <sup>21</sup>
Independent variables	coefficients	weights	sig.	deletes
Constant	26.430		0.00	
Carcass wt.	027	548	0.00	0.78
Round score	0.430	0.144	0.03	0.84
Length of round	1.006	0.426	0.00	0.81
L. dorsi muscle area	0.389	0.142	0.02	0.84
% flank yield	0.410	0.737	0.00	0.46

 $R^2 = 0.85$  Standard error of estimate = 1.84 Value of coefficient of determination if that variable was deleted from the analysis. Table AT. Coefficient of determination between weight of carcass retail yield and a combination of objective carcass measurements and percent flank fat trim, and the regression coefficient for each of the variables (500 to 550 lb. weight grown).

each of the	variables (500 to 5	ou lb. weig	ght group).	
	Regression	Beta	Level of	R21
Independent variables	coefficients	weights	sig.	deletes
Constant	<b>-</b> 78 <b>.14</b> 6		0.13	
Carcass wt.	0.323	0.388	0.00	0.73
Length of round	4.059	0.358	0.00	0.76
L. dorsi muscle area	2.360	0.230	0.00	0.80
% flank fat trim	-1.050	773	0.00	0.49

 $R^2 = 0.84$  Standard error of estimate = 5.82

Table AU. Coefficient of determination between percent carcass retail yield and a combination of objective carcass measurements and percent flank fat trim, and the regression coefficient for each of the variables (500 to 550 lb. weight group).

Independent variables	Regression coefficients	Beta weights	Level of sig.	R <sup>21</sup> deletes
Constant	34.099		0.07	
Carcass wt.	0.006	0.019	0.77	0.88
Length of round	1.401	0.316	0.00	0.82
L. dorsi muscle area	1.010	0.251	0.00	0.82
% flank fat trim	379	713	0.00	0.58

 $R^2 = 0.88$  Standard error of estimate = 2.00

<sup>1</sup> Value of coefficient of determination if that variable was deleted from the analysis.

<sup>&</sup>lt;sup>1</sup>Value of coefficient of determination if that variable was deleted from the analysis.

Table AV. Coefficient of determination between weight of carcass retail yield and a combination of carcass weight and percent flank yield, and the regression coefficient for each variable (700

to 750 lb. weight group). Regression Beta Level of Independent variables coefficients deletes weights sig. 0.03 Constant -85.807 Carcass wt. 0.320 0.465 0.00 0.58 0.22 1.879 0.758 0.00 % flank yield


 $R^2 = 0.79$  Standard error of estimate = 6.28 lValue of coefficient of determination if that variable was deleted from the analysis.

Table AW. Coefficient of determination between percent carcass retail yield and a combination of carcass weight, percent flank yield and percent kidney knob, and the regression coefficient for each of the variables (700 to 750 lb. weight group).

	Regression	Bet <b>a</b>	Level of	R <sup>21</sup>
Independent variables	coefficients	weights	sig.	deletes
Constant	47.476		0.00	
Carcass wt.	0002	001	0.94	0.74
% flank yield	0.436	0.774	0.00	0.20
% kidney knob-right side	677	213	0.02	0.70

 $R^2 = 0.74$  Standard error of estimate = 1.60 <sup>1</sup>Value of coefficient of determination if that variable was deleted from

the analysis.



JECTIVE SCORES  Maturity Marbling Final Carcass	Score		60	12	14	14	15		-	i -	0 0	80 <b>•</b>		11.7	11.7		17	20	21	18	16	15	13	15	21	11	16.7	16,7		18	14	14	20	15	17	18	22	16	27
A CARCASS EVALUATION - SUBJECTIVE Conf. Hind- Conf. Fore- Maturit	Quarter		90	90	60	90	08	60	70	יי ס	3 8	2,9	၅	6 <u>.</u> 3	6.3		07	80	80	60	80	90	05	03	07	05	6.4	<b>6.4</b>		07	. 80	80	60	80	90	07	90	07	•
A CARCASS EV. Conf. Hind-	Quarter		90	90	80	80	60	1	50	S &	88	<b>†</b> 5	6 	7.0	7.0		07	10	66	80	07	05	90	40	60	05	7.0	7.0		08	80	60	: 11	60	07	80	60	60	•
APPENDIX I	Conf.		90	90	60	07	60	10	5.0	ر د	3 8	<b>5</b> 6	70	8°9	8 <b>.</b> 9		07	10	80	08	07	70	05	90	80	05	9.9	9*9		07	. 80	60	10	60	07	80	60	60	
Round			90	90	80	80	60	: =	90	90	8 8	<b>5</b> 6	[6]	7.1	7.1		90	11	80	08	02	05	90	05	80	05	7.1	7.1			90	08	11	10	0.2	08	10	10	
Steer	No. Breed	26-	н 1	2 3	3 1	4 2	5 1	9		~ a	7 (	7 0	10 2	Eng. Mean	Metric Mean	Group II .5175"	11 1	12 1	13 1	14 1	15 1	16 1	17 3	18 3	19 1	20 2	Eng. Mean	Metric Mean	Group III .76-1.00"	21	22 2	23 1	24 1	25 1	26 1	27 1	28 1	29 1	!

2040	6	0.00011	APPENDIX	APPENDIX I A (Continued)	1)	Mouth 1 4mg	Ting October	
No. Breed	Score	Conf.	Ouarter	Ouarter	Score	Score	rinar varcass Grade	
1.0								l
31	60	60	10	80	ო	18	08	
32 1	10	10	11	60	ന	20	60	
33 1	60	80	80	07	ო	21	60	
34 1	90	07	90	80	2	24	60	
35 2	90	07	90	07	7	16	07	
36 1	07	07	80	07	ო	17	07	
37 1	07	90	07	80	4	16	08	
38 2	80	80	80	80	7	13	90	
39	10	60	10	80	ന	21	60	
7 07	07	80	80	080	m	17	08	
Mean	6		\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	2,5	2.8	18.3	8.0	
Metric Mean	7.9	8,1	8.2	7.8	2.8	18,3	8.0	
Group V .2650"								
41	03	03	90	03	7	14	90	
42 2	<b>9</b> 2	07	07	07	5	20	80	
43 1	07	07	07	90	∞	20	07	
44	07	90	07	05	4	18	90	
45 2	80	07	07	80	က	15	07	
	80	90	80	10	ന	21	60	
	07	07	07	90	4	15	90	
48 2	07	90	07	90	Ŋ	13	05	
49 1	60	80	80	80	4	20	80	
50 2	07	07	07	8	41	16	[]	
Mean	7.0	9 <b>.</b> 9	6.9	6.7	4.7	17.2	6.9	
Metric Mean	7.0	9*9	6*9	6.7	4.7	17.2	6.9	
Group VI .5175"								
51 2	80	80	80	07	4	20	80	
52 1	07	90	07	05	Ŋ	17	07	
53 1	90	80	80	80	4	19	80	
54 2	80	60	60	80	ო	17	08	
	<b>2</b> 0	07	80	60	4	13	90	
	90	80	80	80	4	70	08	
57 2	02	05	90	05	4	17	90	
58 2	80	80	80	80	4	16	07	
	60	60	60	10	ო •	73	60	
	ଞ୍ଚ	70	ଞ୍ଚ	9	<b>4</b> 1	91	zi i	
Eng. Mean	<b>7.</b> 6	7.5	7.8	7.2	თ ი ო ი	17.8	<b>7.</b> 4	
Metric Mean	٧•٥	ر•/	/•۵	7./	5.7	0 • / τ	<b>†</b>	

					V	(Continued)			
teer		Round	Cverall	Conf. Lind-	Conf. Fore-	Maturity	Narbling	Final Carcass	
N.O.	Breed	Score	Conf.	Quarter	Quarter	Score	Score	Grade	
VII	76-1.00								
19	2	20	<i>L</i> 0	20	08	7	16	20	
23	H	. 5\ O	10	10	10	$\mathcal{N}$	16	20	
63	2	20	20	20	<b>න</b> ට		77	90	
\. \.\!\	m	90	07	07	07	2	1 5	20	
· Tu	· ~	20	90	07	90	m	, L	90	
99	Н	11	10	ī	60	m	25	12	
23	(~	S	30	60	ထင	, M	16	20	
99	2	08	90	80	20	7	20	90	
69	0	60	08	30	20	Μ	57	60	
20	2	08	07	20	20	7	18	20	
Mean		₽ 1	2.3	្រះ	7.7	W V	ಲ್ಲಿ ಬಿಟ್ಟ	<del>7.</del> 6	
Metric Mean	<b>.</b>	a, 1•	•	ಗ <b>ಿ</b> ಐ	7.7	س بر	18,3	7•6	
VIII	1.01-1.2	10							
71	2		30 •	20	08	7	20	90	
72	<∨	90	20	90	20	ſΛ	77	90	
73	2	30	20	90	20	m	15	20	
77	2	08	08	30	20	7	20	වල	
2	2	20	20	20	90	7	20	90	
92	Н	ဆင	80	30	90	7	15	90	
17	2	80	90	60	20	7	16	20	
78	2	90	<u> </u>	ခင	60	<b>_</b> 7	22	60	
6/	7	C2	20	90	20	7	<b>1</b> 6	20	
റ്റ	႕	0 80	60	တ	10	7	56	10	
Mean		7.8	7.7	7.7	<u>7.</u> 6	C•†	7.31	2.0	
Metric Mean	_	J. E	7.7	7.7	<b>5.</b> 6	0•17	18.4	7.6	

1	1	-192-	
Eth Thor.	66.23.33.00.03.33.00.00	1 22 23 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	44444444444444444444444444444444444444
th Thore		: 425884274754	27 17 17 17 17 17 17 17 17 17 17 17 17 17
T YELSTEES Eth There		2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 3. 3. 3. 3. 3. 3. 3. 3. 4. 4. 4. 4. 4. 4. 4. 4. 4. 4. 4. 4. 4.	2444444444 2444444444 2444444444444444
CETRITUE E.T. Eth Thor.	00 17 00 17	######################################	15 15 15 15 15 15 15 15 15 15 15 15 15 1
Thor	194899999999999999999999999999999999999	11 12 12 12 12 12 12 12 12 12 12 12 12 1	25.67 10.667 10.667 10.667 10.667 10.667
67 H		20 20 20 20 20 20 20 20 20 20 20 20 20 2	7000 11000 11000 1000 1000 1000 1000 10
H	170555555	2	4 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
HE :	10 00 00 00 00 00 00 00 00 00 00 00 00 0		
Carc.		1	-0. 10. 10. 10. 10. 10. 10. 10. 10. 10. 1
Steer	Group 1.26.1 2.22 3.44 7.66 Mean 100	Setric Rean Urcur II •51 12 13 13 14 15 16 17 18 19 20 Fag. Mean Fetric Kean	Group III • 22 22 24 25 26 27 28 29 30 30 30 Xean

	e e e e e e		
		•	
		·	

	1 1		2,50	
	Sth Thor. Avg.	21.00 14.67 17.00 17.33 17.00 17.00 17.00 17.00 18.1	12.67 13.00 13.00 10.00 10.33 11.67	11.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 12.00 13.00 13.00 14.00
	Eth Thor. 12 in.	7. 17. 17. 17. 17. 17. 17. 17. 17. 17. 1	12 12 12 12 12 12 12 12 12 12 13 13 14 15 15 15 15 15 15 15 15 15 15 15 15 15	1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0
	Eth Thor. 8 in.	25 27 27 27 27 20 20 20 20 20 20 20 20 20 20 20 20 20	244224441.44 2444441.44 24	12 52 52 52 52 52 52 52 52 52 52 52 52 52
(p)	Éth Thor. 4 in.	18 13 13 19 19 19 17 18 10	8.3447 8.3447 8.3447 8.3447 8.4447 8.	52122222151 112222151 24222151
B (Continue	Sth Thor. 5th Thor. Eth Thor. 12 in. Avg. 4 in.	25.33 20.00 20.33 20.00 20.00 20.00 20.00 20.00	18.67 19.33 19.67 11.00 11.00 16.00 15.4	18.67 21.00 18.00 18.00 13.33 13.33 13.03 16.3
L XICTERS	5th Thor. 12 ir.	25 17 20 27 27 27 28 27 27 27 27 27 27 27 27 27 27 27 27 27	17 17 17 17 17 17 17 18 17 18 17 18 17 18 17 18 18 18 18 18 19 19 19 19 19 19 19 19 19 19 19 19 19	17 17 13 13 13 15 15 15 15
	5th Thor. β in.	38 30 25 25 23 23 23 24 28 29 29 29 20 20 20 20 20 20 20 20 20 20 20 20 20	23 17 17 17 17 17 17 14	28 5 1 2 2 1 2 2 8 8 2 2 2 2 3 4 2 2 3 4 2 2 3 4 2 3 4 2 3 4 2 3 4 3 4
	5th Thor. 4 in.	77 77 77 77 77 77 77 77 77 77 77 77	11 09 09 08 08 09 11 11 140	11 10 12 12 15 10 10 10 10 10 10 10 10 10 10 10 10 10
	Carc.	75-27 75-27	.50" 740 717 702 703 704 693 708 708 321	737 723 723 736 736 736 726 726 328 328 328
	Steer No.	Croup IV 1.01 32 32 33 34 35 36 37 38 39 40 40 Metric Mean	Group V .26 41 42 43 44 45 46 47 48 49 50 Eng. Mean	Group VI •51- 52- 53- 54- 55- 56- 57- 58- 59- 60- 3ng. Mean Metric Mean

		-194-
gth Thor. Avg.	17.33 11.67 11.67 15.33 23.67 23.67 17.67 17.67	19.67 19.33 17.67 17.67 12.67 16.00 18.00
Sth Thor. 12 in.	25 118 128 128 148 170 170	22 28 2 29 29 29 29 29 29 29 29 29 29 29 29 2
8th Thor. 8 in.	23 12 12 12 12 12 12 12 12 12 13 14 15 15 16 17 18 18 18 18 18 18 18 18 18 18 18 18 18	80 10 10 10 10 10 10 10 10 10 10 10 10 10
ed) Eth Thor. 4 in.	44 44 45 45 45 45 45 45 45	
APranlix I B (Continued) h Thor. 5th Thor. St 2 in. Avg.	17.33 15.00 22.67 26.00 17.33 12.00 24.67 21.33	21.00 21.33 21.33 21.67 21.67 21.00 20.23
APPTNJIX 5th Thor- 12 in.	55 55 57 58 57 58 57 58 58 58 58 58 58 58 58 58 58 58 58 58	.t. 6623382738678
5th Thor. 8 in.	21.50 33.25.	25 27 28 29 20 20 20 20 20 20 20 20 20 20 20 20 20
5th Thor. 4 in.	24442200444 24442200444 344 344	
O	716 656 656 712 712 722 723 323.8	1.01-1.25# 694 736 738 718 702 758 700 728 728 728
Steer No.	Group VII -76 62 63 64 65 65 66 67 69 70 Eng. Mean Metric Mean	Oroup VIII 1, 72 72 73 74 75 76 77 78 79 80 Eng. Mean

		•																									
	4th Lumb.	20	88 6	0 0	တ ဝ	16	1 5	5 전	•41 10•5		בומ	20	H 0	12	01 11	80	0 10 10 10 10	10.6	(	1.7 7.1	07 1/1	18	16	35 15	2 t 5 t 5 t	77.	?• 寸 <del>1</del>
	1st Lumb.	00.60	18,33	10.33	14.33	16.00	12.33	00 K	• 49 12• 5		17.00	17.33	16.67	14.57	14.67	15.67	14-33	15.5		70°00 20°00	10.00	23.67	13.00	24.67	00°02	-72 -1 3 F	• ○ 1
	lst Lumb. 12 in.	90	<b>1</b> 0	07	7 <b>1</b>	21	2,5	당시	.51 12.9		) T c	H F	57.	ን'Ά'	15 21	12,	$\mathbb{Z}_{\mathcal{C}_{\mathcal{C}}}$	14.2	(	Lc 27	12	23	ь Г	28	97 18	02.	<b>5.02</b>
(T)	lst Lund. 8 in.	1 7	20 20 20 20	18	17	17	<u>ي</u> د	12 E	.68 17.4		12	30	24 26	) 61 11	დ ა ი	333	ည်	27.2	ì	575	12 28	£ £ £	7.5 T &	26 33	22	ည မျှင် • (	6.22
B (Cortinue	11th Thor. 1st numb. Avg. 4 in.	20	10	000	SO O	10	00 C	S 21	7.5		77	60	<del>ಟ</del> 5	30	010	11;	3F	11.2	,	15	00 د	14.	75	20	121	-  -  -  -  -  -  -	C• 7T
T SECTED TO	llth Thor. Avg.	05.33	11.33	08.67	10.00	11.33	50 50 50 50 50	11.33	ω φ Μ • • ον		11,00	10.00	10.00 10.67	29.50	12.67 11.00	12.00	8	11.1		16.00 16.00	10.00	12.67	10.0 13.67	22.33 55.05	14.33	9, -	7.77
	llth Thor. 12 in.	1 ~	27 C	60	다 80	11	75 ر	ਰ ਨ <u>ੀ</u>	10.0		10	0 F	010	3 0 0	11	12	∄F	10.5	!	17 16	45	10	07 13	, T	77	9 <del>.</del> -	7.47
	llth ihor. 8 in.	10	11	<b>S</b> O	0.0 0.8	:11	건 건 건 건	721	 		13 61 61	11	디논	, A,	97 11	17,	12 18	12.2		L7 18	60 F	1 H	17 17 17	22	1 K	1	14.0
	llth Thor.	.50" 08	1 8	000	<u>ن</u> 0	12	ግን <sup>ይ</sup>	8 8	9.1	75"		7 [	50 C	71	11 (	£ £ £	와	10.6	.16-1-67.	ָּהְרְ הַלְּהְ	9 r	12	90 L	53 1	큐	٠.٠ ٢٠٧٠	14.0
	Seer Meer	3roup <u>1 .26</u>	0 m	<b>)-</b> :1	᠘᠘ᠺ	2	လ ၁		Trg. Mean Metric Mean	Group II .51	<b>ار</b> ر	13	<b>1</b> 1	16	77 T	) () ()   H (	700 700 700 700	Metric Mean	II	<b>1</b> 2	87	1 00 1 170	900	- 5 - 5 - 5 - 6 - 6 - 6 - 7 - 7 - 7 - 7 - 7 - 7 - 7 - 7 - 7 - 7	% 0% % 0%	Trg. Fean	Tetric Mean

	4th Lumb. 4 in.	цмц- хача	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	20 20 10 10 10 10 10	24 99 11 18 10 10	17 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	00000000000000000000000000000000000000
	lst Lumb. Avg.	31.00 24.00 20.00 33.67	32.00 21.00 28.00	22.33 20.67 .97 24.7	16.67 14.00 14.00 20.33	20.00 13.33 22.00 11.00 17.0	24.67 11.67 24.33 12.63 24.33 26.67 20.67 20.67
	lst Lumb. 12 in.	35 127 27 28	27 22 62	233 200 25.2 25.1	70 70 70 70 70 70 70 70 70 70 70 70 70 7	23 11 24 19 20 20 20	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0
a)	ist Lumb. 8 in.	77 31 20 20 20 20 20 20 20 20 20 20 20 20 20	1402°	27 28 33.33	12 14 14 28 28 16	25 25 25 14 14 21 21 21	32 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
B (Continue	llth Thor. 1st Lume. 1 Avg. 4 in.	15 11 11 12	1809F	17 11 16.8 15.8	00 00 00 00 00 00 00 00	07 07 09 09 38 38	00 00 00 00 00 00 00 00 00 00 00 00 00
A FPW OIX I	llth Thor. Avg.	16.67 24.33 15.67	77777 77777 77777 77777	15.33 17.00 18.2	11.00 09.67 10.33 11.67	12.67	13.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00 10.00
	llth Thor. 12 in.	11 23 31 7	7 M M F C	11.15 10.05	#8####	12 12 10 12 12 12	40027727474 20027727474
	llth Thor. 8 in.	0 m 0 a	2227	15 20 •76 19•3	000000000000000000000000000000000000000	13	21 00 00 00 00 01 01 01 01 01 01 01 01 01
	11th Ther.		0 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	12 12 18-73 16-55	.50" 09 08 09 10		51-75 13 10 07 09 16 10 12 12 13 145
	teer	Group IV 1.0	1 77 (O) (C) (C) (C) (C) (C) (C) (C) (C) (C) (C	39 39 40 3ng. Mean Metric Mean	roup V .26. 11. 12. 13. 14. 14. 15.		Group VI .5 52 52 53 53 55 56 57 58 58 59 60 Eng. Wean

1 1		25,
4th Iumb. 4 in.	74488338844 6-21 7-41	238888888888 60
lst humb. Avg.	26.67 20.09 20.09 20.09 20.09 20.09 20.09 20.09 20.09	
lst Lumb. 12 in	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	
1) lst Lun <b>b.</b> { in.	70.00 70.00	25 20 20 20 20 20 20 20 20 20 20 20 20 20
APPTOIA I B (Continued lith Thor. 1st Lumb. Avg. 4 in.	45444468844 <u>6</u>	
APPTOIA I lith Thor. Avg.	22 23 23 23 23 24 24 25 25 25 25 25 25 25 25 25 25 25 25 25	
llth Thor. 12 in.	244488249 2440 2440 2440 2440 2440 2440 2440 2	89558884988 000 000 000 000 000 000 000 000 000
lith Thor. 8 in.	2248 248 254 254 254 254 254 254 254 254 254 254	80777500 807775700 807775700 8077757
llth ihor. 4 in.		-125 -125 -11-
Steer Fo.	Group VII • 78 62 63 64 65 65 65 66 69 70 Ing• Yean Wetric Yean	Group VIII 1, 72, 72, 73, 74, 75, 76, 77, 77, 78, 79, 80, 80, 80, 80, 80, 80, 80, 80, 80, 80

	3rd Sac. 12 in.	2.00 2.00 2.00 2.00 2.00 2.00 2.00 2.00	000 000 000 000 000 000 000 000 000 00	स्थाद कर विकास कर विकास १
	3rd Sac. 8 in.	7.50 00 00 00 00 00 00 00 00 00 00 00 00 0		80000000000000000000000000000000000000
	3rd Sac. 4 in.	00 00 00 00 00 00 00 00 00 00 00		1159 1188 128 128 128 128 128 128 128 128 12
	€th iumb. ±Vg.	08.67 12.00 08.67 07.67 06.33 16.33 16.33 16.33	23.67 23.67 23.67 23.33 24.88 8.43 8.43 8.43 8.43 8.43 8.43 8.43	22.67 21.67 09.67 15.00 17.00 18.33 18.33
inued)	6th Lumb. 12 in.	1. 1. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.	20 12 10 13 13 15 15 15 15 15	22 22 22 24 24 25 27 27 27 27
(Gort	6th Lumb. 8 in.	00000000000000000000000000000000000000	22 33 12 12 15 15 15 16 16 16 16 16 16	28 128 24 25 24 25 25 25 25 25 25 25 25 25 25 25 25 25
APPENUT	óth Lumb. 4 in.	00000000000000000000000000000000000000	13 00 00 00 00 00 00	3.00 177 177 177 177 177 177 177 177 177 1
	dtli Lumb. Rvg.	112.00 112.00 112.00 100.00 100.00 13.00 13.00 13.00 13.00 13.00 13.00 13.00 13.00 13.00 13.00 13.00 13.00 13.00 13.00 13.00 13.00 14.00 15.00 16.00 1	20.00 16.67 11.00 11.5.67 15.67 15.67	25.00 25.00 25.00 25.00 20.00 23.33 20.00
	lth Lumb. 12 in.	25 24 24 24 24 24 24 24 33 33 34 34 34 34 34 34 34 34 34 34 34	2000 1000 1000 1000 1000 1000 1000 1000	30 10 10 10 10 10 10 10 10 10 10 10 10 10
	ith Lumb. δin.		75 80 80 10 10 10 10 10 10 10 10 10 10 10 10 10	76-1.00" 22 28 23 27 26 26 27 26 23 23
	teer No.	Group 1 .29- 2 2 3 4 4 5 5 6 7 7 7 8 10 10 10 10 Metric Kean	Froup II .51 12 12 13 14 15 16 17 18 20 7ng. Mean	Group III

			-199-	
	3rd Sac. 12 in.	1.50 1.20 1.20 1.20 1.20 1.20 1.20 1.20 1.2	170 170 170 170 170 170 170 170 170 170	8767000400 •~
	3rd Sac. E in.	25 25 25 25 25 25 25 25 25 25 25 25 25 2	9.1000000000000000000000000000000000000	000 000 000 000 000 000 000 000 000 00
	3rd Sac. 4 in.	22 23 12 12 13 14 15 15 15 15 15 15 15 15 15 15 15 15 15	10 11 12 12 13 14 14 15 16 17 17	200 100 100 100 100 100 100 100 100 100
	éth Lumb. ∴vg.	31.00 15.67 17.00 23.00 21.67 21.67 17.00 19.67	16.67 11.33 11.33 11.33 11.33 11.33	22.67 22.67 23.67 23.67 23.67 23.67
inued)	6th Lumb. 12 in.	25.5 20.5 20.5 20.5 20.5 20.5 20.5 20.5	26 129 129 124 129 23 23 24 25 25 27	
APPEDIX I B (Continued)	6th Lumb. 8 in.	100 50 50 50 50 50 50 50 50 50 50 50 50 5	22 23 24 24 24 25 26 27 26 27	8 4 2 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
TO PPE UT	6tn Lumb. 4 in.	20 11 12 12 12 12 12 15 15 15	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	200 000 100 100 100 100 100 100 100 100
	4th Lumb. Avg.	24.67 29.00 17.00 22.67 25.00 19.67 23.3	16.67 13.33 13.33 15.33 15.33 15.33 15.33	13.33 10.00 11.00 124.67 12.00 18.33 18.33 18.33
	Lth Lumb. 12 in.	201120 20	122 122 123 144 164 164 164 164 164 164 164 164 164	00 00 00 00 00 00 00 00 00 00 00 00 00
		01-1.25 34 32 28 34 34 26 26 1-12 26 1-12 26 26	5.50" 18 15 15 12 20 21 21 11:	5175" 21 24 25 25 25 26 26 27 20 20 20 20 20 20 20 20 20 20 20 20 20
	steer No.	Group IV 1.0 31 32 33 34 35 36 37 40 Mean Metric Mean	Group V .26.  11  12  13  143  144  145  148  149  50  Fng. Kean Wetric Mean	Group VI

	× .				
		4	e e e	. , ,	

	3rd Sac.	12 in.		13	11	CT	L N	07	27 27	25	20	50	임	r:	13.0		??	임	19	디	60	12	ા	Ħ	10	្យ	12.7
	3rd Sac.	ರಿ ಸ್ಟಾ		53	တ	31	25	.တ ဝ	23	17	19	25	)Te	C.L.	17.5		50	23	2,5 2,5	20	77	17	77	20	17	<u>1</u> 5	17.3
	3rd Sac.	4 in.		82	ኒ스 러	50	58	بار ا	16	56	16	25	27	52.	20.1		27	23	21	೮	22	23	11	16	22	25 1	./. 19.8
	oth Lumb.	AVG		20,33	16.67	23.00	25,00	16.67	(n)	25.00	22.33	26,00	27.00	£.	23.63		19.33	22.00	26.67	26.00	24.00	29.33	13.33	20.33	25.33	15.33	22.56
nued)	1 23	12 in.		92	77	32	on TU	) (1	36	31	56	ري کر	[2]	95-	24.3		ა ა	12	23	33	23	38	۲٦ ۲۷	19	8	기 기 기	24.9
I B (Conti	Ι'	8 in.		18	50	21	H (C)	200	32	52	22	30	37	1,13	56.9		<b>ા</b>	22	31	56	27	댔	<b>1</b> 6	23 3	27	원	23.7
APPINITE A	6th Lumb.	1.10.		17	12	50	75	יו	17	23	19	23	23	<u>.7</u> 0	19.7		20	<i>رب</i>	56	19	22	19	50	77	19	20 10 10 10 10 10 10 10 10 10 10 10 10 10	• (> 19•1
	4th Lumb.	AV3		H	,	ं	ं	ထ	m	ં	15.67	ď	αĵ	05•	55.9		27.33	_ <del>_</del>	ġ	ထံ	5	ä	9	\$	ň	_= <u></u>	27° 27°
	Lumb.	12 in.		20	02	77	23 72	21	50.	29	13	25 25	24	ထ မာ •	7.23		23	53	23	31	36	25	16	22	27	$\sim 1^{\circ}$	25°5
	Lith Lumb.	8 in.	6-1.00"	96	200	19	C	16	띥	Č	15	21	7:5	C	27.0	01-1.25"	32	31	9	32	덗	23	21	18	23	ದ್ಟ	7 - 00 C
	Steer 4	• O 🔏	Group VII .7	<b>1</b> 9	62	63	9	Ø.	<b>6</b> 6	29	89	69	20	∄ng. I	etric Mean	II	7.1	72	73	7/7	75	76	77	78	62	ప్ ;	ing. Mean Metric Mean
			C												24	(;)										ŗ	#1 <b>}</b>

\				
	Fat Taick.	8.32 8.32 8.32 8.32	122 132 134 134 134 137 137 137 137 137 137 137 137 137 137	24444444444444444444444444444444444444
	Fat Thick B	06 12 07 04 06 09 03 03 03 03 03	9. 9. 71 72 73 74 75 76 76 76 76 76 76 76 76 76 76 76 76 76	84 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
inued)	rat Thick. A	% 2 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	1172 120 120 120 120 120 120 120 120 120 12	2. 1222 2. 122
I B (Cont	5th Sac. 5th Sac. fat 12 in. Avg.	04.67 21.00 04.33 06.67 05.67 06.67 06.67	09.33 11.33 10.33 13.00 12.13	11.00 11.00 11.00 11.00 10.00
APPINIDIA	5th Sac. 12 in.	3.1000th 000333000	00000000000000000000000000000000000000	00 00 00 00 00 00 00 00 00 00 00 00 00
	jth Sac. βin.	00000000000000000000000000000000000000	6.33 1.02 1.03 1.03 1.03 1.03 1.03 1.03 1.03 1.03	## NE 66 L L 19 0 0 L 19 0 L
	5th Sac. 4 in.	11.002.002.003.11.00.002.11.00.000.0	16 26 12 17 19 19 17 17 17 17	12 23 24 25 25 25 25 25 25 25 25 25 25 25 25 25
	3rd Sac. Avg.	65.00 64.33 64.33 64.33 65	75" 11.67 06.00 09.00 09.00 09.33 17.00 11.67 11.00	6-1, CO. 12, 23, 23, 23, 24, 25, 24, 25, 24, 25, 25, 25, 25, 25, 25, 25, 25, 25, 25
	Steer No.	Group 1 .20 2 2 3 4 4 5 6 7 8 9 10 3ng. Mean Metric Mean	Group II •51 12 12 13 14 15 16 17 18 19 20 Eng. Wean	Group III .76 22 23 24 25 26 27 27 27 29 29 30 Eng. Mean

	\												
Fat Thick.	0 7	0087	† 0, 0, 1, 0, 0,	5 <b>2</b> 5	12 17 17	ಎ • ರ • ರ	하 9 0 c c c c c c c c c c c c c c c c c c	4 0 0 0 8 9 9 9	0 0 H • 0 0 0 G • 0	13 06	89 H Z F H	(0.77 <i>)</i>	21. 21.
Fat Thick.	m ć	233	2 0 0 2 0 0	to	36	34.7	ខ្ពង់ង	3 G S Y	25 20 15.8 15.8	776	X <b>A</b> A &	. 612 127	17.5
nued) Fat. Thick.	30	37 77 77	) (7 F	V) 1, (	5 <u>7</u>	6.4.3 6.4.3	177 00 08 08	C H H C	10.6	88	% % & % A	경우다	5년 5년 5년
APPA JIM I B (Continued)	ėvg.	7000 7000 71000 71000	21,00 21,00 12,67	12.67	18.00 14.67 63	15.90	50.00 00.67 00.01 10.67	20 <u>.</u> 67 21 <u>.</u> 00 12.67		10.33 11.00	15.00 14.67 15.33	12.60 12.60 12.60 13.60	17.67 • (3 16.10
Sth Sec.	12 in.	SH 증 F	ឯដទ	08 11 08 11	이라. - 라마	10.0	70 70 70 70	ม	000 000 000 000 000 000 000 000 000 00	10	Ø Ø ॐ ⊅ O O O O	3381 1	
l l	R in.	7 E C C	31 71 71	158	급리	13.8	21 00 07	мω ω г мω ю г	1887 24	44	00 00 00 00 00 00 00 00 00 00 00 00 00	CT 821	17.56
ω, <sub>(2)</sub>	्यः १	7 m/o 6	y H 公 n の に	0 (~ © 1 -1 (V	0 8 C	トーゴ	7795	0, W M F	184126 0.	10 24	25 25 15 15 15	\d # #.	2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 -
1	P- K∕	7440 767	• •		• • 1	16.6	•	12.00	12.00	7.40 8.00	07.07 11.06 17.06 17.09	$t \odot C = t$	$\alpha$ ta $\alpha$
Steer	Group IV 1.	7 W M 7	፣† <b>ህ</b> ነላር ናነ ወን መ	) M M	۵ اندر ۲۰ نځ	Retric Mean	Szeup V .26- 41 42 43 44	1-1-7-	19 19 50 7ng• Meen Metric Mean	Group VI .51	ጠ. ታኒኒዣ ኒኒኒኒኒ	/ カ/ カ/ カ/ / ケーの ウ/	60 Ing. Men Tetric Year

1 12-	APPINIX I B (Continued)  3rd Sac. 5th Sac. 5th Sac. 5th Sac. 5th Sac. but Thick. Fat Thick.  A.V. I. in. 8 in. 12 in. A.V.		15,33 35 25 11 23,67 33 14	23 15 13 05 11.00 22 25	.00 29 16 10 18.33 18 24	•00 23 22 10 16•33 17 35	67 18 06 06 10,00 31 12	3 16 15 18.67 22 30	•67 35 11 19 21•67 37 20	•00 23 14 04 13•67 24 22	•00 34 18 13 21•67 32 22	13 17.67 23 30	7 1.03 .62 .42 .65 1.02 .52	6.1 15.7 10.6 17.47 25.9 23.L	.01-1.25"	0 28 12 15 18,33 37 34	3.67 24 15 15 18.00 40 28	.•67 22 20 16 19•33 45 32	5,33 19 16 11 15,33 34 36	5.00 3½ 14 11 19.67 ¼0 26	.•00 38 25 12 25•00 27 33	<b>2</b> 5 <b>1</b> 2 07 <b>1</b> 4.67 20 36	33 17 11 08 12.00 29 30	.33 39 27 13 26.33 32 26	21 18 24•67 25 30 23	
88 Mean 111. 11. 12. 13. 14. 15. 15. 15. 15. 15. 15. 15. 15. 15. 15	1 2	17	15.3	•	16.00	22,00	<b>29°5</b> 0	21.33	27.67	14.00	•	9	-67	Н	-1.2	•	9	•	3	15.00	17.00	9	12,33	16.33	1,5,00	

I B (Continued)	average A, B, C	09.33 12.67 11.67 10.33 11.33 07.00 13.00 10.67 10.67	16.33 16.67 16.67 16.67 16.33 16.33	22.67 18.67 18.67 22.33 20.00 21.00 23.67 20.67
APPENDIX I B (C	Fat Thick. G	25. 11. 25. 25. 26. 26. 26. 27. 27. 27. 27. 27. 27. 27. 27. 27. 27	98 110 12 13 13 14 15 15 15 15 15 15 15 15 15 15 15 15 15	854445683448 854445683448
14 14	Fat Thick. F	00000000000000000000000000000000000000	09 1331 1990 1990 1990 1990 1990	235 235 235 235 245 25 25 25 25 25 25 25 25 25 25 25 25 25
	Fat Thick. E	00000000000000000000000000000000000000	13 10 10 10 10 13 13 13	103884444 1038844444444444444444444444444444444444
	Fat Thick. D	.55." 02 10 09 08 06 08 08 08 08	75" 11 12 12 12 12 12.9	.76-1-00" 111 199 199 199 199 199 199 199 199 199
	Steer No.	Sroup 1 .26- 2	Group II .51- 12 13 14 15 16 17 18 19 20 Erg. Nean	Group III -70 22 22 23 24 25 26 27 27 28 27 28 27 28 27 27 27 28 27 27 27 27 27 27 27 27 27 27 27 27 27

(pg	nge C	0.000m00000		67 67 67 67 67
Continue	Average A, B, C	24 24 24 24 24 24 24 24 24 24 24 24 24 2	27 20 20 20 20 20 20 20 20 20 20 20 20 20	15.67 16.67 16.67 16.67 16.67
APPENDIM I B (Continued)	Fat Thick. G	20000000000000000000000000000000000000		22 22 22 24 24 25 24 25 25 25 25 25 25 25 25 25 25 25 25 25
15. T.	Fat Thick. F	5.4 5.4 5.4 5.4 5.4 5.4 5.4 5.4 5.4 5.4	951 198 198 198 198 198 198 198 198 198 19	00000000000000000000000000000000000000
	Fat Thick. E	4974888888999°°°°°°°°°°°°°°°°°°°°°°°°°°°	2422448492 <b>9</b>	28888888888888888888888888888888888888
	Fat Thick. D	다. 아. 아. 아. 아		2000 2000 2000 1000 1000 1000
	Steer No.	Group 17 1.0 31 32 33 34 34 35 36 36 59 50 Eng. Mean	Group V •26.	Sroup VI -51 52 52 53 53 55 56 59 60 Eng. Mean Yetric Yean

ontinued)	Averase A B C	C6U 6A	20		22.67	19.67	22,33	19,33	25,33	0.0 0.0 0.0	(C+1)	して・ハー	23.67	23,33	23.	22.0		30,33	25,33	33,00	25.67	26,00	25.67	25.67	26.33	25.67	26.00	1. 1.	25. 0
APPEDIX I B (Continued)	Fat Thick.	5	00	7.0	77.2	31	25	19		, - - - -	ع <u>ب</u>	<u></u>	31	53	1.06	27.00		56	33	26	27	24	56	17	17	29	쏬	35.	25.0
APP	Fat Thick.	4	ኒ (•	J ,	ĭì	56	18	30	C	) (d	) L	) <del> </del> (	20	<b>2</b> 5	1.02	25.8		13	36	22	23	16	15	16	17	22	25	59•	22.7
	Fat Thick.	1	00	77	7	16	21	17	· C	- t	) L	^ · <b>F</b>	ري <b>ا</b>	56	82.	22.5		20	30	27	25	20	16	76	27	25	27	ر ا ا	21.8
	Fat Thick.	-	: ;;;;; =	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	15	23	16	19	0	) _ -	) t	- ì	15	17	-74	16.9	1.01-1.25"	19	. 23	18	21	23	16	77	17	22	미	94.	13.7
		1.0	roup vir • (c	₹.	62	63	79	65	, 99	62	-0	000	69	70	Eng. Mean	Metric Mean	Group VIII 1	77	72	73	172	75	75	27	78	42		Eng. Mean	Metric Mean

TS																					
CBIENTIVE NEASTRE ENTS	hrea.	70	<u>ွ</u>	36 51.	# 00 10 10 10 10 10 10 10 10 10 10 10 10 1	1 (V	Z†	25	70	1 <u>2</u> 2		7,7	ار د	26	86	17	 	200	α α ι ο	2	238255888888 238888888888888888888888888
BUECTIVI	٠,١٠١	)• 50	07.	12.00		12.	7.50	000	ည် သ ()	10.78	•	60	11 10 10 10 10 10 10 10 10 10 10 10 10 1	12.	000	07.	~ ` ` ` ` ` ` ` ` ` ` ` ` ` ` ` ` ` ` `	30	9.92 63.98	)	11 CS
110	Koura	31.9	32.5	022 0.00	ر د د د د د	35.4	31.5	32.1	33.0	36.73	) 1	32.1	35°6	37.00	32°. 7°. Le	33.1	32°tr	9,4 4,5 7,0	32.76	• ()	
1	Kound	27.8	28.9	2.00	200	27.8	28.0	59.0	27.6	26.25	) •	26.6	26 26 27	27.0	0 0 0 0 0 0	28.5	0 0 0 0 0 0		27.52	1.0	25 25 25 25 25 25 25 25 25 25 25 25 25 2
Pradin I c	Erisket	3.10	장	w . 0 .		ν. ν.ν. ν.ν.	4.40	ල ද	00°-	01.17	1	2.40	4.50	, 4, 0, 0, 1,	4.70	1.05		1. 50	10.01	† O • • • • • • • • • • • • • • • • • •	11. 60 11. 60 11. 60 11. 60 11. 73 11. 73
at B	12th Rib	08.10	04.50	07.70	07°-11	07.50	00.70	01.70	00° 07°	7.56	•	08.50	08.10	08.30	07.50	07.75	09.70	07.70	<b>I</b> OLX	•	08.80 06.90 08.10 07.50 07.50 07.50 07.50
Depth at		6.90	5.75	06- 00-	, de 0	6.75	6.80	07.9	09 <b>.</b> 2	6.51 6.51	#C•0#	L75" 8.10	0 <sup>1</sup> - 2	7.60	7.30	9.00	07°9	0.09 0.09	7.20	<b>1</b>	.76-1-00" 6.75 6.75 7.70 8.50 7.30 7.30 7.30 7.30 8.50
Steer	0 1	uroup 1 .co.	8	m-:	<b>4</b> 70	<b>\</b> \	2	ω	ه د	Ing. Mean	וופסו דכ ו:פמוו	Group II .51	12 51	14	Z Z	17	<b>ი</b> ლენ	20 20 20 20	Eng. Mean	וייתמ	Group III .7 21 22 23 24 25 26 26 27 27 28 29 30 4ng. Wean Retric Mean

(pet	L.J. Area	05.33 06.33 06.03 06.13 06.73 07.53 07.53 07.53	242 242 242 242 242 242 242 242 242 242	2524 2524 2524 2524 2524 2524 2524 2524
I C (Continued)	Circ. of Round			$\begin{array}{cccccccccccccccccccccccccccccccccccc$
APPENDIX :	Lergth of Round	25.50 25.50	88888888888888888888888888888888888888	422445426925 505644564269
	Depth at Brisket	17/1/20/20/20 24/00/20/20/20/20/20/20/20/20/20/20/20/20/	44 www.www.u.u.u.u.u.u.u.u.u.u.u.u.u.u.u.u	ルフログログログログログログログログログログログラウル (1) (1) (1) (1) (1) (1) (1) (1) (1) (1)
	Depth at 12th Rib	08.10 07.60 07.20 08.50 07.70 07.70 19.61	00000000000000000000000000000000000000	
	Depth at 10th Rib			17777777777777777777777777777777777777
	Steer 10.	Group IV 1.0 31 32 33 34 35 36 37 38 10 Eng. Mean Metric Fean	Croup V .26 11. 12. 13. 14. 14. 14. 14. 14. 14. 16. 14. 16. 17. 18. 19. 50. 50. 50. 50. 50. 50. 50. 50	Group VI 522 523 523 524 525 525 525 525 525 525 525 525 525

ued)	L.D. srea	() ()	12°03	11.44	0,11	13.51	11.23	13.63	07.01	10.30	11.02	36-11	73.40		11.65	12.18	10.25	12.70	11.75	10.59	10.22	77.	10.25	11.00	12.11	72.30
I C (Scrtinued	Jire. of Round		36.3 36.3	37.7	35.7	25.7	38.1	ಗ್ ಎ സ	37.2	~1 C: 보기	77	36.35	3. W.		37.6	34.7	70.2	37.1	36.95	34.5	0. 10.	34.2	35.6	26.2	72.08	70°25
RICESS C	Length of Round	7 00	) (V	20.3	25.	31.3	28.3	20.32	30.5	30.1	30.8	83.52	05•11		28.6	25.8	30.0	30.5	30.5	30.5	?? ?\•	25° 57	30°	0.73	72.73	75.75
	Jephnach Brisket	< \ \ -	1 M	, 60 60	7. 0.	07.1	6.50	2.00	5.40	7.50	5.10	51.5	14.53		ე• 1.	3.60	00•9	6.10	5.20	5.70	4.30	м. 10	•	6.10	•	$\omega$
	Derin at 12th Bib	() () ()	000	07.50	07.70	09*50	05.60	07•30	08.10	08.10	08.25	ा <u>.</u> ा	21.34		07.50	10.40	00•60	03.70	0E•30	05.70	09.50	07.50	05.30	07.70	C5.3	21.59
	Jepth at 10th Pib	<del>-</del> 5										12	ري	.01-1.25"	7.20	6.50	00°3	6.80	7.40	C0•3	\$\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	6.39	7.60	6.80	70.7	16.40
	Cteer Fo	Group VII .7	J 0	63	t/9	53	99	29	39	69	0.2	Eng. Nean	Metric Mean	Croup VIII	<u>.,</u>	72	73	77	75	92	77	رم ص	79			Netric Mean

1			ı													•	- 41	U-																
	wt. Trmd.	Chuck		07.70	200	65.30 67.30	00	76.50	63.20	68.00	72.00	70.60 81.33	30.52		64.70	63.20 20.50	06°07 20°80	64,00	72.20	04.70 60.70	74.60	64.30	67.01 30.39		V3 87	62.20	64.70	76.50	68°59	62.50	67.90 73.50	000 000 000 000	52.33	ゴス・つう
	Wit. Untrad.	Shuck	0) 1)	00.40	30.00	65°30	000	76.50	63.40	68.20	72.75	70 <u>.</u> 60	31.01		64.70	64.40	71.20	64.70	72.80	03. 70. 70.	75.10	64.30	67.54 30 <b>.</b> 63	\	0° 39	65.20	64.70	01°00 01°00 01°00	70.20	65.70	6% 30	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	F. 53	TG*TE
	Wt. Trmd.	Rib	()	07.25 76.86	ก C ว น ว น ร ด		23.70	21.50	20.50	22,30	21,00	21.40 04.10	05.6		26.70	21.55 23.80	27.10	18.40	24.00	Oハ•ON	23.80	21.20	22.48 10.20	•	0.1.00	22.40	25.00	24.20	21.10	22.60	25.00 26.00	21.80	22.78	10.33
LE C'IT YIRLD	Wt. Untrmd.	Rib	Q Q	00.77	00°98	2. C	24.20	22.00	20.70	55.40	21.20	22.00	10.13		27.20	09.22	27.80	18.40	24.00	17. 50 00.00	25.10	21.40	23.14 10.15	:	טר אס	27.50	26.80	26.60	22.40	26.50	26.60 20 1.0	21.20	05.42	42 <b>•</b> TT
1	wt. Trmd.	Loin	00	34.60	ָּהָלְיּהְ הַאָּרְיּהְ	77. 37. 37.	33.70	32,30	37.50	39.10	32°50	37.80	16.17		35.60	34.20	35.30	39.20	36.70	ころい	29.30	35.50	36 <b>.</b> 13 16.38	1	00 57	34.50	36.10	30.10	36.60	31.30	36.40	かい ・	35.45	15.84
I A RIGHT SIJE	Wt. Untrmd.	Loin	( f	35.10	00° W	00°	35,10	32.50	37.50	40.10	36.80	37.80 36.50	16.56		37.70	37.50	37.10	05.14	00,000	30•10	32.60	05 <b>•</b> 9€	37.52 17.20	•	00 6.7	35.00 35.00	37.20	34.40	00°90 30°90	36.20	35.10 27.25		36.10	17.28
APPENDIX II	wt. Trmd.	Round	()	0) • To	0 0 1 1 1 1	0/1° (2)	67.55	62.50	55.00	65.70	05•19	7,000 CCC CCC CCC CCC CCC CCC CCC CCC CCC	27.52		54.10	5. 5√. 5√.	57.80	52.00	55.10	70° 00° 00° 00° 00° 00° 00° 00° 00° 00°	55.10	57.50	25° 35° 35°	\ \ \	70 00	15.00 10.00 10.00	03.60	0 0 0 0	54.70	49.20	51,00	52.70	53.37	24.20
	Wt. Untrmd.	Round		00.09		00,10	61,10	00.49	55.00	65.70	67.60	59.30	28.01		57.10	07.00	79.40 79.40	54.50	59.40	70°40	58.10	안	56.24 26.41		υ 13	57.10	24.80	0 2 2 3 3 3 3 4 3 4 3 3 3 3 3 3 3 3 3 3 3	57.60	53.55	53.50 EX.30	02°53	56.38	25.57
	Rt. Side		Ιĩ	25. 25. 25. 25.	0.002 0.002 0.002	200 200 200 200 200 200 200 200 200 200	06.552	256.20	00.743	264.50	260,30	252.50	116.79	•		0/1 a a c c c c c c c c c c c c c c c c c	265,60	260.70	273.70	257 TO	265.05 05	21,9,85	263.43		"00•T-9 <i>)</i> .	271.50	260.80	10/2 10/2 10/2 10/2 10/2 10/2 10/2 10/2	263.50	270.140	263.50 260 260		276.40	
	Steer	NO.	Group I .26	<b>⊣</b> c	3 6	<b>)</b> ~	<u>+</u> \	\\O	7	ω	0	2	Metric Mean	Group II .51	T i	2 T C	7 T	17	16	<u> </u>	15	20	Ing. Mean Metric Mean			7.2	23	700	26 26	27	<b>ω</b> (	300		Metric Mean

			-211-	
	wt. Trmd. Chuck	32000000000000000000000000000000000000	######################################	12128688888888 111386788888888
	wt. Untrad. Chuck	66.25 73.10 68.25 71.15 72.15 73.15 74 74 75 75 75 75 75 75 75 75 75 75 75 75 75	503559886586 5035599865858	11448848888888888888888888888888888888
	at. Trad. Rib	00.000.000.000.000.000.000.000.000.000	31.70 27.70 27.70 27.70 27.70 27.70 21.30	68888888888888888888888888888888888888
nd )	at. entrad. Pib	######################################	50000000000000000000000000000000000000	25 25 25 25 25 25 25 25 25 25 25 25 25 2
I A (Continued)		888888888888 607988888838	256000000000000000000000000000000000000	20 20 20 20 20 20 20 20 20 20 20 20 20 2
IL MICAGAN	wt. Untrmd. Loin	25.83.43.55.55.55.55.55.55.55.55.55.55.55.55.55	3043343444 8988988838	F1000000000000000000000000000000000000
	i.t. Trmd. Round	257777577755 257777777775 20122222222	68.77.77.77.77.77.77.77.77.77.77.77.77.77	76.50 77.50 77.50 77.50 77.50 83.06
	wt. Matrud. Round		25-55-55-55-55-55-55-55-55-55-55-55-55-5	874588888888888888888888888888888888888
	Kt. 3			1.36.77.33 3.77.33 3.77.33 3.77.33 1.60.73 1.60.73
	Steer No.	Group Land Restriction Restric	Oroup V • 26  11  12  13  14  15  16  16  17  19  19  19  19  10  10  10  10  10  10	Shoup YI . X Z Z Z Z Z Z Z Z Z Z Z Z Z Z Z Z Z Z

																-	21	2-									
nt. Trmd.	Chuck		24.30	102.30	51.85	87.20	69.80	67.40	82.50	69.68	51.65	65,60	36.06	41.23		67.40	00•95	09.05	93.40	55.25	85.90	65.60	65.40	85.20	80.10	න න න	70.62
at. intrad.	Chuck		94.30	104.30	93.50	90•30	92.50	93.00	86.30	61.60	93.50	09*16	93.43	42.37		05•63	00•86	09*1/5	97.40	00•75	67.50	98.10	21.45	88.00	87.60	07.57	11.50
Wt. Irmd.	Rib		25.70	27.20	28,20	26.80	30.10	28,00	25,00	25.90	30,00	25,30	27.62	12.53		25.60	32,00	25,20	28.60	27.40	22.50	28,50	29,00	25,30	28.75	27.08	12.55
wt. Untrad.	Rib		31.50	25.20	31.20	30.60	32.40	32.30	30.50	28.10	33.60	28,30	30.77	13.55		28.60	35.10	32.50	33.00	31.50	24.80	32.25	32,10	25.50	33.80	31.36	14.22
wt. Trmd.	Loin		42.70	48.50	56.44	42.10	7-40	36.55	36.10	1:4,10	70.80	14.10	12.57	15.49		39.70	47.50	42.20	00°44	35.70	50.20	45.40	43.40	43.50	47.60	90°ti	20.12
wt. Untrmd.	Loin		77.00	24.90	50.70	4.7 - 80	51.10	17.70	14.20	47.70	45.70	76.67	16.34	21.92		4.7.50	57.00	02.54	0£•30	15.60	02.09	75.80	75.60	50.70	55.30	51.37	23.30
Wt. Trnd.	Round		$\alpha$	$\alpha$	71.70	65.40	78.00	66.40	05.59	09•179	00*92	68.15	70.14	31.81		70.70	71.40	66.50	54°00	70.75	00*69	27.00	00•99	02.07	55.60	65. •58	31.55
Wt. Intrmd.	Round		79.20	•				•	•	•	•	•		•		<b>.</b>	<u>~</u>		Ö	9	ညံ	i,	C)	ده	ν,	٠.	34.55
Rt. Side	1. t.	1.00°T−92	352.10		•	•	•	352.70	353.00	•			1 •	•	.01-1	351.40	05°59E	365.70	365.25	352.75	376.60	380.75	340.20	358.15	u い い い	356.60	161.72
Steer	No.	>	9	62	63	719	65	99	29	99	69	20	Eng. Mean	Metric Mean	III	71	22	73	77	75	75	77	78	52		$\geq$	Metric Mean
	Rt. Side Wt. Intrmd. Wt. Trmd. Wt. Untrmd. Wt. Trmd. Wt. Untrmd. Wt. Untrmd.	Rt. Side Wt. Intrmd. Wt. Ind. Wt. Irmd. Wt. Wrind. Wt. Wit. Round Round Loin Loin Rib Ri	Steer Rt. Side Vt. Intrmd. Wt. Trmd. Wt. Trmd. Wt. Untrwd. Wt. Irmd. Wt. Untrwd. No. Vt. Round Round Loin Loin Rib Rib Chuck VII .76-1.00"	Rt. Side Wt. Tritmd. Wt. Trind. Wt. Trind. Wt. Untired. Wt. Trind. Wt. Trind. Wt. Untired. Wt. Trind. Wt. Trind. Wt. Untired. Wt. Untired. Wt. Trind. Wt. Trind. Wt. Untired. Wt. Unti	Side V.t. Intrmd. wt. Trmd. wt. Trmd. wt. Untred. Wt. Trmd. at. Untrad.         No.       A.t. Round       Round       Loin       Loin       Rib       Rib       Chuck         VII. 76-1.00"       VII. 76-1.00"       42.70       31.50       25.70       94.30         61       352.10       79.20       74.20       54.90       48.50       29.20       27.20       104.30	Stde       V. Introd       W. Irmd       W. Irmd       W. Irmd       W. Introd       W. Irmd       W. Introd       W.	Stde       V. Introd.       W. Irmd.       W. Introd.       W. Introd.       W. Introd.       W. Introd.       W. Introd.       W. Irmd.       W. Irmd.	Stde         Vt. Intrmd.         Wt. Untrmd.         Chuck         Chuck	Side W. Intrmd.       W. Irmd.       W. Irmd.	Stde       V. Intrmd.       W. Irmd.       W	No. N.t. Gound W. Trmd. N.t. Untrmd. W. Trmd. W. Untrmd. W. Trmd. N.t. Untrmd. No. N.t. Round Bound Loin Loin Rib Chuck Chuck VII .76-1.0011  61 352-10 79-20 74-20 47-00 42-70 31-50 25-70 94-30 65-40 71-50 56-40 47-80 48-50 31-20 25-20 27-20 104-30 65-40 71-00 65-40 47-80 42-10 30-40 32-40 30-30 65-50 66-10 44-20 36-55 32-40 30-30 25-30 86-30 66-30 66-30 66-30 66-30 64-60 47-70 14,-10 28-10 28-10 25-90 51-60	No. N.t. Side N.t. Trund. N.t. Untrind. Wt. Trund. Wt. Trund. Mt. Trund. At. Untrind. Wt. Trund. Mt. Trund. At. Untrind. At. Untrind. Mt. Trund. At. Untrind. At. Untrind. Mt. Trund. At. Untrind.	No. http://doc.org/lineary.com	No. ht. Side Nt. Intrmd. wt. Untrmd. wt. Trmd. wt. Untrrd. Wt. Trmd. it. Untrrd. Wt. Inch Ch. 30	teer Rt. Side Nt. Intrmd. Wt. Irmd.	teer Rt. Side	Rest Side N. Introd.         <	teer         Rt. Side         Lt. Irind         At. Unitude         At. Unitude         At. Unitude         At. Irind         At. Irind <t< td=""><td>Loser         Rt. Side         L. Introd.         At. /td><td>toer         Rt. Side         Lt. Trund         At. Untrud         At. St. O.         At. St. O.</td><td>Leer         Rt. Side         Lt. Tetrnd         At. Trnd         &lt;</td><td>Res         Rt. Side         Lt. Trund         Lintund         <th< td=""><td>Rt. Side (L. Trumd, wt. Trumd, w</td><td>Res         R. Side M. Trund. Mt. Mt. Mt. Mt. Mt. Mt. Mt. Mt. Mt. Mt</td><td>total         R. Side         H. T. Side         H. T. Littrid.         At. Trmd.         At. 30         A</td><td>Steer         R. Side         U. Trimit         A. T</td><td>  No.   State   C.   Tend.   A.   A.   Tend.   A.   A.   Tend.   A.   A.   A.   A.   A.   A.   A.  </td></th<></td></t<>	Loser         Rt. Side         L. Introd.         At.	toer         Rt. Side         Lt. Trund         At. Untrud         At. St. O.         At. St. O.	Leer         Rt. Side         Lt. Tetrnd         At. Trnd         <	Res         Rt. Side         Lt. Trund         Lintund         Lintund <th< td=""><td>Rt. Side (L. Trumd, wt. Trumd, w</td><td>Res         R. Side M. Trund. Mt. Mt. Mt. Mt. Mt. Mt. Mt. Mt. Mt. Mt</td><td>total         R. Side         H. T. Side         H. T. Littrid.         At. Trmd.         At. 30         A</td><td>Steer         R. Side         U. Trimit         A. T</td><td>  No.   State   C.   Tend.   A.   A.   Tend.   A.   A.   Tend.   A.   A.   A.   A.   A.   A.   A.  </td></th<>	Rt. Side (L. Trumd, wt. Trumd, w	Res         R. Side M. Trund. Mt. Mt. Mt. Mt. Mt. Mt. Mt. Mt. Mt. Mt	total         R. Side         H. T. Side         H. T. Littrid.         At. Trmd.         At. 30         A	Steer         R. Side         U. Trimit         A. T	No.   State   C.   Tend.   A.   A.   Tend.   A.   A.   Tend.   A.   A.   A.   A.   A.   A.   A.

	ı	1	-213-	
	Ext. Fat Trim		777777 770777101010110	949468469494 6585684469466
	Trad. Lean Cuts	182.20 174.20 185.20 185.20 195.20 195.20 195.20 195.20 195.20 195.20 195.20	131 131 131 131 131 131 131 131 131 131	22000000000000000000000000000000000000
(Continued)	wt. of Belly Cuts + K.K.	75.00 70.25 70.25 70.25 70.10 81.25 81.25 81.25 81.25	77 73 73 73 74 74 74 74 74 74 74 74 74 74 74 74 74	ロットキャスのアオロロド たののがかのがあるようだ ないのがかのかのものです
APFELUIX II н (Sc	irtrm Cuts	184.50 180.10 180.10 180.10 180.10 180.10 196.00 17.83	0.000 0.000	
TA.	wt. K.K.	06.20 06.20 06.20 06.20 06.20 06.20 06.20 06.20 06.20 06.20 06.20	11.000.000.000.000.000.000.000.000.000.	
	iwt. Flank	16.80 17.40 18.30 17.40 17.40 17.40 8.31 8.31		COCOCOCOCOCOCOCOCOCOCOCOCOCOCOCOCOCOCO
	wt. Plate	22.33 21.33	23000000000000000000000000000000000000	10000000000000000000000000000000000000
	wt. Brisket	16.46 10.70		54.75.55.55.55.55.55.55.55.55.55.55.55.55.
	ut. Shank	26-59" 10.20 10.20 08.05 09.00 09.00 09.60 11.30 11.30 08.10 11.30 11.30 11.30	4.000000000000000000000000000000000000	
	teer 5.	6 2 2 6 2 4 2 2 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Group II 11 12 12 13 14 15 16 17 19 20 3ng. Kean Year	200 20 20 20 20 20 20 20 20 20 20 20 20

	Ext. Fut Trim	24.52 25.52	101 005 005 005 005 005 005 005 005 005	0.00 11.00 11.00 11.00 10.00 1
	Trad. Lett. Cuts	174.50 173.70 181.65 161.65 171.70 174.70 17.63	217.00 217.00 253.70 235.20 226.05 2214.20 237.15 215.80 106.47	258.00 233.60 235.70 227.70 227.70 236.15 107.35
Confirmed)	.t. of Belly Juts + N.K.	82 25 25 25 25 25 25 25 25 25 25 25 25 25	111.65 100.10 100.40 090.40 090.40 090.40 090.40 090.40 090.40 090.40 090.40	057.10 109.40 107.40 100.50 100.50 101.10 101.65 101.05
) Y II VICE	f Entrad. 3 John	25.05.05.05.05.05.05.05.05.05.05.05.05.05	252.0 257.30 257.30 257.30 241.00 240.10 250.10 250.10 250.10 250.10 250.10	266.30 240.30 240.30 260.30 260.30 260.30 260.30 260.30 260.30 260.30 260.30 260.30
	X な X X	######################################	12.00 14.00 15.00 15.00 16.00 17.00	12.00 11.00 12.00 12.00 12.00 12.00 12.00 13.00 10.00
	i.t. Flerk	26.00 20.00	31.00 26.00 26.00 26.00 27.10 23.10 23.10	27.70 27.70 31.00 27.00 27.00 27.00 31.00 12.75
	n.t. Plate	67.75.01 67.75.01 67.75.01 67.75.01 67.75.01 67.75.01	37.60 83.82 83.82 83.83 83 83 83 83.83 83 83 83 83 83 83 83 83 83 83 83 83 8	25.00 20.00
	.t. Brisket	11	18.50 11.20 10.50 10.50 10.50 10.50 10.50 10.50 10.50	116.00 118.00 118.00 118.00 118.00 119.00 128.00 128.00 128.00
	i.t. Overk	3.46 3.46	66-70-11-70-	12. 25. 11. 25. 25. 25. 25. 25. 25. 25. 25. 25. 25
	Stoer No.	Group I/ 1 32 32 33 33 35 35 35 40 40 8et. Meen	Oroup V .2  11  12  13  14  15  16  17  19  19  19  10  20  En . Mean	Group VI.

	Ext. Fat	367 77	ä	10		GD.	O.I	Ċ	10	i	+		• •	~		•	•	•	•	•	•	•	•	•	•	•	6°•6
	Trid. Lean	3   3	•	•	•	•	•	•	•	•	•	•		105,06		•		•	•	•	•	•	•	•	216.05		10t 8%
Continued)	to of Belly	2	100.10	105.10	084.35	101.70	106.00	111.60	114.60	05.701	109.30	57.250	103 83	1,7.09		•	•	•	•	•	•		•	•	36° 560		47.89
Predict II A (do	t. of Untrad	001. 000	•	•	•	•	•	•	•	•	•	•		112.44													113.97
· :	۲. چ	•1	$\sim$	സ	ごヽ	$\omega$	CA	アノ	O.	$\sim$		$\mathcal{L}_{i}$	14.53	6.78		$\sim$	r	$\sim$	$\sim$	CA		$\sim$	$\Gamma$	9	12.85	- 7	6.75
	7. t. 1. r. r.	4	NO.	~	ıΩ	$\overline{}$	$\sim$	$\sim$	m	$\sim$	$\overline{}$	$\sim$	29.58	13.60		ρ. •	7.5	ο, •	9•0	1.2	い。シ	ω, ω,	6.9	1.1	28.20	1.3	T- 10
	4t. 10ء اتا	3	ω)	O	$\sim$	$\circ$	9	m	$\vdash$	⊣	Ŵ	$\circ$	32.16	14.58		o	ſν	$^{\circ}$	<u>.</u>	<u>-</u>	نی	$\sim$	۲.	'n	00°82	,	_: <u>†</u>
	1: t. n. 000	0 0 7 7	\Q	$\rightarrow$	CV	いノ	$\alpha$	$\omega$	$\circ$	77.	$\sim$	-1	16.31	7.41	ĵ.	$\widetilde{\dashv}$									16.20		7.64
	: t.	3	10.40	08.70	09.60	10.80	13.30	10.70	09.60	10.40	10,30	10,00	10.40	4.72	1.01-1.2	12,00	07.50	08.30	10,40	10,55	10.20	14.00	10.65	10,85	10.70	10.81	7.55
		Group VII	႕	62	63	79	65	99	29	68	69	70		Kea	Group VIII	71	72	73	77	75	92	77	78	79	63	Eng. Mear	Met. Mean

Bone Shank			00000000000000000000000000000000000000
Fa <b>t</b> Shank			E7488228888
Shank Yield			
Bone Chuck	000 000 000 000 000 000 000 000 000 00	88888888888888888888888888888888888888	777888888
IDE Fat Chuck	84588686668 846886866668	70698888986 80898888 80898888	002.50 002.50 002.50 002.50 003.50 00
RIGHT S Chuck Yield	4 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	0.4446464666666666666666666666666666666	272 2000 2000 2000 2000 2000 2000 2000
p p	240256322220 2402563222220 24025632222220	77108082808644 4710808288888484	
IL CUTCUT Fat Bo	000000000000000000000000000000000000000	044400404040 0345000404040	
B RETA Rib Yield	17.60 20.70 20.70 11.50		
IX II Bone Loin	2000 2000 2000 2000 2000 2000 2000 200	20000000000000000000000000000000000000	
APPENDIX Fat Bo Loin Lo		84988888888888888888888888888888888888	00110777777
Loin	22 25 25 25 25 25 25 25 25 25 25 25 25 2	25.20 23.20 23.20 25.20	50000000000000000000000000000000000000
Bone Reund	02.00 02.00 03.00 03.00 03.00 03.00 03.00 03.00 04.00 05.00 04.00 05.00		00000000000000000000000000000000000000
Fat Round	00000000000000000000000000000000000000		
			26-1-9-1-9-1-9-1-9-1-9-1-9-1-9-1-9-1-9-1-
ee o	1 . 26 . 3 . 3 . 56 . 56 . 56 . 56 . 56 . 56	. 111 122 123 123 123 123 123 123 123 123	111 - 22 - 23 - 25 - 25 - 25 - 25 - 25 - 25
	Group Ing. K	Group Frg.	Group etry

and the control of the 

		-21/-	
Spank	6002050505050505050505050505050505050505		
Fat Shank		000000000000000000000000000000000000000	01011111111111111111111111111111111111
Shank Mold			00000000000000000000000000000000000000
bone Obyck		117.00 11	111.70 112.70 112.70 113.70 113.70 113.70 113.70 113.70 113.70
rat Chack	05.70 08.65 08.65 07.60 07.70 07.70 07.70 07.70 07.70 07.70	06.30 06.30 06.30 06.30 06.30 06.30 06.30 06.30 06.30 06.30	07.60 09.95 111.90 12.00 12.00 12.00 12.00 12.00
d) Chack Yield	25.50 25.50	72.60 77.50 62.00 61.10 69.15 69.15 31.15	78.20 68.00 71.40 66.00 72.65 72.65 72.65 72.65 72.65 72.10 73.10
Cortinued Bone C Rib Y		~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	7001000010014 200100010014
) स म भूकर भूग	10001000000000000000000000000000000000	00000000000000000000000000000000000000	88898988888888888888888888888888888888
Richar I	00000000000000000000000000000000000000	22.20 24.60 20.20 20.20 20.15 20.15 20.15 20.10 20.10 20.10	22.10 22.10 22.10 20.13 117.50 117.50 117.50 117.50 117.50 117.50 118.30 117.50
n por Pone Loin	00000000000000000000000000000000000000	87798777998 8000000000000000000000000000	00000000000000000000000000000000000000
Fat Loin	23.000.000.000.000.000.000.000.000.000.0	17 00 00 00 00 00 00 00 00 00 00 00 00 00	84444448884488888888888888888888888888
Loin Yield	25.70 26.70 26.70 27.50	32 22 23 24 25 25 25 25 25 25 25 25 25 25 25 25 25	50 50 50 50 50 50 50 50 50 50 50 50 50 5
Bone Reard	87777778878 888888888888 8888888888888	21111111111111111111111111111111111111	21111111111111111111111111111111111111
Sat Pound	23000000000000000000000000000000000000	24598855888 24598859888	20008300073C
und eld	12. 10 10. 70 13. 20 38. 10 38. 10 37. 10 37. 10 17. 67	**************************************	25 55 55 55 55 55 55 55 55 55 55 55 55 5
ree No	Group 14 1. 32 33 33 34 35 36 36 36 36 10 Eng. Mean	Croup V .26- 11 12 13 14 15 15 17 18 19 50 50 For Mean	Oroup VI -52 52 52 52 53 55 56 57 58 58 56 56 56 56 57 57 57 57 57 57 57 57 57 57 57 57 57

bone Shank																											
bone Shank																											
		4.30	9.50 50	3°6	4.40	7.50	3.60	3.60	4.50	4.40	3.95	01.1	1.86			1 c	(1)	3.65	14.70	4.25	1.05	5.20	4.10	=1 (A (A)	3.70	4.12	1.87
rat Shank		1.20	0.40	1.50	0.30	1.00	1.60	0.20	1.10	1.00	1.17	8 0	77.0				0.00	1.25	07.0	1.20	0,0	1.60	1.70	1.70	1.60	1.14	0.52
Shank Yield																•	•	•	•	•	•	•	•	•	•		•
bone Chuck		11.45	12,00	11.70	10.70	13.35	04.50	05.30	12.20	12,00	12,30	11.47	5.20			•	•	•	•	•	•	•	•	•	•		•
ohack Ohack		10.60	16,60	10.75	14.65	08.20	13.10	12,10	10.40	09.85	12.65	53.11	5.39			) L	^>.	11.40	08,20	13.60	07.70	07.50	14.40	12,50	10.50	77.CI	4.73
Yield		71.80	73.40	65.55	61.85	68,20	64.10	02.09	67.00	00•59	06.99	67.25	30.50			•	•	•	•	•	•	•	•	•	•		•
bone Rib		7.50	93. E	5.40	4.55	5.30	3.70	7.00	7.60	5.10	5.20	90.1	2.11		<u>-</u>	1 / 1 (	3.	4.25	6.25	л, 00	00.4	7.80	5.15	5.30	77 S	4.54	2.24
D1		2.30	2.50	3.40	3.85	1.40	η <b>•</b> 00	3.40	1.20	3.20	2,00	2.72	1.23			•	•	•	•	•	•	•	•	•	•	•	•
Yield		22.40	50,50	19.40	16,30	23.40	19.70	17.25	19,80	21.70	17.70	20.02	80.8			•	•	•	•	•	•	•	•	•	•	•	•
$\circ$		•	•	•	•	•	•	•	•	•	•		•			•	•	•	•	•	•	•	. •	•	•	١.	•
3 O		2	4.15	3.45	7.00	1.70	3.20	2.20	1.40	3°00	4.25	3.12	1.41		6	•	•	•	•	•	•	•	•	•	•		•
) •~ (		•	•	•	•	•	•	•	•	•	•		•			•	•	•	•	•	•	•	•	•	•	١.	•
Round		11.60	05.50	11,20	10,80	12,00	08.60	08.30	12,10	12,35	10,60	10.52	4.95		02			11,00	12,30	10,80	11.50	12.40	10,50	12,50	0 <b>2•</b> 50	11.22	رن د د د د
Pound	=	•	•	•	•	•	•	•	•	•	•		•	: L	^\ _	) \ • • •	ひ•く	ි. දි	2.55	7.60	5.60	3.70	00•9	4.50	5.60	70.0	1.84
Yield Yield	6-1	Ü	m	-	ä	٦.	Ċ	9	6	ထံ	Ö	E	ţ.	-		000	54.30	55.60	58.70	54.15	54.80	60,80	149.140	53.30	1:4.30	2T°06	24.52
יט ע בייל כ	VII.	61	62	63	719	65	99	29	68	69	20	Nean	ic Mean	F + + 1.	T T T A	٦ (	V	$\sim$	_:;†	ጉ/	9	2	œ	٥,	ව	ean	Kean
	Group											_	· .	Č	grou											Eng B	letr.
	If nound fac bone bone brank fac bone outck fat bone Shank Yield Pound Round Yield Loin Loin Xield Rib Yield Chuck Yield	veer nound ray bone bone say bone star bone once fat bone Star bone Star bone Star bone Star bone Star bone St VII .76-1.00"	Veer nound rate Done File Fig. 2018 The Rib Rib Yield Chuck Tield Shank VII .76-1.00"  VII .76-1.00"  61 55.50 3.00 11.60 32.50 3.50 6.70 22.40 2.30 4.50 71.80 10.60 11.45 4.80	VII .76-1.00"  61 55.50 3.00 11.60 32.50 3.50 6.70 22.40 2.30 4.50 71.80 10.60 11.45 4.80 6.80 6.80 6.80 6.80 6.80 6.80 6.80 6	VII .76-1.00"  61 55.50 3.00 11.60 32.50 3.50 6.70 22.40 2.30 4.50 71.80 10.60 11.45 4.80 62 53.00 3.50 05.90 37.30 4.45 6.05 20.50 2.50 3.40 65.51 10.75 11.70 4.25 63.50 5.90 11.20 34.95 3.45 6.55 19.40 3.40 5.40 65.55 10.75 11.70 4.25	VII .76-1.00"  61 55.50 3.00 11.60 32.50 3.50 6.70 22.40 2.30 4.50 73.40 16.60 12.00 4.20 6.3 54.60 5.90 11.20 34.95 3.45 6.55 19.40 3.40 5.40 65.55 10.75 11.70 4.25 61.65 3.00 10.80 32.00 6.10 16.30 3.85 4.55 61.85 14.65 10.70 10.70 6.10	No. Yield Pound Round Yield Lein Lain Yield Rib Rib Yield Chack Chuck Yield VII .76-1.00"  KI .76-1.00"  KI .55.50 3.00 11.60 32.50 3.50 6.70 22.40 2.30 4.50 71.80 10.60 11.45 4.80 62 53.00 3.50 05.90 37.30 4.45 6.05 20.60 2.50 3.60 73.40 16.60 12.00 4.20 64 51.60 5.90 11.20 34.95 3.45 6.55 19.40 3.40 5.40 65.55 10.75 11.70 4.25 64 51.60 3.00 10.80 32.00 4.00 6.10 18.30 3.85 4.55 61.85 14.65 10.70 6.10 65 61.90 4.00 12.00 37.90 1.70 7.80 23.40 1.40 5.30 68.20 08.20 13.35 7.40	Fig. Yield Found Round Yield Lein Lein Kield Rib Rib Yield Chack Chuck Yield VII .76-1.00"  VII .76-1.00"  61 55.50 3.00 11.60 32.50 3.50 6.70 22.40 2.30 4.50 71.80 10.60 11.45 4.80 62 53.00 35.90 37.30 4.45 6.05 20.50 2.50 3.60 73.40 16.60 12.00 4.20 6.3 54.60 5.90 11.20 34.95 3.45 6.55 19.40 3.40 5.40 65.55 10.75 11.70 4.25 64 51.60 3.00 10.80 37.90 4.00 6.10 18.30 3.85 4.55 61.85 14.65 10.70 6.10 65 61.90 4.00 12.00 37.90 1.70 7.80 23.40 1.40 5.30 68.20 08.20 13.35 7.40 65.20 4.30 79.30 30.80 3.20 4.30 15.70 4.00 3.70 64.10 13.10 09.70 5.20	No. Yield Pound Round Yield Lein Lein Kield Rib Rib Yield Chack Chuck Yield VII .76-1.00"  VII .76-1.00"  61 55.50 3.00 11.60 32.50 3.50 6.70 22.40 2.30 4.50 71.60 10.60 11.45 4.80 62 53.00 3.50 05.90 37.30 4.45 6.05 20.60 2.50 3.60 73.40 16.60 12.00 4.20 63.60 5.90 11.00 37.30 4.45 6.55 19.40 3.40 5.40 65.55 10.75 11.70 4.25 64 51.60 3.00 12.00 37.90 1.70 7.80 23.40 1.40 5.30 68.20 06.20 13.35 7.40 65 52.20 4.30 09.30 30.80 3.20 4.30 15.70 4.00 5.30 68.20 06.20 13.35 7.40 67 56.40 3.80 09.30 29.10 2.20 4.10 17.25 3.40 4.00 60.70 12.10 09.30 5.60	No. Yield Pound Round Yield Lein Lein Iield Rib Rib Rib Jeld Chuck Fat Bone Shank VII .76-1.00"  VII .76-1.00"  61 55.50 3.00 11.60 32.50 3.50 6.70 22.40 2.30 4.50 71.80 10.60 11.45 4.80 62 53.00 3.50 05.90 37.30 4.45 6.55 19.40 3.40 5.40 65.55 10.75 11.70 4.25 64.56 5.90 11.20 34.95 3.45 6.55 19.40 3.40 5.40 65.55 10.75 11.70 4.25 64.50 61.90 4.00 12.00 37.90 1.70 7.80 23.40 1.40 5.30 68.20 06.20 13.35 7.40 65.20 4.30 09.30 30.80 3.20 4.30 15.70 4.00 3.70 64.10 13.10 09.70 5.20 67 56.40 3.80 09.30 29.10 2.20 4.30 17.25 3.40 4.00 67.00 10.10 09.30 5.60 68 4.50 56.70 12.10 35.40 1.40 6.70 19.80 1.20 09.30 29.20 4.60 6.70 12.10 09.30 5.60	Fig. Yield Pound Round Yield Lein Lein Xield Rib Rib Yield Chuck Jett Bone Shank VII 76-1.00"  VII 76-1.00"  VII 76-1.00"  61 55.50 3.00 11.60 32.50 3.50 6.70 22.40 2.30 4.50 71.80 10.60 11.45 4.80 62.50 3.50 73.40 16.60 12.00 4.20 62.50 3.50 73.40 16.60 12.00 4.20 63.50 3.50 3.50 37.30 4.45 6.55 19.40 3.40 5.40 65.55 10.75 11.70 4.25 64.50 5.90 11.20 34.95 3.45 6.55 19.40 3.40 5.40 65.55 10.75 11.70 4.25 64.50 5.20 4.30 12.00 37.90 1.70 7.80 23.40 1.40 5.30 68.20 06.20 13.35 7.40 6.50 55.20 4.30 12.30 37.90 1.70 7.80 23.40 1.40 5.30 68.20 06.20 13.35 7.40 6.50 55.40 3.80 05.30 25.10 2.20 4.30 15.70 4.00 6.70 13.10 09.70 5.20 68 69 50 2.70 12.10 35.40 1.40 6.70 19.80 1.20 4.60 67.00 10.40 12.20 4.60 67.00 10.40 12.20 4.60 67.00 09.85 12.35 31.50 3.00 6.00 71.70 3.20 5.10 65.00 09.85 12.00 4.60	Fig. Yield Found Round Yield Lein Lin Xield Rib Rib Yield Chack Chuck Yield VII 76-1.00"  VII 76-1.00"  61 55.50 3.00 11.60 32.50 3.50 6.70 22.40 2.30 4.90 71.60 10.60 11.45 4.80 62 53.00 3.50 37.30 4.45 6.05 20.60 2.50 3.60 73.40 16.60 12.00 4.25 64.55 19.40 5.40 65.55 10.75 11.70 4.25 64.55 10.70 11.20 34.95 3.45 6.55 19.40 3.40 5.40 65.55 10.75 11.70 4.25 64.50 12.00 12.00 37.90 1.70 7.80 23.40 1.40 5.40 65.50 13.35 7.40 65.50 12.00 12.00 37.00 12.10 35.40 1.40 6.70 12.10 37.00 12.20 4.50 65.00 67.00 13.10 09.70 5.20 68.50 12.30 12.10 35.40 1.40 6.70 19.80 1.20 67.00 10.40 6.70 12.20 4.60 67.00 10.40 6.70 12.20 4.50 67.00 10.40 67.	Fig. 1961 Found Round Yield Lein Lein Yield Rib Rib Yield Chuck Yield VII 76-1,00"  VII 76-1,00"  61 59,50 3.00 11.60 32.50 3.50 6.70 22.40 2.30 4.50 71.60 10.60 11.45 4.80 6.50 3.50 3.50 3.50 3.40 6.50 3.60 73.40 16.60 12.00 4.20 6.20 6.50 3.60 73.40 16.60 12.00 4.20 6.20 6.50 3.60 73.40 16.60 12.00 4.25 61.50 4.55 61.65 10.75 11.70 4.25 61.50 4.00 12.00 37.90 1.70 7.80 23.40 1.40 5.30 68.20 68.20 68.20 13.35 7.40 6.50 67.50 4.30 69.30 29.10 2.20 4.30 17.75 3.40 4.00 5.30 68.20 68.20 68.20 13.35 7.40 6.50 67.50 12.10 35.40 1.40 6.70 17.25 3.40 4.00 67.00 10.40 12.20 4.60 67.00 17.25 3.40 4.00 67.00 10.40 67.00 10.40 67.00 10.40 67.00 10.40 67.00 10.40 67.00 10.40 12.20 4.60 67.00 10.40 67.00 10.40 67.00 10.40 67.00 10.40 10.40 67.00 10.40 67.00 10.40	Fig. 1914 Found Round Yield Lein Lein Isial Rib Rib Yield Chuck Field Chuck Yield Chuck Signo 3,500 3,	VII 76-1.00"  VI 7	VII 1.01-1.25"  VIII 1.01-1.25"	VII 76-1,00"  VI 7	VII 76-1.00"  Viel Found Round Yield Lein Lein Field Rib Rib Yield Chack Chuck Field  VII 76-1.00"  VII 76-1.00"  VII 76-1.00"  VII 1.01-1.25"  VIII 1.01-1.25"	Fig. 1904 Found Round Yield Lein Field Rib Rib Yield Chuck Jene Shank VII 1.01-1.25"  VII 1.01-1.25"  VII 1.01-1.25"  VIII 1.	Fig. 1 (2) 1. (2) 1. (3) 1. (4	Fig. 1. Solution 1	Fig. 1 Sound Bound Yield Lein Lein 1901d Rib Sib Chack Fig. 2010 Sib. 176-1.00**  (c) 55.50 3.00 11.60 32.55 3.56 6.70 22.40 2.30 4.50 71.60 10.60 11.45 4.80 6.55 51.00**  (c) 55.00 3.00 11.60 32.55 3.56 6.70 22.40 2.30 4.50 71.60 10.60 11.45 4.80 6.55 51.60 3.60 5.90 11.20 34.95 34.45 6.55 12.40 34.0 54.0 64.55 10.75 11.70 4.25 61.50 11.20 34.95 34.45 6.55 12.40 14.0 54.0 64.55 10.75 11.70 4.25 61.50 11.00 12.00 37.90 11.00 6.10 6.10 6.10 6.10 6.10 6.10 6.	Fig. 1 (2014)  Fig. 2	Fig. 1 (2) 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.0	Fig. 71-10.  10. 57-10.  10. 5	THE POINT BOUND WILLS AND LONG STATE THE STATE STATE CHARK STATE AND STATE STA	Fig. 19 and Found Round Mich Lein Field Rib 19 1913 Chark Chuck Field WII .76-1.00"  (1) 59.50 3.00 11.60 32.50 3.50 6.70 22.40 2.30 4.50 17.40 10.60 11.45 4.80 6.5 59.00 3.50 11.20 34.50 6.50 2.50 3.40 73.40 16.60 12.00 12.20 6.5 59.00 3.50 11.20 34.50 6.50 2.50 3.40 73.40 16.60 12.00 12.20 6.10 6.10 12.00 12.20 6.10 6.10 12.00 12.20

j	1					
Tetal Bone	36.30 42.20 32.20 32.20		6 5 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	24 C C C C C C C C C C C C C C C C C C C	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	ンクエエエコ
Int. Fat Trim	28 28 28 28 28 28 28 28 28 28 28 28 28 2	22.55 23.55	11 10 14 14 13 13 13 13 13 13 13 13 13 13 13 13 13	10000000000000000000000000000000000000	23.50 22.50 22.50 25.50	20.00 20.00
Retail Yield	166.70 164.35 168.15 175.40	1444444 166778474 146084474	166.45 165.00 174.30	186.60 172.30 174.80 164.90 174.90 78.55	190.90 171.60 176.40 174.10 167.45	1000 1000 1000 1000 1000 1000 1000 100
Tield Lean Outs	0 4 7 4 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9	66. 57 66. 57 66. 57 66. 57	137.75 137.35 14.35 18.80	64. 35. 55. 56. 56. 56. 56. 56. 56. 56. 56. 5	153.30 155.30 155.30 137.30 15.75 15.00 15.00	139.05 131.15 131.60 62.33
Done Flark	}	poddinie poddarelep	00000		000000000000000000000000000000000000000	00000000000000000000000000000000000000
(Sectioned) ank rat eld Flank	00.00 00	05 70 00 00 00 00 00 00 00 00 00 00 00 00	~ 60 € 67 60 €	000 11 11 12 12 13 13 13 13 13 13 13 13 13 13 13 13 13	13.80 13.70 07.60 12.90 13.30	13.50 13.50 14.10 14.10 6.35
∞दिः इ	11.60 08.40 08.60 10.35	00000000000000000000000000000000000000	27.500 20.00	000 000 001 000 000 000 000 000 000 000	13.00 00.50 10.80 00.55 00.25 00.75	06.20 09.30 07.70 08.40 5.15
Bone Flate	0 2 2 2 0 2 2 2	100000000000000000000000000000000000000	000000 000000	2 2 2 2 2 2 2 2 3 2 3 3 3 3 3 3 3 3 3 3	00000000	14196 200 200 200 200
APPER I	00.00 00.00 00.00 00.00	иделемием СССССОВИИ СССССОВИИ		20000000000000000000000000000000000000	08.20 09.00 09.00 09.00 08.30 08.30	13.85 07.80 08.55 12.50 14.14
Plate Field	*	12.50 10.00 10.00 10.10 10.10 5.01	12.50 12.50 11.60 11.60	12 12 15 15 15 15 15 15 15 15 15 15 15 15 15	12.30 10.00 10.10 10.60 10.60 09.60	10.00 10.00 10.70 09.90 10.37
Bone Erisket	2.00 1.00 1.00 1.00	144444 222864444	44046 07000	1890 1890 1890 1890 1890 1890 1890 1890	44.00 00.00 00.00 00.00 00.00	00000000000000000000000000000000000000
Fat Brisket	4.00 6.20 3.50	14000000000000000000000000000000000000	7,1,44 0,000	10000000000000000000000000000000000000	474676 6006944	20020 20020 20020 20020
Erisket Yjeld	1.5	1	,	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	76-1-00. 6.70 7.50 1.70 1.85	6.10 6.10 6.13 7.75
Steer No.	Group I .26.	5 6 7 8 10 Erg. Yean Metric Mean	Group II .51	16 16 17 19 20 Eng. Mean Metric Mean	Group III 22 23 23 24 25 25 25 25 25	27 28 29 30 Sng. Mean Metric Mean

		0 W W 0 0 W 0 0 W 0 0 ~	M	070000000000000000000000000000000000000
	Total Bene	25.10 27.65 28.15 38.65 30.55 30.55 30.65 13.17	22 46 46 46 46 46 46 46 46 46 46	24444444444444444444444444444444444444
	Int. Fat Trim	25.00 25.00		20222222222222222222222222222222222222
	Retail Yield	155.55 176.35 176.35 176.35 117.20 11	22222222222222222222222222222222222222	236. 226. 226. 220. 220. 230. 230. 230. 230. 230. 230
	Nield Lean Cuts	133.10 143.95 143.95 143.95 132.10 124.70 125.45 127.96 127.96	162.50 204.30 175.60 173.60 177.50 184.50 82.31	24444444444444444444444444444444444444
	Bone Fiert	010000000000000000000000000000000000000	5120101010101010101010101010101010101010	200000000000000000000000000000000000000
tinued)	Fat Flank	18.10 19.20 19.20 10.20	20000000000000000000000000000000000000	200 200 200 200 200 200 200 200 200 200
B (Continued	Flank Jield	06.10 06.10 06.10 06.10 06.10 06.50 07.20 07.20	12.10 13.10 13.20 13.20 13.20 10.20	11111111111111111111111111111111111111
П	0	22 22 22 22 22 22 22 22 22 22 22 22 22	11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	00000000000000000000000000000000000000
APPE	Fat Bone Flate Plat	12.20 111.00 111.00 111.00 10.50 112.50 07.60 08.60 10.36	23.00 20 20 20 20 20 20 20 20 20 20 20 20 2	111.50 113.20 113.20 113.20 113.20 114.70 115.20 117.70 117.70
	Nate Yield	05.25 113.60 113.60 113.60 06.70 06.70 06.70 10.10 10.10	25.54.64.64.65.65.65.65.65.65.65.65.65.65.65.65.65.	12.00 13.00 13.00 13.00 13.00 13.00 13.00 13.00 13.00 13.00
	Bone Brisket	00000000000000000000000000000000000000	64564646466666666666666666666666666666	44444444444444444444444444444444444444
	Fat Brisket	7.70 8.70 8.70 8.70 7.70 8.70 8.30 8.30 8.30	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	00000000000000000000000000000000000000
	Y Y	11.25.00 5.40 5.40 5.40 5.40 6.50 6.	~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~	27.20 8.80 6.20 6.20 6.20 6.20 6.20 6.20 6.20 6.2
	er.	Group IV 1.0 31 32 32 33 34 35 35 36 37 10 10 Metric Mean	Group V .26.   11   12   13   14   15   15   15   15   15   15   15	Sroup VI -51 52 52 52 53 55 55 55 55 55 55 55 55 55 55 55 55

and the second of the second o 

1	Bone	ł	ロン・バン	32.05	12.10	70.20	び。 上	34.10	34.40	13.50	11 11 11 11	11.28	77.17	10.45	Č	ウリ シ・ マハ-	10.00 10.00 10.00	38.60	2.4	13.05	C:• 17	47.70	30° (A	10° 0	3.0		15.46
Tot. 299	Trim	7 6 7	00° (V	OT•0).	50	66.85	58.45	73.75	64.20	09•09	65.50	64.20	64.33	25.16	_	ઇ	۱	٠.	Š	ä	ં	<u>-</u> -1	•	$^{\circ}$	<b>1</b> .	0.1.0	زن
Retail	Tield	) () () ()	2772	20.422	210.15	202,20	229.75	205.15	207。12	201.50	220.80	271.25	212.47	56 <b>.</b> 36	-	07. 4.70	07.57 07.57 07.57 07.57 07.57 07.57	200.10		205.60	275.00	23, 60	100 M	() () ()	755.5	S. 7.	9 <b>5.</b> 95.
[]	Lear Cuts	•	02.001	Tel4.30	178,50	163.75	151.40	166.80	163.45	171.70	181,10	167.60			, , , , , , , , , , , , , , , , , , ,	1/6.50 2/6.90	ि• <u>२</u> १२	173.70	101 101 101	168.60	179.10	152.80	166.30	144,10	02°公司	20.771	2; •66
BOLE	Flank	ر ب		0.7.	01.	•10	01.	֖֖֖֭֭֭֭֭֭֭֭֭֭֭֭֭֭֭֭֭֭֭֓֞֝֞֝֝֟֟֝֟֝֟֝֟֝ <del>֡</del>	.10	.10	62.	5	, (2)	ω\ C •	ć	<b>.</b>		- - -	۱۱\ ط •	() ()	03•	.20	id N	() 다	C.L.	i.	.07
Continued)	Flank	76 71	10° (7	CO• ZZ	15.70	17.70	15.40	27.12	20°00	17.50	15.60	17,15		.t .o. .o.		•	ំា	•	ر	Ö	•	دی	·	22,10	• د ين	10.01	O)• √
्म् ) ह	Yield	) C	シャン・	15.15	09.85	13.40	10.40	11.15	13.10	ල <b>ී</b> දිට	11,20	0 <b>9.</b> 3克	11.67	٠. م	1		77.	11.50	11.50	10,3%	0%.50	N O	00. 20.	57.30	0E 50	S = 13	ν. ΕΟ•
DOIN II	Plate	ć		67.5	5.60	2.10	7.60	2,10	0. 0.7	2.40	3.10	3.30	(J	T.33	(	ू १	(), ,	2.70	7.50	3.40	2.60	3.10	رم س بر	3.33	S - 70	ري د د د د	ار سا ا
H 17	Fl.ate	00 7 5		13.55	12.20	11,65	19,55	16.10	13.20	18,10	15.50	17, 20	15.32	6.5	( (	07.27	1.0° 5.1° 1.0° 1.0° 1.0° 1.0° 1.0° 1.0° 1.0° 1	17.35	13.70	16.10	14.00	15.80	13.50	16.20	13.20	14.02	£.63
Plate	Tield											12.80		6.14	ì	70	7.0	Δ, Ω	ry.	7.7	ري ا	m)	۲. در	3.7	ئے	13.75	N
Вопе	Brisket		•	•	•	•	•	•	•	•	•	1.50	•	<u>.</u>		•	•	•	•	•	•	•	•	2.00	•	7:5T	\$9 <b>.</b>
Pat	Brisket	1	•	•	•	•	•	•	•	•	•	S S	•	•	\	•	•	•	•	•	•	7.60	•	10.50	7.60	†:P• <u>/</u>	
Brisket	Yield	1	•	•	•	•	•	•	•	•	•	6.70	•	•	1,01-1,25"	•	•	•	•	•	•	•	•	7.10	•	•	त् ()
<b>2</b> € € € € € € € € € € € € € € € € € € €	? 22,	· II,	J (	ç5	<b>%</b>	79	65	99	29	89	69	<i>:</i>	AT.	Metric Mean	H H									<i>6</i> 2		รักฐ. รัยลภ	Fetric Jean

	ht. Lean Rel. + Remp	43.90 35.30 35.00	C	12.20 10.70 10.70 18.20	28 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	64466666666666666666666666666666666666
ECTS THEIR	Mt. Rrd. + Rump	61.79 60.39 57.70	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	67.30 67.30 61.60 77.71	2272827488256 2425237788256 24552777886756	~&~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~
の 記 1	kt. Bone Rump	2.60 2.50 2.10	0 0 0 0 0 0 0 0 0 0 0 0	23.30		44444444444444444444444444444444444444
F RET BOXES	l.t. Fat Rump	8 7 9 10 8 10 00 00 00 00 00 00 00 00 00 00 00 00	44 m w e 0 17 0 6 7	250 mm 1	HTTWAWWTWWTT RESONCE	74 5 5 6 6 8 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
BP (BATIOT) OF	kt. Lean Rump	12.90 08.30 09.30	00000000000000000000000000000000000000	05.50 05.50 07.00 07.00 07.00 07.00	00 00 00 00 00 00 00 00 00 00 00 00 00	79.000.000.000.000.000.000.000.000.000.0
3.000 S	i.t. Rump	16.70 15.80 14.80	17.20 13.75 17.70 11.60	16.20 16.20 15.20 7.11	14.80 12.00 14.10 12.00 14.70 14.70 16.70 17.70	12.30 14.30 14.30 14.30 14.30 14.30 14.30
	M.t. Bone Reund	6.10 6.50 6.50	~~~~ ~~~~~ ~~~~	2000 m		
I TOIS	i.t. Fat Round	05.10 10.10 06.10	0.000000000000000000000000000000000000	00-70 07-15 07-15 07-15 07-15	05.15 07.50 07.50 07.50 07.50 07.50 07.50 07.50 07.50 07.50 07.50 07.50	00.00 00
5	lean it. Round	31.00 27.00 29.70	2000 2000 2000 2000 2000 2000 2000 200	31.00 31.00 31.00 31.00 13.00 13.00	26.80 28.10 28.10 28.10 27.30 28.40 28.40 28.40 28.40 28.40 28.40	28.55 27.50 27.50 27.50 27.50 26.50 26.30 26.30
	1 .	6.50°°°°°°°°°°°°°°°°°°°°°°°°°°°°°°°°°°°°	2007 11 12 12 13 14 14 14 14 14 14 14 14 14 14 14 14 14	21.27 15.60 15.60 15.10 0 20.60	12.75" 142.10 143.10 143.30 143.30 143.30 143.50 143.50 143.50 143.50 143.50 143.50 143.50	.76-1.00" 144.50 146.10 150.00 122.50 122.50 122.50 122.10 123.10 123.10 123.10 123.10
	teer No.	C 🔾	コンクレロ	<b>n</b> 60	Group II .  12  13  14  15  16  17  17  18  20  Eng. Mean	Group III 22 22 23 25 25 25 25 25 26 27 28 29 30 30 Xetric Mean

	ht. Lean Rnd. + Rumn		31.70	ルン・ スペー みな。 みな。	33.50	31.10	31.25	31.40	31.20	7.5.70 20.70	() () () () () () () () () () () () () (	1	\ - -	は、一つで、一つで、一つで、一つで、一つで、一つで、一つで、一つで、一つで、一つで	27.70	04.84 14.	14.50	18.50	69.75	19.15	03 <b>-</b> 51 03-51	22.59	18.75 16.50 16.10 17.00	50 77 87 87	. ду (. . ч. с. . С. д.	12.69	ניייי ביייייייייייייייייייייייייייייייי	7 2 A
	Wt. Rimn	4.000	7. 7. 3. 3.	ייער סעייער סעייער	27.00	XX.20	52.30	53.30	17°-70	いる。 かん かん	57.172 57.172	1	) )	70.75 70.75	81.70	cZ-9Z	70.70	63.70	80.60	75.60	75.30	34.12	76.83 74.80 74.20	81.40 78.30	61,60		77.60	- 01
	Ut. Bone Rumn		2.10	05°L	1.88	2.20	1.45	1.15	02° H	00.2		•	(	7.0 7.0 7.0 7.0 7.0	3.25	3.20	0 0 0 0 0 0 0 0 0		3.65	3.10	3.16	1.43	8.5.30 8.5.30 5.5.30	0.00	5.00 C.1.00	1 CA	Ci (-	) () ) ()
Cortinued)	Mt. Fat	2.	v. 55		700.7	6.10	ლ ლ	4.20		かっている。	7 T V	) •	\(\frac{1}{2}\)	ん ひ い い い	1.95	CZ•77	у, С С Л	, w	3.65	0S. 7	3.70	2.13	7,0 V,0 V,0 C,0 V,0 C,0	1.10 2.50	74.4 0.0 0.0	. 6 . 6 . 6 . 6	2. C	() (U) () (U)
) w III	kt. Lean Rumn	7	07.70	04.65 06.65	06.20	05.50	04.50	05.15	<b>୍ଦ</b> ୍ଧ ଅଧ	00°,400	06.43	1	(	11,30	13.30	08.50	0.40 0.40 0.40	02.00	05.70	00°00 00°00	000000000000000000000000000000000000000	04.43	11.35	08.10	(C)	か。 う う う う う う う う う う う う う う う う う う う	유. 	1 m.
APPENDIX	i.t. Rumn		15.40	11.70	12.20	13.90	05.50	10.90	13.40	5.01	12.53 5.86	)	0	20.00	21.70	17.50	12.00 17.00	13.70	17.00	16.10	17.75	8.05	21.50 21.50 18.40 50.50	16.10	01.71	S. S	다. () ()	
	wt. Bone	Siri A	ν.	1 L	, М	5.20	ር	5.20	۲. ۱. ۱.	ул У.У.	7 <u>1</u> 7			• •	•	•	•	• •	•	•	07-0 07-0 07-0	•	8.30 7.10 7.20 7.50	• •	•	• •	•	
	Wt. fat Round	31	•	• •		•	•	•	•	•	00/ (12 01/ 1/5	•	-	2, ω	3.0	2.0	ے ر ک (	) ()	. 5	900	00.00	4.5	11.50 10.55 13.20	12.50 15.10	11.70	11.50	11,00	) () ) () • () • () • () • ()
	Lean Mt. Round		÷.	<b>1</b> 0	, <sub>[</sub>	$\mathcal{V}_{\mathbf{v}}$	ġ	ø.	_i	ນໍ ດ	26.32	1	C	n O	4	~	<u>ن</u> -	1 TV	O	٦,	10.05 10.05	4	37.75 38.00 38.00 39.00	\.\.\.\.\.			0	•
	Wt. Round		•	•		•	•	•	•	•	19.06	ì	S (	~\~	Ö	င်္သံ ۔	-i c		ี่ ค้	0,0	60.00 58.48	26.		υλα	ت +ت	. (V	di	1 80
	Steer	>	31	J (.	ابر آبرا	32	36	37	∞ M	λ ( Υ) –	Ing. Mean Vetric Mean	) )	Group V .26	10	13	777	7 T	277	168	2.1.2 2.1.2	50 ng. Mean	Metric Mean	oroup VI SI SI SI SI	ኒ ኒኒኒኒ	. <b>Γ</b> ~ α	. o.	•	Fotzje Tesn

	1																-	22	24-	•								
	t. Lean	7rd. + 3.mp		○c • 1 ::	1.6 .80	05.54 5.4	다. 다.	1.7.30	03.04	06.00	スペッパイ 2.5.4.4.2.5.5.4.4.4.4.4.4.4.4.4.4.4.4.4.	15.10	16.50	17.5	20 <b>.</b> 15		42.00	4.1.5	43.70	75° 85	15.10	1,1.75	48.60	43.05	7.60	1.0 .65	35. 11.1 00 €0	77.7
	: t. R.S.	1 1 1 1		€ 00°	75.70	77.10	72.10	24.80	70.40	74.10	76.10	75.57	76.20	(1.27)	37.06		•	•	•	•	•	•	•	•	•	• !	75.76 21. 26	•
	Et. Bere	Rimin		س د. پرز	2.85	ა ლ•	2.20	3.10	S. S.	C 7 .	3.10	3°00	) -:'	2.00	1.21		2.30	0 N	2.60	3.20	2.40	س آن آن	3.30	2.1.0	3 <b>.</b> 00	2. FO	ر. اور اور	† , • <b>+</b>
(Cortinued)	:: :::::::::::::::::::::::::::::::::::			9.50 9	€.1,≎	00°2	⊖ <b>2</b> •9	S. 23	6.30	رب ور• برب	6 <u>.</u> 00	ر دن دن	ν. C	97 <b>.</b> 9	£.53		09 <b>•</b> 9	7.90	7.60	양	4.60	7.50 CX.7	00.9		7.70	6.00	ئ ئ ئا ر	<b>J</b>
V III	it. Icar	J. 11. 1		OL:30	11.80	10°60	03.30	02.50	11.20	10.10	10,80	06.30 00.30	07.20	90. めC	04.47		10.20	05.15	10.70	03.11	08 <b>°</b> 30	12,00	13,00	06.05	09•30	04.00	) a () a () a	1.40
XIC Reserve	٠. تا	R.zm.		15.50	21.20	02°03	16.75	18,00	20,20	05.03	20 <b>.</b> In	19,50	15.15	() () ()	22.3		cí•51	1,9,60	21.50	20.20	16,00	22.70	22.50	17.50	15.60	15.50	17.42 19.83	1.00
	at. Bone	Round			نرين •	1	(* )	α, )	r:	-1	$\mathcal{I}$	7	-	100	C )		•	•	•	•	•	•	•	•	•	• 1	ري. دد د	•
	.t. Fat	Round		٦.	Ċ	~,	_=1	Ċ.	;	ं	۲۰۰۱	1,5	_ = :	13.3	•		<u>.</u>	w	υŇ	ς,	$^{\circ}$	<u>.</u>	$\sim$	ij			L. かい。 かい。 かい。 で、	
	Loan Wt.	Pennd	1	υ,	λέ,	_:1	رن.	<u>.</u>	UN	ਂ	_=	١٤٠	O.	17. 17.	L'\	254	-i	c u	m	۲-	Š.	α	$\dot{\mathbf{w}}$	น้	~	v.L	0 -1 1 1 1 1 1 1	•
	7.t.	Rennd	) ]													H	55°55	55.50	56,50	C3.62	57.80	54.60	55.30	56.75	60,20	5 10 10 10 10 10 10 10 10 10 10 10 10 10	ສຸນ ທຸນ ກຸນ	60.63
	Staer		Group VII.	61	(1)	69	(F)	<u>(</u> ح	\$	67	<b>9</b>	69	62	:: ::	Metric Mean	⊬4 ⊬1 ⊬4	77	72	73	7.4	77.		77	78	52		Mean	) T.T.

1 1		-226-	
wt. Chack	65 12 12 12 12 12 12 12 12 12 12 12 12 12	60000000000000000000000000000000000000	
wt. Bone Rib	00000000000000000000000000000000000000		mobbook combo
wt. Eib	2011 2011 2012 2012 2013 2013 2013 2013	0.0011100000 0.001110000000000000000000	
at. Lean	0.1111200000000000000000000000000000000	0 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
: 요: 나 한 한	68888888888888888888888888888888888888		
(Continued) .t. Jone	20000000000000000000000000000000000000	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	
III A t. Fet Loir	25. 25. 25. 25. 25. 25. 25. 25. 25. 25.		66666666666666666666666666666666666666
APPENDIA At. Lean A	00000000000000000000000000000000000000		0.000000000000000000000000000000000000
조 다 다 다 라 :	15 00 00 00 00 00 00 00 00 00 00 00 00 00	######################################	00000000000000000000000000000000000000
wt. Bone Rad. + Rume	79977999477799 79777999797779		
13 Pu		: : : : : : : : : : : : : : : : : : : :	THE TOROUGH TO THE TRANSPORT OF THE TRAN
Steer No.	Group 1V L., 32 32 33 34 35 35 35 30 40 40 Metric Kean	Group 7 .26- 11 12 12 143 145 145 145 145 145 145 145 145 145 145	The state of the s

i	1	-227-
+ ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ;		87.72 23.72 23.73 24.73 25.23 27.73
.t. Bone	TANDONNONTA ANDONNONTA Laurananian	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
t. Fat		
t. Lean		5444444444 5646888866888 78688886888
. t.e	######################################	\$5550000000000000000000000000000000000
-: د.		20000000000000000000000000000000000000
	I row or repersion	60000000000000000000000000000000000000
ing Archin		286988888888888888888888888888888888888
رب اب در اب در		24888484488888888888888888888888888888
t.t. Cone	5088455511664 5088455511664	
Et. Fat		10.00 10.00
۲۱	Sean Rean	Group VIII 1. 72 73 74 74 75 76 77 77 78 78 78 78 80 Eng. Mean

	W.t. bone		C17.	1.30	1.50	1.20	1.15	1.20	07°-1	) () () ()	122.		76~0	1.00	1.20	ц.	1.37	2.70	щ О		1 S C C	<b>T</b> ⊃ • O		0.00	1.00	יע אר ס ה ריין ר	1 H	1.15	0.75 CC L	0 H	100	0.7°C
	it. Fat Erisket		CC 4	20°20	ነ ፓ./ - (ሳ - ሲ./	5.40	4.10	01.0	2.5 M	1.00	1.55 2.21		09°5	4.70	09•9	S S	7.00	3.30	η Ο γ	00.0	1 1 0 0 1 0 0	( † • 7	,	ر ا ا	C	7.55	7.10	2.60	7. 7. 7. 5.	л С	[] []	2.5
ľ	Mt. Lean Eristet		ار و و و		03.4	6.10	ਨ. ਨੂੰ	0.00 0.00 0.00	7. 66.	) (1) (2)	2.25 2.29		01.10	50 7	, 0	φ • • • •	79.7	3.30	w- ℃ 0	200	60-	○ <b>⊤•</b> γ	:	7 00 10 10 10	C. C.	м, и 0 с	6. 5. 5. 5. 5.	5.20	ν, Τ. Ξ. ς.	L.10	5.20	2.30
	∵t. Arisket		03. C3.	13,20	11.70	12.80	10.70	11.45	10,40 04,40	17.00°	5.17		0.50	10.75	12.80	14. 00.0 10.0 10.0	10,70	05,50	10 <u>.</u> 50	16.50 10.30	11.10	14.		40.40 13.30	12,10	00	15. 8. 8.	11,000	12.20	9.0H	13.02	ر ا
มูล	at. Bone Shank	:	ci a	9		$\sim$	<b>.</b>	ر د	<u>ٽ</u> ر	. w	6.4 6.7		C	. T	9	Ç C	? -7	6	ر. د	•	107 c	•	•	2 C 2 C 3 C	3.10	0 0 0 0	3.00	2.60	2°60 8°0	2.60	2.00	7.56
4	st. Fat Shark		•	• •	•	•	•	•	٠ ٢ ٢	• •	1.28 .58		01.1	1.20	1.05	1.	37.1	1.30	1.5 0.00	20.	727	•		1.40	1.30	02.4	1.35	1.65	1.25 1.25	1. 1. 1.5	下。 一	£9•0
I VIORIA	at. Lean Shank			٠ ٠	10 11 12	w. س	<u>ر</u> ش	09.	つん・ コー コー	3 to 0.0	3.55 1.76		3,35	3.17	3.20	3.40	7.7	3.20	1. 0.02 0.02	٠ ٢٠ ٢٠	3.57	7 • 1		4.40	3.60	۳ ۳ ۳	, <u>–</u>	3.45	۳. ورز		3.76	1.70
	Shank							a . (		် ယ	3.51		•	•	•	•		•	•	•	25.6	•		ည် ပြင် ပြင်	•	•	• •	•	•	• •	•	•
	at. bone Chuck		•			•	•	ن ند	•	ران ز	76.4													20 20 20 20 20 20 20 20 20 20 20 20 20 2	•	•		•	•	• •		•
[	tt. Fat Chuck		•	•		•	•	•	•		85.9		•		•	•		•	•	•	18-41	•		22°60 20°10	رين	∾° 0	, c	Ci.	۰ د	, ,	0	•
	kt. Lean Chuck	· (C) ·	26.10	0 0 0 0 0 0	06. 1.1	15.20	55°55'	1.0.75	99°27	10	15.05 20.71	· · · · · ·	- č	10.10	39.80	14.50	70°77 16°50	34.80	37.65	10~10~1	101 a	<b>((-)</b>	6-1.	37. 77. 70.	12.50	10.60 28.30		35.25	02°96	57. CZ	10.16	18.21
	Steer	20	r-1 (	y r	1-17	<b>ኒ</b>	9	~	ల ల	ν ΟΤ	Ing. Mean Netric Mean	<b>⊢</b> F	• tr dno.	12	13	4	75 16	17	8 8 8	2,6	Keen Keen	ပ	Group III .7	12 26	23	けいつつ	52	22	න <b>ර</b>	0E	, e	Metric Mean

1	}	-229-	
it. Bone Brisket	60000000000000000000000000000000000000	20000000000000000000000000000000000000	444444444 8888888778877
Kt. Fat Brisket	w -   -   -   -   -   -   -   -   -   -		~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~
.t. Lean Brisket	7,0005000000000000000000000000000000000	00000000000000000000000000000000000000	がい ことのようとくらる こうさい ひょう こう の で こう の の が の の で は な
Wt. Brisket	23 44 44 44 44 44 44 44 44 44 44 44 44 44	66556666666666666666666666666666666666	8228844484 8228844484 822844484 8228444 8228444 82284 82486 82486 82486 82486 82486 82486
.t. Bone Stank	88888888888888888888888888888888888888	44600000000000000000000000000000000000	22000220000000000000000000000000000000
wt. Fat Skark			44446646666666666666666666666666666666
د د		02000000000000000000000000000000000000	
Kt. V Shank	80077780007 180885000000000000000000000000000000000	00000000000000000000000000000000000000	00000000000000000000000000000000000000
Mt. Bone Chack	<ul><li>これののことをできる。</li><li>これのことをはなる。</li><li>これのことをはなる。</li><li>これのことをはなる。</li><li>これのことをはなる。</li><li>これのことをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これをはなる。</li><li>これ</li></ul>		
Wt. Fat Chuck	23.22.22.22.22.22.22.22.22.22.22.22.22.2		22 22 22 23 24 25 25 25 25 25 25 25 25 25 25 25 25 25
Wt. Lean Chuck	1 でははならならないに	25665447566459 6666747566459	**************************************
Steer No.	الم	Group V .26- 111 112 113 114 115 116 116 117 118 119 50 50 50 50 50 50 50 50 50 50 50 50 50	Fromp WI • 51 62 63 60 Eng. Mean Motric Lean
	teer Wt. Lean Wt. Fat St. Bone Mt. Wt. Lean Mt. Fat Wt. Bone Wt. St. Lean Mt. Fat Mt. St. St. St. St. St. St. St. St. St. S	No. Chuck Chuck Shark Stark Stark Brisket Bris	Steer Mt. Lean Nt. Fat it. Bone Mt. it. Lean it. Mt. It. Lean it. Fat it. Bone Mt. Stark

	M.t. Kidney Knob		7-70	00.00	7.60	7.90	2°C	- w	5,10	6.40	3.55	11.50	04-90	0%•64 01-01	11,00	13.30	11-30	08.40 04.40	11.72	,	05.30	12.40 12.10	09.01	00° d.t.	17.50	05.41	) () V V M H	(0) (1) (0) (1) (1) (1)	ن المراجعة المراجعة
	wt. Bone Flank		01.	0.5	10	010	٠ د د	01	.20	505	<b>7</b> 0°	01.	٠ ا	01.	2,5	12	010	) 5	Ci	† •	01.	٠ <u>.</u> د	10 10	0 0 0	• • • • •	ນ oʻ			<u>.</u>
(p)	wt. Fat Flark		08,10	10,50	10,10	13.70		00.50	06.50	05 05 05 05 05 05 05 05 05 05 05 05 05 0	511-110	00°75°	13.00	16.70	13.00	10.10	00° 60° 60° 60° 60° 60° 60° 60° 60° 60°	06.00	13.16		18.45	15.60 60	07.41	16.50	15.70	17. 30.	٥ ٢ ٢ ٢	15.16	ਹ <b>਼</b>
A (Continued)	it. Lean Flank		က္ ကို (	ુ િ અ	8,30	07.6		2.40	6.10	20°3	3.64	09*2	8.50	တ္ လ တ ု	, , , , ,		000		3.75 3.76	) •	8.70	7.72	7.40	°.60 00.70 70.70	, v	7.05 205		1-1 1-1	77) 71) 71
APPENDIX III	™t. Flank		16.70	19.05	18.50	23.40	10 TO	16.50	1.1.75	16.90	8.20	08,33	23,30	25,30	22.40	10.20 22.50	22,00	16.50	21.77	- - •	27.60	24. 00.00	00.53	25.20	21.60	22.75	77.25	(S)	77.07
APFE	.t. Bone Flate		2.10	1.75	2. .W.	1.95	0 <b>T.</b> 2	<b>2</b> •60	2.50	2-1:0	65.	C.C.	1.50 0.50	1.80 2.10	2.10	70°7 10°5	2°.	00.0	2.08	1	1.30	ч у у	1.75	90. در و	1.80	1,60	0 to	1.75	€7.
	Wt. Fat Plate		7.00	00 00 00 00 00 00 00 00 00 00 00 00 00	7.30	00.5	2.40		ሊ ር	7-65	3.52	ر برن برن	09.50	12.20	02.50	07.65 07.60	11.40	08.20	07.50		12.60	다. 원급	12.40	01.01	09.11 11.60	다.	10.00 17.65		₹.
	wt. Lean Plate		10.75	00 00 00 00 00 00 00 00 00 00 00 00 00	06.60	08.20	04. 80. 80. 80.	11,10	08 • 60	10 cg	1. Lo	۲ د د د د د د د د د د د د د د د د د د	10.60	08 <u>.</u> 50	00.50	ू.   	09,40	10.00 01.00	9.72 Fil.1	1 7 •	06.10	လူ ရ ၁	03.50	09°50	05.70	C5. 50			[5.*;
	.t. Flate		21.00	02.22	18,60	20,30	100° 00° 00°	21.90	18,10	01 13 00 03		•	22.60	23.00	21.30	24.00	& C	10.4.01 10.4.01	22.1E	֓֞֜֞֜֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓֓		25 <b>.</b> 00	23.80	00°68	27.72	23.70			7. C
	Steer	Group I 26	ਜਿ	// m	ケノ	w	0 6	– య	6	10 Fnc. Mean	ric	Group II .51	12	<b>در</b> ر	, vi	9 C T	⊕ . ⊕ .	20 Z	Ing. Mean	)	J. TIT dnoar	() ()	24	ርሳ <u>ር</u> ፓ./ ለ	22	∞ ( <b>0</b>	N 6	Tag. Zoen	Metric Fean

			-232-	
	t. Edaes Zrob	2482444244 2482468886 2483468888	218114884414 6888478888888	25.000 25
	Westk	enceconnonien enceconnonien	2525555555	000000000000000000000000000000000000000
	lvt. Fat Slank	2007808824479 2007808826 2007808		447898449 6478984498 6478984498
(Continued)	ht. Dean Flank		#448888888848 #4488888888	88888888888888888888888888888888888888
√ 	1. t. Flank	80888888888888888888888888888888888888		
	Mt. Bone Plate	44444444 766666676746	EDWOODS SHORTS HOW WON TO WOOD	00000000000000000000000000000000000000
	Mt. Pat Plate	44444444444444444444444444444444444444		
	lt. Lean Flate	てきるなるできるならです。 できるのでのがあるのではい		
	1.t	22.53 23.53	5.6.6.2.6.3.6.8.6.6.4.4.6.6.6.6.6.6.6.6.6.6.6.6.6.6	
	teer No.	Group IV 1.7 31 32 33 33 35 35 35 36 36 36 10 808 Yean Netric Yean	Crewy V76  11  12  13  14  14  15  15  16  17  16  17  17  18  19  19  19  19  19  19  19  19  19	Group VI • 红 522 523 526 526 526 526 526 527 528 528 528 528 528 528 528 528 528 528

	Nt. Kidney	MOO	02°57	15.60	70° 50	13.00	11.90	17.50	07.31	22.40	13.50	16,00	11.30	6.71		77 <b>.</b> 60	03•71	16,30	13.90	13.83	16.00	17.60	13.30	17.00	다. 60 60 1	₹ <b>1</b> 0.57	7.01
	Lt. Hone	71	01.	01.	νς Ο	ال ا ا	•10	.1.	•10	.10	ाः	О. Г.	60	70.		•10	<b>ा.</b>	• ሊ	.10	٠ ٢٧	ਨ 0 •	• 50	• 70	•10	01.	11.	٠ کر0
(pe	int. Pot	actor a	20.75	27.13	16.50	21,00	19.70		21,30	19.85	21.15	19.30	19.03	9.35		21.00	25,00	27.80	19.35	21.00	C3.61	18,60	17.30	23,00	21.20	01.13	9.57
. (Continued	ht. hean	AGET	10,30	11,60	02.50	05.30	10,50	10.70	01.30	10.90	00° 00	04°60	10 10	75.		10.25	10.45	11,30	0,.15	11,30	08.30	10.90	ce*50	07.60	07.50	34.40	4.53
III YIG.	: • t• • 100 P. J	A LOUK	31.70	33.70	26,30	30.70	30,30	0%. 3%	30.60	31.50	31.40	00.32	31.03	14.07		31.75	32,80	36.90	59 <b>.</b> 00	32.90	26.20	55.90	27.00	32.65	30.80	31.49	14.28
APPWDIA	it. Eune	rrane	2.50	S.50	2.30	2.50 50	3.40	2,30	1,83	2.30	2.75	2,80	2.60	1.18		2.10	2.55 5.55	2,4 0,4 0,4	2.90	62.3	2.50	2.50	2.40	2.75	2.50	50 1. KN	1.17
	i.t. Eat	TERUF	NO.	20.00	cr)	زن	1.	( <u> </u>	[-	<b>~</b>	<u>~</u>	۴,	16.22	7.36		14.40	ث	زن	~	Š	20,30	œ,	'n		(r)	0	7.61
	1-4	ets se	٦.	50°7∟	1.1	2.7	8	· - 3	0.0	C. C	3.6	C\i	12.06	ኒኒ		Ö	-7	o	į.	- = =	11.80	<u>-</u> 寸	~	i,	r.	C J	5.53
	- C-	-11-	اج ا	~	·		-	2	• نان:	~	-	0.5	H	[二]	- 4	<u>.</u> .	۱ń	$\sim$	-	0)	35.10	N	Ú,	-1	-i	CU	-
	Steer	t.		62	63	64	65	éé	67	68	69	20	2 b	ric M	TII J	2	72	52	7.1	٦/	76	77	<b>3</b> 2	79	ဓ	Eng. Tean	Netric Kean

ed) Total Bone	12 12 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	2000 00 00 00 00 00 00 00 00 00 00 00 00	28 28 28 28 28 28 28 28 28 28 28 28 28 2
A (Continued) Total To Fat Bo	3377778 33777778 33777778 33777778 33777778 33777778	000 000 000 000 000 000 000 000 000 00	104.50 104.67 104.50 106.25 106.25 106.25 105.15 105.15 105.15
PPEDIX III / t. Total de Loan	64 54 54 54 54 54 54 54 54 54 54 54 54 54	136.00 126.00 126.00 126.00 126.00 136.00 136.00 136.00 136.00 136.00 136.00 136.00	135.17 132.22 131.25 134.35 131.20 131.20 131.20 131.60 61.08
APPE Total Mt. Left Side	256.25 26	261 262 262 262 262 262 262 262 262 262	277-70 276-50 276-50 271-72 271-75 271-75 271-75 271-75
Mt. Bone Hidner Ynch	00000000000000000000000000000000000000	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	36%883888884
ht. Fat Kidrey Enob	6.00 6.00 6.00 6.00 6.00 6.00 6.00 6.00	12.20 05.40 05.30 05.30 112.30 14.20 17.20 17.20	
Steer No.	uronp 3 .20- 2 .2 3 .4 4 .5 6 .7 8 .7 9 .10 30 .3.ean Metric Mean	Croup II • 51 12 13 14 15 16 17 18 19 20 Tng. Mean	Group III • 22 23 24 25 24 25 25 27 27 28 29 30 30 30 Mean Metric Mean

ed)	Total Bone	22222222222222222222222222222222222222	12. 12. 12. 12. 12. 12. 12. 12. 12. 12.	21000000000000000000000000000000000000
(Continu	Total To Fat En	一多1031350~~~	131 101 101 101 101 103 103 103 103 103	
ਚ ਜ਼		124.10 124.10 142.45 1123.30 123.30 125.00 125.00 125.00 125.00 125.00 125.00 125.00	89. 19. 19. 19. 19. 19. 19. 19. 19. 19. 1	
- 4.	Total W.t. Laft Side	24 N D D O C O N D P O C O N D P O C O D D P O C O D D P O C O C O D D P O C O C O C D D P O C O C O C O C O C O C O C O C O C O		
	it. Eone Kidney Knob	828888888		
	ا بکر پ	25.70 12.20 12.30 13.35 13.35 13.35 13.35 13.35 13.35 13.35 13.35	07 11 22 22 22 22 22 22 22 22 22 22 22 22	
	Steer. No.	でよるろれどらて8908m	Group V .26  11  12  13  14  15  16  17  18  19  19  10  10  10  10  10  10  10  10	See a

(ed)	Total Poro	0.00	05.01	10.70	36.67	34.50	1.6.65	30.00	30.60	40.00	59·th	00°36	38.38	17.38		-		37.1.5	• (\)		(:)	~,	- +		15		~
(Cortinued)	Tough Mak			150.05T	126.75	124.00	126.60	157.30	161.60	153.55	116.85	127.33	141.14	64.01		136,000	153,05	163.80	133.85	134.55	150.60	137.20	142.55	1148.85	0.7 <del>1</del> 111	17. HI	65.73
OIA III AIC	Total	i.	ر. ان ن	187,30	(C)	S. 35	7. CO.	ਹ <b>਼</b> ਹ	() () () ()	で ()	12.0	3.92	•	78.5		J.	,	168.75	ن	_	π,	Λ,	·	ری	-	6	ω̈́
ei ei	10 ts] . 0.	ן ה ט	この。ないの	362,10	0/・///////////////////////////////////	TO OF	365.00		の で で で で で で で で で で で で で	349.70	365.10	00.100 100.000	357.82	162.00		310.55	362,90	374.10	٥٢. ١٠	360.20	358.60	375.65	DY. LYK	360.25	ው ነሳ ነሳ ነሳ	360.13	163.32
	orie Fr	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1.10	ن ٦•دن ۲-دن	03.0	1.20	, C?•u	0.4.0	0.7.0	0.0	<b>1.</b> ⊗	0.80	0 V O	277.0		వ <b>్</b>	1,00	09.0	03.0	0.30	0.00	1.00	03.0	05.0	0.50		0°36
	3at	11.00 H	12.6		C		-	Q	7.7	١,٠	SO.	N.	$\vdash$	0.1	01-1-5g"		(1)	~	œ,	2,5	•	6.7	() []	6.1	J.,10	NO.	6.62
		Group VII 70	15	62	63	150	Li N	92	67	<b>6</b> 00	69	C2	Ing. Mean	Metric Mean	Proud VIII 1.	7.7	2)	73	77	75	()	77	- S	37	CS	Sng. Mean	Metric Mean

	12th Rib Jore	whaawwawaman whoomocoonbo	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	5 W 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
Jeth Ri	1000	7.6.6.2.4.6.4.6.2.2.2.2.2.2.2.2.2.2.2.2.2	65.000000000000000000000000000000000000	44466446664666666666666666666666666666
10-11th, 3	12th Rib Lean	44644444444444444444444444444444444444	444444444 6464666666666666666666666666	
6-7-8th, 5-	CUE o	66777886666669	444444444 7068888768887	986866688888
	6-7-(4h Pat	00000000000000000000000000000000000000		4 w w 4 4 w w 4 4 4 4 4 4 4 4 4 4 4 4 4
	6-7-cth Lean	CACACACACACO CACACACACACACA CACACACACACA	70000000000000000000000000000000000000	VVOONNW-VVNNO UNUUNUONOONOO
	5-10-1135E Fone	44444444444444444444444444444444444444		
PART OF ETT	5-10-1148 Fet	~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~	4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	
SET Edd	S-10-11th Lean			44444444444444444444444444444444444444
	Steer No.	Group 1.20 23 44 55 66 7 88 9 10 30 Eng. Mean	Group II •57 11 12 13 14 15 16 17 19 20 3cg. Mean Metric Mean	Oroup III • 22 22 23 24 25 26 27 26 29 30 30 30 Metric Mean

	12th Rib Eore	00. 00. 00. 00.		\$	SHOUSESHE TO
	12th Rib Fat	2.1.4.0 2.1.4.0 2.1.4.0 1.4.0.4.0	14444444 2000000000000000000000000000000	44444444 668668774687	
	12th Rib	1140 001 002			aamaahaaaa Axoooooonnar
tinged)	bh 6-7-rth Sore	4.00.1.00.00.00.00.00.00.00.00.00.00.00.0		~ • • • • • • • • • • • • • • • • • • •	
	6-7-3 Fat	8889 4400	innnama Seesesines	HENTHERMINER CANDONNERS CANDONNER	
T PPST LTX T	6-7-5th Iean	м.м. 64. 64.00 64.00 64.00	00 W W W W W W W W W W W W W W W W W W	ト <b>ケ</b> ケケル 3 3 7 3 7 1 7 8 8 9 9 7 9 7 9 7 9 7 9 7 9 7 9 7 9 7	またなるのでででで ごのものはありもののでは
	5-10-11th Bone	7000 1000			444444444 20002200200000000000000000000
	9-10-11th Fat	n'3 w= ñ'58€		2002 W C C C C C C C C C C C C C C C C C C	Nawaznannem SVSSSSSHAZEH
	9-10-11th Iean	1.25" 1.25" 1.60 5.10	12/2002/1027 12/2002/1027	0.888888888 8888888888	MARCECAGESES MARCECAGESES
	Steer No.	ri ri	24 36 37 38 39 10 10 10 10 10 10 10	V . 26 1.1 1.2 1.2 1.4 1.4 1.6 1.6 1.6 1.6 1.6 1.6 1.7 1.8 1.8 1.8 1.8 1.8 1.8 1.8 1.8 1.8 1.8	Group VI PROPERTY SAN

	}	1														•	-2:	39	-								
	12th Rib	90.02.	03•	٠ ا	01-	٠ بر	C9 <b>•</b>	0,	0;	(T)     	<u>ල</u> ]	} <del> </del>	's 0 0	7,•		Ç L	5 , € 14 –		() •		A C	(.C.+	O (3)			= · ·	67.
	1255 Tib	13.0	2.50	и\ -) -d	ይ) ተ! (ህ	Q € €	(V)	ጥ ው ህ	3.60		ر روز روز روز	0 0 0 0				() ( E) (	[] } [] (]		۵۱ ( ۱۰ • • • • • • • • • • • • • • • • • • •	) (4) (4)		- JT • 2	[] []			· (	ुन्•्न
	12th Rib	The state of the s	20.57	) ) , ,	(A)	ા.?ે	61.5	2.30	2.60	्    	٦٠ ١٠ ١٠	5.00	3 <b>7</b> € (∃ ∪ .∪	<u> </u>		164 (0 164 (0 164 (0	[] () 	(;;;• ⊣ (	τ <b>1</b> , ν (	ir (C	્ર • • •	\\ \\ \\	(T• 7	9. ; d			``
Contineed)	13 )-1-9	HLION	1.90	2.20	1.70	1.70	と、 (v) (v)	S.10	© H		2.50			ر د •		Ç.	o (0 - 1 \ - 1 \	ું - - -	) ( V		] \  -  -  -  -	15. T	ار انج اح			ન દ • •	•
) pa est	(-7-5th	3 3 3 4	5.10	100°	_1 • → Γ./	<i>ກ</i> ໄ ວິ	η <b>)</b> (1) (1)	ኒ/ O E-	С У.,	(급) (급)	L'\ ( d\ ( L'\=		ی رد. - از د م رد.	) 1		O <b>\</b>	61 200 201	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	ν.÷ π/ (	- - - }\	7) ( 3) () 1)	) • <del>• •</del> •		O (	92. 4	) () -}-	ر • باد
T MICHE SY	6-7-7th	ii da T	7.20	: ਜੂ	6.50	<b>့</b>		ω <b>\</b> 0 <b>-</b> υ <b>-</b>	6+ <del>1</del> -5	ු දු	[  		٠ - ا	ς: • •		C (	1 V V	ο \ ( , \ ( ) \ ( , \ ( , \ ( , \ ( ) \	30. 0		် ဦ ပ	() () ()	ري• <i>ز</i>			਼ੋਂ - -	
	7-10-11 th	pone	•	•	•	•	•	•	•	•	•	Ç.  -  -		) 0		() ( - ; - - d r	•	•	•	•	•	•	•	•		•	•
	)-10-11th	7 . T	100 V	() เมื่ ไปไ	л <b>,</b>	л, С.	ይ \ ይ \ ይ \	ኬኒ ኬኒ ኒስ	03•9	٥ ٢٠٠	00° 00°		다 <b>, c</b> 이 년 이 년	( · )		か/ヾ う.	11	•	(	۱ د٠.	٠,٠,٠	' '	ا ۱۰۰	w.	٠.١	_ `, î	•
	11 (	Lean	• ັ• • ኒ.՝	O. W	5.70	CX•1	6.75	0¦7•‡∤	0 m	М. С.	パ <b>ノ</b> .	8	::)	7	1.01-1.25#	ក្ស ទីវ	ر الم الم	3. A)	Ç; ; 5√; 14√; ;	); ); (3)		ົວ. ວັ	L <sub>ር ነ</sub> ር ነነ ይ ነ .	7,•€©		(\) \ -\$- -\$ -	· · · · · · · · · · · · · · · · · · ·
	Steer	Cacha WTT	4	62	63	73	59	9)	13	<b>9</b>	69		ď,	umar otagen	Group VIII	77	04 C	~ <u>} -</u> ;		''- <b> </b>	0 1	- (	(1) <b>(</b> **	υ. ί -	C .	4 80 T	Stric Form

		-240-	
Psoas Major	00000000000000000000000000000000000000	\$2000000000000000000000000000000000000	20101110 2000 2021888891508
Supra- soinatus	2.15 1.95 2.25 2.25 0.94		
Tricens Erachii	1.30 1.30 1.30 1.65 1.10	10000000000000000000000000000000000000	8058884486555
Rectus Fenoris	00000   6001   00000   6001   000000   6001	ENERGERESSES	20000000000000000000000000000000000000
NTSOLE AND BAR m- Bicens sus Femoris	20000000000000000000000000000000000000	03486568669868 03486666669868	20000000000000000000000000000000000000
III C MTS Semimem- brancsus	2.00.00 2.00 2.00.00 2.00.00 2.00.00 2.00.00 2.00.00 2.00.00 2	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	4004WW4WWWW0 00000H00H00H
Seminemb.	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	70.00.00.00.00.00.00.00.00.00.00.00.00.0	60000000000000000000000000000000000000
Quad <b>ri</b> ceps		01000000000000000000000000000000000000	0,000000000000000000000000000000000000
Semiterd- inosus		12222222222222222222222222222222222222	
itee Fo.	Group I .76- 2 3 4 5 6 7 7 10 10 Eng. Mean	Group II •51 12 13 14 15 16 17 18 20 20 4ng• Wean Wetric Mean	Group III .7 22 22 23 24 25 26 27 27 28 29 30 30 30 Frg. Mean

	1 1		-241-	
	Psess Najor		832000000000000000000000000000000000000	waaraaaaaaaa wee maaaaaaaa
	Spire tes	00000000000000000000000000000000000000	88888888888888888888888888888888888888	
	Triceps Erechii		wwommocmocho Hoginochocho Www.minwinwino	5.5.46.5.6.6.6.6.6.6.6.6.6.6.6.6.6.6.6.6
ned)	Rectus Remoris	10000000000000000000000000000000000000	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	
C (Continued)			884966666666666666666666666666666666666	Woden were well to
III XIQ.53	ិត្តក្រុក ឯកឧកស	ママンション・マック・マック・マック・マック・マック・マック・マック・マック・マック・マック		30000000000000000000000000000000000000
A. A				
	્રેપ્ટલેજ લ્લુક	01/444440000000000000000000000000000000	WG67277760 PT 2020700700 PT 2020700700	1.001,414,414,60 4.646840488
	(%)		Saawaaawaaaa gywoogkookoo gywoogkookoo	
	eer o	oroup 1v 1. 32 33 33 34 36 37 38 39 10 Metric Kean	Group V .26  11  12  13  143  144  155  165  165  165  165  165  165	Tales of the state

	13	្នាះ		lin lin	八 八 八 八	3.33	2.15	05 <b>°</b> 2	2,00	2,30	S. S.	0 بار بار		C1	0 F		5.10	2.70	10 C-	S. S.	2.60	2.1V	ርረ ም. ፓ/	2.80	2°.57	2. J.S	Ç;•3	1.18
	-eating	soir etus		S. C.	ر. کار در	01.3	ኒ የ የ	ው ው	(C)	() () ()	ኒ.ነ (ና)	( ) ( )	2.50	710	1.1		2.20	2,30	01°2	8 안.	5.40	03.5	2.1:0	8.30	S. 30	C ( )	C	1.C7
	rice	Frachi		Li \ Li •	00•9	о. С.	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\		ນາ (ຕ	() () () ()	(C)	いり	\(\lambda\)	(L)	2.30											전 (연 (연		
(Pe	Rectas	Femoris		<b>.</b>	r',	(J	•				. (7	c,	· C ·		-3		9 <b>.</b> 6	۵. ار	ত জু	િ જ	5.10	ሊ	<u>ر</u> ن • 2	00.0	ربر. ه	3.10	30.	1.37
C (Septinged)	6 100 n	Ç		다.	ፓላ ሌ!	10.70 10.70		() 	10.01	C2.	S.	100 100	(S. 5)	(C)	ا ا ا		つ い い	7.80	О М И,	છ. જે	6.50	7, 1,7,	00°9	6.10	€.00 •	6.17		2 <b>.</b> 60
III XICE	Cerimon-	इसक्र अध्यत्		7.70	09•9	00 <b>°</b> 9	000	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	. (C) . ນ .	(N. 1)	C) U,	ڻ. وي	10 10	1000	2.00		C (*)	6.70	6.10	က် တ်	6.E0	6.60	ج 99.	6.45	5.7.2	7.30		3.77
	Sumita onb.	+ Adduct.			( )	_	. ( )	Lı	. ( \	05.70	. 1	٠,			7:•05											() ()		رن س س
		sioojajan*		ር	6.10	ਨ• • •	6.10	CO <b>°</b> 9	01. V	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	C3•2	( <u>.</u> 75	ر. بر	6.37	ි. වේ		•	•	•	•	•	•		•	•	ኒ./ (~ ሀ ነ	•	•
	Cantibero-	incesse	17-1.65"	た) で)	). ().	•	•	•	•		•	•		200		.01-1.25"	γŌ	'nĴ	~	Ç	4	7	୍ଦ	$\sim$	Q	6) D.	ربا	[
	Steer	*C.	7	<del>U</del>	<del>2</del> )	<del>(</del> 3	ائد دیلا	n NG	99	<i>(</i> , )	er.	59	02	Jng. ∑ean		1 1111 1111	77	72	73	-1	ር~ ፓ/	92	77	r- ω	42	ධ	žai.	Natric Mean

_	•
Continued)	4
Ò	Ś
~	ź
-	:
7	4
٠,-	ŀ
-	>
(	•
7	1
	′
ζ.	)
_	-
Ċ	١
_	•
_	1
⊦-	١
<b>⊢</b> ⊢	1
١.,	
ľ	:
Г.	1
	3
<u>.</u> -	
Çτ	1
1	i
á	•
	•
• =	i

, ,				
( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( )	Humerus	20000000000000000000000000000000000000		41010000000000000000000000000000000000
	Scapula	\$388900K08XX	00550000000000000000000000000000000000	444444444 73674687888
0 ! K ! &	+ Fibula	00000000000000000000000000000000000000	444966666466 64496666666666666666666666	28.30 28.30 29.10 29.10 29.10 10.10
	Femur		HWWWWWWWWWWWWWWWWWWWWWWWWWWWWWWWWWWWWW	20000000000000000000000000000000000000
25.00	+ Ulna	2000 2000 2000 2000 2000 2000 2000 200	119999999999 200000000000000000000000000	440040040440 86886868666
	L. Dorsi	# 50 H H H H H H H H H H H H H H H H H H	00000000000000000000000000000000000000	6-1-00-00-00-00-00-00-00-00-00-00-00-00-0
\$ 0 10	No.	3   26-   2   2   26-   3   4   26-   5   6   26-   6   7   26-   7   8   20-   8   20-   9	Croup II .51 12 13 13 14 15 16 17 18 19 20 Eng. Wean	Group III .7 21 22 23 24 25 26 26 27 29 30 5rg. Vean Netric Mean
		,		

C (Continued)	Humerus	35	2,10	ν ν Θ Μ	20.0 20.0 20.0 20.0 20.0 20.0 20.0 20.0	2.15	2 <u>.</u> 15	2.10	0.30	101		3.50	3.20	3.70	w w 30,00	w. v.	ر م م	03.60	00°		2.55 2.55	<u>ر</u> د		3.60	00°†7	2.60	en to	○ C ↑₩ ↑ H
DIX III	Scapula	1.52	07.	1.05 1.05	1,60%	1.35	۲. گرار	다 다	1.30	19.		2. 2. 2.	2.5 2.10	2.20	2.05 2.05	2.33	ν ς ο ω ο ν	2.5	1.04	,	2.50 2.10	2000	2.75	2.25	<b>เง เ</b> เง เ เง ห	0 0 0 0 0	2.k	0 (V V C V H
APPE	Tibia + Fibula	2.30	رب ا السيار السيار	۲. در در در در	7. Co.	2.00	20 10 70	2.10	1.85	35.		3.60	3.50 3.20	2.50	9.10 2.10	G. (	~  	3.30	7.00 1.16	,	w v w v	2.00	02.0	3.10	w w	 	<u>्</u> र	1) (Q 1_J 1) (T
	anweg	3.05	W.C.	2,00 3,00	, ~	2.70	0 0 0 0 10 10		2.60 2.60 2.60 2.60 2.60 2.60 2.60 2.60	1.30		4.15	10°17	1, 80 1,		1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1	1 0 0 0 0	1.20	200 100 100 100 100 100 100 100 100 100		4.55 3.75	0.5°	1 to 1	05.	7. 01. 01.	. w.	ر س س	ş₩ Ş₩ AH
	Radius + Tha	2.20	•	•		•	•		2.10	• •		₩. 127.	0.7°	2.60	2.5 2.01.	3.40	3.40	5.00 C	7.00		•	•	• •	•	•	• •	• 1	↑ (d)
	L. Dorsi	1-1.25" 8.10	0. 0. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.	20. 0.63	8. 60.	స్త్రి	ణ	3	000	4.11	.50"	•			12.70		- <del>-</del> ~	• •	6.10	75m	13.67. 7.00.00	12.70	12.15	13.15	12.25 10.05	12.65	12.30	ン の い い い い い に っ い っ い っ っ っ っ っ っ っ っ っ っ
	Steer No.	○ H	(M) (						1:0	ug	Group V .26-	4	77	177	77 70 70 70	77	100		Mcg. Mean Netric Mean	덗	ር የ	1 <b>(</b> ),	±¥ Ľ\ť	\X	<b>γ</b> ν η <b>ς</b> - α	ን ዕላ አውነ	Оў.	Ing. Mean Metric Mean

(Jortinged)		ingerus			C.W.	ر. در•	3.10	J. 60	2.75	65.50	w) w; p;	다. 다.	3 - 1 - C	26.2	7:-		2.70	2.50	ر د د	07.0	000	01.01	0.00	0.0	\_``\ -`↓ • ('\	2.10		1.32
PRAIDIX III C	1	Scanนใจ		5.30	2.00	07 در• در•	() 라	2.50	1.70	€. 1	2. 1.	01.3	년 ( 8	20°Z	. ` ` ` · ·		•	•		•	•	•	•	lo∖ Ov ed	•	• 1	T. 3	رن درن •
Sat Table	Tibia	+ Fibula		∑ • •	5.60	ි. ල	S. 80	00°	?. .50	ूर <b>े</b> इ.स	ر د د	٠. د.	00.6	( ) · ;	1.32		C) L J('	2.10	ان م ان	ი ლ.	10 0	3.00	3.12	2. 2.	3.10	ر. د اند د اند	۲. د.	7.52
		ក្រុមពេល		%. €. €.	D. 65只	3.10	೧೫ <b>.</b>	05.4	3.10	00.0	3.40	1.17	1,00		93.• ∃		7.00	w N	0. 0.	1 <u>0</u>	00.4	05.0	07.7	ህ . ማ	1.35	3.70	3.57	1.00
	Redins	+ เกิกล		ኒ ኒ •	س رہا ہیں	2.80	2.70	2.60	2.70	2.05	3.30	3.10	2.73	2°-0	3 <b>.</b>		2.00	<b>ሪ</b>	<b>6</b>	3.20	05.3	0 <u>1</u> -2	3.50	2.20	₩. 20.	2 - 2 - 3	5	J. 56.
		1.3	6-1.00	ری د	2.7	3.1	$\mathbb{C}$	(J)	7.5	2.0		2.2	r1	1 -1 (	rv,	()	-12	1.2,15	11.25	ኒ ር	12.10	in 라 터	10.05	12.30	11.70	12.15	( ) ( )	กัก นา
	Cteer		Croap VII .	<b>U</b>	62	(3	<del>[]</del>	65	9	67	68	ίÿ	70	2	Metric Mean	Croug VIII 1	77	72	73	71,1	۱ ۱ ۲ -	<u>ر</u> ر	27	36	S.	င့	(G)	Metric Mean

		1											-2	46-												
	Temerness Stoor		07.60	10,82	(\) \(\) (\) (\) (\) (\) (\) (\) (\) (\)	100 100 100	0.5.65	10.07 20.	11.37	10.01 10.03 10.03	\(\frac{1}{2}\)	06.94 12.30	10.07 04.00	10.17	07. 10. 18.	11,10	0% . WS	10.44 14.60		10.77 08.20	で で で で で で で で で で の で の で の の の の の の	11. 10. 10.	08.31	20·20	08.16 20.2	50
	Indian ness Tooks Pond		7 (S) V	6.37	© \ D \ O \	0.	7.12		) S	<b>い</b> の	×:	ာ့ ကု	رن بر س	ر. ا	a) Ca	AD M	7.0.7	( (		6.87 7.12	<b>~</b> % % % %	6.62 5.62 5.62	7.62	2. 2. 2. 2. 2.	7.37	\$ C
	iat oom prot breis	u L	ು ದು ಗ (v	(A)	20°	コ・レン	26.11	٠ - - - - - - - - - - - - - - - - - - -	- S.	21.12		64. 67. 7.	ლა და და და და	) - 1   1   1	26.7	625 1000	2 B.	MW MW		<b>ນ</b> ເພີ່ ທີ່ເພື່	2 <b>9</b> 3	24.7	26.5	5.4.5 5.4.5	27.00 20.00 20.00	26.50
min problem	100 M	ł	1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -	1.55.1.	7,17 0,0 0,0	7. L. O. C.	رم: دم:	0 9 9 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	100 160 160	73-15. 123-15.	(_	-, CA	W.	, מו	いんへつ	C	-! (~\l		,	517 878 1. 8.	7.00 2.00 2.00 2.00 2.00 3.00 3.00 3.00 3	712°2	- 600 - 600	0 00 0 00 0 00 0 00 0 00 0 00 0 00 0 0	ज्यातिक स्थापित	200mm 20mm 20mm 20mm 20mm 20mm 20mm 20m
PER CAR TOE	K+ ppa rot. bosis	0	131.2	3.691	C-Σ V() V M - Σ	의 급 의 () 의 () 금 급	165.3	9 = L W L W	i ci	126.71	α 6	- (조) - (2) - (2)	יולר הייר ממ	0 00 c	153°-7	166.9	168.0	から ない。 ない。		158.9	a o v · v	1,5,6	0.0 9.0 10.0	ે જે. • જે. જે.	1,40 0 1,40 0	154.97
TIMES AT RICAG	May Man		4000 4000 4000	3500	رب می می رم	ひ と さ さ さ か	3100	₩ ₩ ₩ ₩	3213	3508 3368	0	3181	9508 - 1366	300 to 00 to	3335	3451	3555	3347		3253 3157	0 60 0 70 10	322¢ 3001	3000 2000 2000 2000	9000 0000 0000	TU TE	3 03 7 11 7 10 7 10
G.	Foistare	(	a u	$\mathcal{C}_{\mathcal{A}}$	CIC	$\gamma \wedge \epsilon c c$	$(\mathbf{c})$	€.	1 (1)	73.67	Ú	$i_N \odot$ .	ゴッ	`	( (_)	$\alpha$	71.17	46.69 46.69		70.32 71.25	$\omega_{\omega}$	C 11	100 C	` α		\ <u>\</u> \
	1 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5		7.71	ر 0. 0.	~; c ~; c	5.40 2.40 2.40 2.40 3.40 3.40 3.40 3.40 3.40 3.40 3.40 3	3.70	ر ان ان	1 T	رد 109	ن بر د	$\sim$	<b>23.</b> 28. 00. 00. 00. 00. 00. 00. 00. 00. 00. 0	7.14	0 17.0 17.0	74.12 12.12	せる。	 		07.63 08.30					• • • •	
	Frotein	::0: ::0:	$\sigma$	. C	r+c	5 ⊢	C:	65 10 10 10 10 10 10 10 10 10 10 10 10 10	; r-41	₩ ₩ ₩	15.5°	98	17.			•	20.05	21°51 9.62	-1,00	20°58 20°12						
	Steer 5 No.		<b>⊣</b> ∾	m	<b>-</b> :∄	AVC	<b>C</b> -	<i>ن</i> دی	H	Brg. Mean Metric Mean	Group II 500-	구 C2 구 <b>근</b>	W-:	† ነገ \ ት	17	α) ( Η r		Eng. Mean Metric Mean	3roup III .76	53 53	23 12	N 00 N/00	222	) () ()	() () () () () () () () () () () () () (	, ; ; ;

irned)	
(Cont.	
Ν	
A PERMOTIC	

ļ	)	i	· ·	
	Terderness Stear		00.00000000000000000000000000000000000	000 11100 11000 11
	Tenderness Taste Parel	00100000000000000000000000000000000000		
	ier pom prot. besis		60000000000000000000000000000000000000	まとしているとののない。 こうこうしょう こうしょう こうりん できる できる かいごう こうしょう こうしょう こうしょう こうしょう こうしょう こうしょう はいい ない こう ない こう ない しょう
กหคิด )	Rad Ipm fresh basis	577 577 577 577 577 577 577 577 577 577	は の の の の の の の の の の の の の	0.000000000000000000000000000000000000
(Jont1	K+ ppm prot. basis	055 055 065 065 065 065 065 065 065 065	0.000000000000000000000000000000000000	
TO BEAU	K+ ppm fresh basis	3318 3141 3141 3151 3125 3125 3126 3726 3274.7	3357 32515 32515 32511 3503 3173 3173 3173 3173	
	3 Moisture	00000000000000000000000000000000000000	72.80 65.30 65.30 70.11 68.08 70.17 70.17	77. 66. 25. 67. 67. 67. 67. 67. 67. 67. 67. 67. 67
	Fat		4.50 6.34 6.34 7.50 7.50 7.50 7.60 7.60	SESSENT SESSEN
	Pretein		20.22 20.12 20.12 20.12 20.05 11.05 20.05	25. 25. 25. 25. 25. 25. 25. 25. 25. 25.
	Stear No. 9	いころろりとらいのはい	Group 7 .26 11 12 12 13 14 14 15 16 17 16 16 16 16 16 16 16 16 16 16 16 16 16	Eng. Month of Manner

	}	1											_										
	Tenlerness Strr	ارار ابار ابار	٠ ١٥٠ ١٥٠ ١٥٠	09.145	17°07		() ()	0; •0 •0; •0; •0; •0; •0; •0; •0; •0; •0; •		-1 / (- -1 / (- -1 / (-)		10.00	10.20	10.60	رين 13	07. 07.	27・30	00°27	12.05	13.51	10.37	10.12	7.50
	Terdemess Terte Pani	13°9		() • <u>.</u>	_t_ 000 000 000 000 000 000 000 000 000		6.15	(C)				•	•	•	•	•	•	•	•	•	7.07	•	•
	]್ಷೀ ಗ್ಲೀಪ ೧೯೮೭ ಕ್ರಿಂಡ	r-4 r ( - \ ( - \	. α ν ην - Ο	0.00	61 0 (C) 71 (C) 71	ή (Ο ) (Ο ) (Ν	0 v (	Li ja T Cu (		20°53			- `	1	W/	ON	(T)	NO.		V:	31.6	$\circ$	$\mathcal{L}$
nged)	्रेडन हो आ क्रिक्टो, वेह छोड		かいこと	(V) (V) (上)	Մ <b>ի</b> « «. Ն Հ. Ն	<b>1</b> 0	- * · · · · · · · · · · · · · · · · · ·	00 = 0 = 0 0 \ n = 1   0		1 (N) 1 (N) 1 (N)		で <u>。</u> (1) (1)	0.177	578.lı	712.6	623.7	573.0	57.7.5	546.7	٢٠,٦,	618.7	さ・ここ	57.7.4
IV (Corti	it jes mot. nesis	[ ] [ ] [ ] [ ] [ ] [ ] [ ] [ ] [ ] [ ]	100.01	10 10 10	01- 02- 12- 12- 12- 12- 12- 12- 12- 12- 12- 1		17.	mi c Lila Lila Fila	13 h	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\		<b>2</b> 0.	ου -2 <b>1</b>	$\mathbb{C}$	<b>ሮ</b> ፕ) ኣ() .	0	-7	닸	$\Gamma$	<u>.</u>	161.1	3). 3)	ツ
ATCE SA	Tt Jan Roese	ω. σ. ε ε	3318 3318 3318	330F	(~ L เกษ เกษ	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\		31:50 5:10 6:10 1:00 1:00 1:00 1:00 1:00 1:00 1		) a, ' ( ' ( ) ' ( )		1120	725	3501	5 당 6	3056	-ក្ ម <sub>្ព</sub>	3343	3361	3261	3210	M3544	3343
	S Moisture			~	CU (	, 😘	!	t~- (	- (	, (\cdot		ੰ	H	زين	C J		ਂ	,	۳,	ં	67-25	، ڈ	C.Y. 87.1
	43 (3) (5) (5)	UN I	17. S	( C.)	CAR	. در)	( '	رن ا	1.	ا بما ر		(; )	10	· ·	o.		•	· .	O		12.52	5 5	26.3
	Frotein	( - <del>, </del> -	•	C	ri.	, (3	, 1	٠ (	-	• • U\	r'	$\sim$	Cu	r-!	r 1		-:	$\sim$	Ü	C4	15.46	-1 (	
	J. 0.1	3rous 772 - 78 61	N (C)						ż	Tetric Meso		L'-	72	73	11:	<u>ኒ</u> ስ [ ~	, i –		သ 1	52	S ;	,:,	Metric Mean

-5

. .

## List of Variables on Data Cards

1.	Av. 12th rib fat thickness	51.	Wt. of un-trimmed lean outs.
2.			Wt. of belly cuts * KK,
3.			Wt. of trimmed lean cuts.
4.	Carcass conformation		Wt. of external fat trim.
	Hind Quarter conformation		Wt. of K.K R.S.
6.	Fore Quarter conformation		Wt. of boned, trimmed RLRC.
7.	Maturity score		Cutting loss - R.S.
8.	Marbling score		R.S. weight.
9.	Carcass grade		Wt. un-trimmed round- R.S.
10.	5th Thoracic - 4 in.		Wt. trimmed round - R.S.
11.	5th Thoracic - 8 in.		Wt. fat trim round - R.S.
12.	5th Thoracic - 12 in.		Wt. un-trimmed loin - R.S.
	5th Thoracic - av.		Wt. trimmed loin - R.S.
14.			Wt. fat trim loin - R.S.
	8th Thoracic - 8 in.		Wt. un-trimmed rib - R.S.
16.			Wt. trimmed rib - R.S.
17.			Wt. fat trim rib - R.S.
18.			Wt. Un-trimmed chuck - R.S.
19.	_		Wt. trimmed chuck - R.S.
20.	11th Thoracic - 12 in.		W. fat trim chuck - R.S.
21.	11th Thoracic - av.	•	Wt. shank - R.S.
22.	1st Lumbar - 4 in.		Wt. brisket - R.S.
23.			Salable yield - R.S.
24.	1st Lumbar - 12 in.		
25.			Total internal fat trim- R.S.
<b>16.</b>			Wt. bone - R.S.
	4th Lumbar - 4 in.		Wt. plate - R.S.
27.	4th Lumbar - 8 in.		Wt. flank - R.S.
28.	4th Lumbar - 12 in.		Yield round - R.S.
29.	4th Lumbar - av.		Fat trim round - R.S.
30.	6th Lumbar - 4 in.		Bone round - R.S.
31.	_		Yield loin * R.S.
32.			Fat trim loin - R.S.
	6th Lumbar - av.		Bone loin - R.S.
	3rd Sacral - 4 in.		Yield Rib - R.S.
	3rd Sacral - 8 in.		Fat trim rib - R.S.
36.			Bone rib - R.S.
37•	3rd Sacral - av.	87.	Yield chuck - R.S.
38.		88.	Fat trim chuck - R.S.
39.		89.	
	5th Sacral - 12 in.	90.	
	5th Sacral - av.	91.	
42.		92.	Bone shank - R.S.
	Av. at 8 in.	93.	Yield brisket - R.S.
	Av. at 12 in.	94.	Fat trim brisket - R.S.
45.	Depth of 10th rib.	95.	Bone brisket - R.S.
46.	Depth of 12th rib.	96.	% protein in L.D.
47.	Depth of Brisket.	97•	Yield Plate - R.S.
48.	Length of round.	98.	Fat trim plate - R.S.
49.		99•	
50.	Taste panel-flavor	100.	Yield flank - R.S.

```
153. Wt. fat 9-10-11 rib.
101. Fat trim flank - R.S.
                                       154.
                                             Wt. bone 9-10-11 rib.
102. Bone flank - R.S.
                                       155.
                                             Wt. 6-7-8 rib.
103. Wt. Rump-less round - L.S.
                                       156.
                                             Wt. lean 6-7-8 rib.
104. Wt. Lean rump-less round - L.S.
                                       157. Wt. fat 6-7-8 rib.
105. Wt. fat rump-less round - L.S.
106. Wt. bone rump-less round - L.S.
                                       158.
                                             Wt. bone 6-7-8 rib.
                                             Wt. 12th rib.
                                       159.
107. Wt. Rump - L.S.
108. Wt. lean rump - L.S.
                                       160.
                                             Wt. lean 12th rib.
109. Wt. fat rump - L.S.
                                       161.
                                             Wt. fat 12th rib.
110. Wt. bone rump - L.S.
                                       162.
                                             Wt. bone 12th rib.
                                       163.
                                             Fat measurement A.
111. Wt. intact round - L.S.
112. Wt. lean intact round - L.S.
                                       164. Fat measurement B.
113. Wt. fat intact round - L.S.
                                        165.
                                             Fat measurement C.
114. Wt. bone intact round - L.S.
                                       166.
                                             Fat measurement D.
115. Wt. loin - L.S.
                                       167. Fat measurement E.
                                             Fat measurement F.
116. Wt. lean loin - L.S.
                                       168.
                                       169.
                                             Fat measurement G.
117. Wt. fat loin - L.S.
118. Wt. bone loin - L.S.
                                       170. Av. measurements D, E, F, G.
119. Wt. rib - L.S.
                                        171.
                                             Av. all fat measurements.
                                       172.
120. Wt. lean rib - L.S.
                                             Loin eye area.
                                              Taste panel- Juiceness.
                                       173.
121. Wt. fat rib - L.S.
122. Wt. bone rib - L.S.
                                       174.
                                             Wt. semitendinosis
123. Wt. chuck - L.S.
                                       175.
                                             Wt. Quadriceps
124. Wt. lean chuck - L.S.
                                        176.
                                             Wt. semimembran. + adduct.
125. Wt. fat chuck - L.S.
                                        177.
                                             Wt. semimembranosus
                                       178.
126. Wt. bone chuck - L.S.
                                             Wt. biceps femoris
                                             Wt. rectus femoris
127. Wt. shank - L.S.
                                        179.
128. Wt. lean shank - L.S.
                                             Wt. triceps brachii
                                        180.
129. Wt. fat shank - L.S.
                                        181.
                                             Wt. supraspinatus
                                             Wt. psoas major
Wt. longissimus dorsi
130. Wt. bone shank - L.S.
                                        182.
131. Wt. brisket - L.S.
                                        183.
                                        184.
                                             Wt. L.D. 9-10-11 rib.
132. Wt. lean brisket - L.S.
133. Wt. fat brisket - L.S.
                                        185.
                                             Wt. L.D. 6-7-8 rib.
                                             Wt. 1.D. 12th rib.
134. Wt. bone brisket - L.S.
                                        186.
135. Wt. plate - L. S.
                                        187.
                                             Wt. L.D. loin.
                                        188.
136. Wt. lean plate - L.S.
                                             Wt. radius and ulna.
137. Wt. fat plate - L.S.
                                        189.
                                             Wt. femur.
                                        190.
                                             Wt. tibia and fibula.
138. Wt. bone plate - L.S.
139. Wt. flank - L.S.
                                        191.
                                             Wt. scapula.
140. Wt. lean flank - L.S.
                                        192.
                                              Wt. humerus.
                                              Est. % Retail yield (Briedenstein)
                                        193.
lbl. Wt. fat flank - L.S.
                                              U.S.D.A. Est. % RLRC.
142. Wt. bone flank - L.S.
                                        194.
143. Wt. K.K. - L.S.
                                        195.
                                              Est. RLRC (Brungardt & Bray)
144. Wt. fat K.K. - L.S.
                                        196.
                                              Est. Lb. lean (Cole)
145. Wt. Kidney - L.S.
                                        197.
                                              Est. Lean (H & H).
146. Wt. left side.
                                              Est. Fat (H & H).
                                        198.
                                              Est. Bone (H & H).
147. Wt. total lean - L.S.
                                        199.
148. Wt. total fat - L.S.
                                        200.
                                              Taste panel - tenderness.
149. Wt. tetal bone - L.S.
                                        201.
                                              Warner-Bratzler shear tenderness.
150. Cutting loss - L.S.
                                        202.
                                              % Ether extract.
1515 Wt. 9-10-11 rib.
                                        203.
                                              % Moisture of L.D.
152. Wt. lean 9-10-11 rib.
                                        204. Steer number.
```

		•
	•	
•		•
·	•	•
•		
•		•
•		•
•		
•		•
•	•	•
		•
•	•	
		•
-	••	•
		,
	•	
		•
•	•	•
	• •	· ·
٠	•	•
	•	•
•		•
	•	
		•
		•
	· · · · ·	
		•
	•	
•	•	•
		•
		•
•		•
•		•
•		•
		•
	· · · · · · · · · · · · · · · · · · ·	•
		•
	•	
		•
_		_
•	•	•
·	•	•

## Transformed Variables

Variables below are expressed as 205. % K.K. of total carcass. a percent of the cuts themselves: 254. % Yield of round -R.S. 206. % Untrimmed lean cuts - R.S. 255. % Fat of round - R.S. 207. % Belly cuts - R.S. 256. % Bone of round - R.S. 208. % Trimmed lean cuts - R.S. 209. % Ext. fat trim - R.S. 257. % Yield of loin - R.S. 210. % Untrand.round of R.S. 258. % Fat of loin - R.S. 259. 211. % Trimmed round of R.S. % Bone of loin - R.S. 260. % Yield of rib - R.S. 212. % Total rnd. fat trime R.S. 213. % Untrimmed loin R.S. 261. % Fat of rib - R.S. 214. % Trimmed loin R.S. 262. % Bone of rib - R.S. % Fat trim from loin. R.S. 263. % Yield of chuck - R.S. 215. 264. 216. % Untrimmed rib- R.S. % Fat of chuck - R.S. 265. 217. % Trimmed rib - R.S. % Bone of chuck - R.S. 218. % Rib fat trim - R.S. 266. % Yield of shank - R.S. % Fat of shank - R.S. **319.** % Untrimmed chuck - R.S. 267. % Trimmed chuck - R.S. 268. % Bone of shank - R.S. 220. % Chuck fat trim - R.S. 269. 221. % Yield of brisket - R.S. 222. % Shank - R.S. 270. % Fat of brisket - R.S. % Brisket - R.S. 223. 271. % Bone of brisket - R.S. 22L. % Plate - R.S. 272. % Yield of plate - R.S. % Fat of plate - R.S. 225. % Flank - R.S. 273. % RLRC - R.S. % Bone of plate - R.S. 226. 274. % Retail Yield - R.S. 227. 275. % Yield of flank - R.S. 228. 276. % Internal Fat trim - R.S. % Fat of flank - R.S. 229. % Bone - R.S. 277. % Bone of flank - R.S. 230. % Yield of round - of R.S. 231. % Int. fat trim round - of R.S. Following variables are expressed as a percent of the left sides 232. % Bone of round - of R.S. 278. % Rumpless round of L. S. % Yield of loin - of R.S. **233.** 234. % Int. fat trim of loin - R.S. 279. % Rump of L.S. % Bone of loin - of R.S. 235. 280. % Intact round of L.S. % Yield of rib - of R.S. 281. % Loin of L.S. 236. % Int. fat trim rib - of R.S. 237. 282. % Rib of L.S. % Bone of rib - of R.S. % Chuck of L.S. 238. 283. 284. 239. % Yield of chuck - of R.S. % Shank of L.S. 240. % Int. fat trim chuck - R.S. 285. % Brisket of L.S. % Bone of chuck - of R.S. 286. 2归。 % Plate of L.S. % Yield of shank - of R.S. % Flank of L.S. 242. 287. 243. % fat trim of shank- of R.S. 288. % K.K. of L.S. 244. % Bone of shank - of R.S. 289. % K.K. of R.S. % Yield of brisket - of R.S. 290. % Lean of rumpless round of L.S. 245. % Fat trim brisket - of R.S. 291. % Fat of rumpless round of L.S. 246. % Bone of brisket - of R.S. 247. 292. % Bone of rumpless round of L.S. 248. % Yield of plate - of R.S. 293. % Lean of rump of L.S. 249. % Fat trim plate - of R.S. 294. % Fat of rump of L.S. 250. % Bone of plate - of R.S. 295. % Bone of rump of L.S. 251. % Yield of flank - of R.S. 296. % Lean of intact round of L.S. 252. % Fat trim of flank - of R.S. 297. % Fat of intact round of L.S. 298. % Bone of intact round of L.S. 253. % Bone of flank - of R.S.

% Lean of loin of L.S. 299. % Fat of loin of L.S. 300. 301. % Bone of loin of L.S. 302. % Lean of rib of L.S. % Fat of rib of L.S. 303. 304. % Bone of rib of L.S. 305. % Lean of chuck of L.S. 306. % Fat of chuck of L.S. 307. % Bone of chuck of L.S. % Lean of shank of L.S. 308. % Fat of shank of L.S. 309. 310. % Bone of shank of L.S. % Lean of brisket of L.S. 311. 312. % Fat of brisket of L.S. 313. % Bone of brisket of L.S. 314. % Lean of plate of L.S. 315. % Fat of plate of L.S. 316. % Bone of plate of L.S. 317. % Lean of flank of L.S. 318. % Fat of flank of L.S. 319. % Bone of flank of L.S. 320. % Fat of K.K. of L.S. 321. % Kidney of L.S. % Total lean of L.S. 322. 323. % Total fat of L.S. 324. % Total bone of L.S. 325. % Cutting loss of L.S.

## Following variables are expressed as a percent of themselves:

326. % Lean of round - L.S. 327. % Fat of rumpless round - L.S. 328. % Bone of rumpless round - L.S. 329. % Lean of rump - L.S. 330. % Fat of rump - L.S. 331. % Bone of rump - L.S. 332. % Lean of intact round - L.S. 333. % Fat of intact round - L.S. 334. % Bone of intact round - L.S. 335. % Lean of loin - L.S. 336. % Fat of loin - L.S. 337. % Bone of loin - L.S. 338. % Lean of rib - L.S. 339• % Fat of rib - L.S. 340. % Bone of rib - L.S. 341. % Lean of chuck - L.S. % Fat of chuck - L.S. 342. % Bone of chuck - L.S. 343. % Lean of shank - L.S. 3LLL. 345. % Fat of shank - L.S. % Bone of shank - L.S. 346. 347. % Lean of brisket - L.S. 348. % Fat of brisket - L.S. 349. % Bone of brisket - L.S. 350. % Lean of plate - L.S.

351. % Fat of plate - L.S. 352. % Bone of plate - L.S. % Lean of flank - L.S. 353. % Fat of flank - L.S. 354. 355. % Bone of flank - L.S. 356. % Lean in 9-10-11 rib- L.S. % Fat in 9-10-11 rib - L.S. 357• 358. % Bone in 9-10-11 rib - L.S. 359• % Lean in 6-7-8 rib - L.S. 360. % Fat in 6-7-8 rib - L.S. 361. % Bone in 6-7-8 rib - L.S. 362. % Lean in 12th rib - L.S. 363. % Fat in 12th rib - L.S. 364. % Bone in 12th rib - L.S. 365. % 9-10-11 rib of intact rib- L.S. 366. % 6-7-8 rib of intact rib - L.S. \$ 12th rib of intact rib - L.S. 367. 368. Volume of round.

## Following variables were added to card # O after original punching:

369. Lb. total fat trim of R.S.
370. % total fat trim of R.S.
371. Est K.K. Wt. (only L.S.).
372. PPM K<sup>+</sup> (fresh basis)
373. PPM K<sup>+</sup> (protein basis)
374. PPM Na<sup>+</sup> (fresh basis)
375. PPM Na<sup>+</sup> (protein basis)

