
THE ETIOLOGY AND PATHOLOGY OF CALF PNEUMONIA

Thesis for the Degree of M. S William T. S. Thorp 1937 THESIS

Calveo-Ducases Serves



THE ETIOLOGY AND PATHOLOGY

OF

CALF PNEUMONIA

THE ETIOLOGY AND PATHOLOGY

OF

CALF PNEUMONIA

THESIS

Submitted to the Faculty of Michigan State College
in partial fulfillment of the requirements for the degree of Master
of Science in Animal
Pathology

By
William T. S. Thorp
June 1937

THESIS

. . .

TABLE OF CONTENTS

Introduction	-	_]
Review of Literature	-	- 2
Etiological and Pathological Study	-	- 3
Sources of materials and data Method of study in gross Method of microscopic study Case reports		
Discussion	-	- 52
Summary	-	- 55
Tables and Charts	-	- 59
Photomicrographs	-	- 74
Bibliography	-	- 119
Acknowledgments	_	- 120

INTRODUCTION

One of the diseases which has a most demoralizing effect on the dairyman is calf pneumonia, since his future production depends upon the well-being of the young stock. In the two years that pneumonia occurred in one large dairy herd, the following was noted:

The first year, thirty-six calves out of fifty developed pneumonia, thirteen of these resulting fatally. The second year, seventeen calves out of thirty-one developed pneumonia with eight terminating fatally (7). With a mortality of approximately twenty-five per cent over the two-year period, the effect on the dairyman can be easily understood.

It has been well established that any abnormal changes in the structure of an organ are usually followed by disturbances in the function of that organ. There has been considerable study of calf pneumonia from an etiological point of view and in associating it with other diseases. Yet a study in which the structural or pathological changes are the paramount thought seems to be lacking. With this thought in mind, but not ignoring the etiology of the disease, the following study was undertaken.

REVIEW OF LITERATURE

There is a small amount of literature dealing with calf pneumonia as a sequel to calf scours. A pneumonia of calves associated with white scours has been described by Nocard (1) in 1901 and 1902 although his description of this pneumonia does not compare with that of American workers; namely, Hagen (2) and Carpenter (3). In 1917, Hagen (2) described a pneumonia associated with white scours in calves at which time he stated that in some cases scouring is negligible while joint lesions and a peculiar type of bronchopneumonia may occur. He also reports the isolation of the colon bacillus from lungs of many of the cases of calf scours. Probably the most important literature (in English) in which calf pneumonia has been described as a separate disease is the work of Carpenter and Gilman (3) in 1921. They state that the lesions in the lungs vary considerably as to the type of pneumonia and the duration of the disease. Carpenter states in this work that he has isolated from the calves' lungs micrococci, B. Pyogenes, colon bacilli and streptococci, the streptococci predominating. Smith reports a pneumonia in fetuses and calves in which Brucella abortus is the causative organism. He suggested at this time that Brucella abortus may be a predisposing factor when pneumonia is caused by other organisms (4). In an earlier report, Smith mentions the isolation of a pleomorphic bacillus from a case of calf pneumonia (5). Hallman, Sholl, and Delez describe the lesions of fetal pneumonia associated with Brucella infections (6).

ETIOLOGICAL AND PATHOLOGICAL STUDY

Sources of Material and Data

The materials used in this work were obtained during the years 1935 and 1936. They were taken from cases of calf pneumonia brought to the laboratory for autopsy during the two-year period.

Method of Study in Gross

In a larger number of the cases, photographs were taken of the gross specimens to make a more complete record of the extent of the gross lesions. As far as possible, descriptions were made separately of the left and right lungs. Blocks of tissue were taken from various parts of the different lobes (11) depending on the extent of the pneumonia and gross variations in consolidated areas. The following key was used in identifying the different areas from which tissue was taken:

- #1 Diaphragmatic lobe.
- #2 Posterior portion of the cardiac lobe.
- #3 Anterior portion of the cardiac lobe when in two parts.
- #4 Apical lobe.

These numbers were prefixed by the letter L or R to designate the left or right lung. The letter M was used to designate the mediastinal lobe. As accurately as possible, each lobe was equally divided into four planes; namely, A, B, C, and D, A being the plane nearest the hilus of any particular lobe. The four planes mentioned above were cut

at right angles to the main or central bronchial tube of the lobe from which the specimen was taken. For example, 10 R4C would be a section from Case #10, the right apical lobe, in a plane about half way from the hilus to the apex of the lobe. The mediastinal lymph nodes were designated by MLG and the bronchial by BLG. Brief gross descriptions were made at the time each block of tissue was obtained.

Method of Microscopic Study

Blocks of tissue were taken in such a way that the sections would be in the same plane as those previously described as A, B, C, and D. The number of blocks per case depended upon the extent of the pneumonia and the gross variations which could be recognized, the number of areas represented per case ranging from 6 to 27 in number and usually averaging from 12 to 15. Blocks were taken from the mediastinal and bronchial lymph nodes. All sections of tissue taken for microscopic examination were placed immediately in Zenker's solution and embedded in paraffin for sectioning. Sections from all cases were stained by the eosin hematoxylin method. Sections from some cases were also stained for bacteria.

Bacteriological Study

The lungs were cultured only in those cases which had been dead less than six hours. When postmortem changes were evident, the material was not studied either bacteriologically or histologically. Blocks of tissue for the inoculation of culture plates were taken from areas that presented decided variations upon gross examination. These

blocks were taken before those for use in the microscopic study. If the pathology was uniform throughout the pneumonic portion of the lung, a representative number of blocks were taken, usually one or two from each lobe. The inoculated plates were identified according to the location from which the blocks of tissue were taken. The same key was employed here as that used in identifying blocks of tissue for microscopic study. The contiguous lymph glands were cultured in all cases but the heart blood only in those which had died recently or had been killed.

Sterile precautions were taken as follows:

- (1) The surface of the lung from which a block was to be removed was seared. This was done by using a piece of flat metal, 4 cm. in diameter, which was heated until red-hot.
- (2) Using sterile forceps and scissors, a small block of tissue approximately 1 x 2 cm. was removed and three culture plates were inoculated with each block.

The media used was a dextrose blood agar. The addition of .5% dextrose tended to give a more abundant growth at the end of the twenty-four hour period. Transfers were made to blood and plain agar slants. These were used as stock cultures for the bacteriological determinations.

The bacteriological determinations shown in Tables I and II indicate, in general, the procedure followed with each culture. Many similarities based on Bergy's classification will be noted in a comparison of Tables I and II. In the following case reports, the bacteriology of each case will be briefly given.

CASE REPORTS

Case #1 (Autopsy 3028), a female Holstein, eight months old, which had shown a slight cough for four days before slaughtering in the course of an experimental project.

Gross examination:

The only gross lesions noticeable are widely scattered congested areas in the apical and cardiac lobes of both lungs. Upon section of the lobes, these are distinctly lobular in their outline. They become more numerous toward the hilus of the lobes, forming massive areas of congestion in several lobes. The mediastinal lobe shows only one congested lobule at its hilus. The mediastinal and bronchial lymph nodes are normal.

Microscopic examination:

There is a slight increase in the number of lymphocytes surrounding the small bronchi and bronchioles. Sections from the grossly involved areas show a marked congestion of the alveolar walls with considerable atelectasis. No abundant fluid or cellular exudate is present in the alveolar sacs. Considerable normal tissue is seen in this case (Figs. 17 and 18). The lymph nodes are normal.

Case #2 (Autopsy 3036), a male Holstein, ten weeks old which had a pneumonia of one week's duration before death.

Gross examination:

There is red hepatization of the cardiac and apical

lobes of both lungs. The cardiac lobes seem slightly
firmer than the apical lobes. A few grayish-yellow areas
1 to 2 mm. in diameter are seen on the pleural surface of
the cardiac and apical lobes. These areas are largest
toward the hilus of the lobes and upon section are more
numerous on the cut surface.

The diaphragmatic lobes of both lungs show pneumonic areas in their anterior one-fourth which, upon section, appear to radiate from the hilus of the lobe. The posterior borders of the involved portion of the diaphragmatic lobes appear lobular in nature. The mediastinal and bronchial lymph nodes are enlarged and upon section a thin yellowish fluid exudes from the cut surface.

Microscopic examination:

In both lungs there is an acute bronchitis with much desquamation of the epithelium. All the bronchi and larger bronchioles which show any congestion of the mucosa contain a large number of desquamated epithelial cells in their lumina. In a number of the terminal and respiratory bronchioles there is evidence of productive tissue changes in the bronchial wall.

The majority of the alveoli are filled with a recent inflammatory exudate consisting of a coarse network of fibrin, mononuclear cells and a few polymorphonuclear cells.

The inflammatory process is of longer duration in the cardiac and apical lobes than in the diaphragmatic lobes.

The apical lobe shows more evidence of a prolonged pneumonia

than the cardiac lobe. The mediastinal and bronchial lymph nodes contain large numbers of polymorphs and mononuclear phagocytes in their sinuses.

_ _ _ _ _

Case #3 (Autopsy 3028), a female Holstein, three months old, which had shown clinical symptoms of pneumonia for ten days prior to death.

Gross examination:

The cardiac and apical lobes are in a state of grayishred consolidation with very prominent interlobular septa,
probably due to edema. In the left apical and cardiac lobes,
there are more grayish areas than in the corresponding lobes
of the right lung. The anterior one-third of the right
diaphragmatic lung is in a state of red hepatization, only
a few small grayish-yellow areas are seen on the pleural
surface at the hilus of the lobe. The anterior one-fifth
of the left diaphragmatic lobe appears the same as the
cardiac and apical lobes of that lung.

The mediastinal and bronchial lymph nodes are very much enlarged with the mediastinal nodes showing numerous petechial hemorrhages on the surface.

Microscopic examination:

The pleura is practically unaltered. Considerable atelectasis has taken place. In the areas of very recent involvement, there are a large number of mononuclear phagocytes in the stroma of the bronchi and many lymphocytes in the peribronchial tissue. Proportionally, there is more mononuclear infiltration in the mucosa of the terminal

bronchioles than in the mucosa of the bronchi. In the apical lobe, and to some extent in the cardiac lobe, there is considerable evidence of productive tissue changes in the walls of the terminal and respiratory bronchioles. The alveolar walls show an increase in the number of fibroblasts and macrophages. The exudate in the alveoli consists of mononuclear phagocytes (Fig. 63) and polymorphonuclear leucocytes. The above changes are more noticeable adjacent to the bronchioles. The alveoli at a greater distance from the bronchioles do not show any evidence of thickening of the alveolar walls. In many areas, the alveoli appear as if lined with mononuclear phagocytes. Lymph nodes show only a mild congestion.

case #4 (Autopsy 3085), a calf which was one of a set of triplets and the first one to become sick. Just prior to the pneumonia, the calves were changed to another nurse cow and developed scours. This calf became sick on March 14, 1935, and died on April 5, 1935, at the age of one month. When the animal showed the first signs of pneumonia, (Temp. 104.8) it was given 40 cc. of calf scour antiserum. Three days of improvement followed.

On March 25, 1935, the calf was noticeably worse, (Temp. 102). Loss of weight and persistent coughing were accompanied by accelerated respiration and diarrhea. Daily injections of calf scour antiserum were made until April 5, 1935.

Gross examination:

The cardiac and apical lobes of both lungs show a grayish-red consolidation with a pleura studded by numerous reddish-brown areas ranging in size from 1 to 2 mm, in diameter which suggests necrosis. The diaphragmatic lobes are consolidated only in the anterior one-third which appear much the same as the adjacent cardiac lobes. The mediastinal lobe is similar in appearance to the cardiac Throughout the cardiac and apical lobes of both lobes. lungs, many small abscesses, 5 mm. to 1 cm. in diameter, appear just beneath the pleural surface. The abscesses seem more numerous toward the hilus of the lobe. involved portions of the left lung show more gray areas indicative of an abundant cellular exudate. Upon section of any of the lobes, a foamy purulent exudate exudes from the smaller bronchi and bronchioles which is considerably thicker than that from the larger bronchi. Many abscesses similar to those noted on the pleural surface can be seen upon section. If a lobe of the lung showing numerous abscesses on the pleural surface is cut so that the main bronchi and some of their branches are divided lengthwise, several of the abscesses will appear at the termination of the small bronchi and bronchioles. On the cut surface of the anterior lobes, many opaque yellowish patches are observed which include an entire lobule. The mediastinal and bronchial lymph nodes are enlarged and have a distinct zone of hemorrhage around the peripheral border.

Microscopic examination:

The pleura is slightly thickened. The interlobular septa are wide and edematous and contain lymphatics distended with mononuclear phagocytes. In many of the small bronchi, the mucosa is definitely thickened by a marked congestion accompanied by considerable desquamation of the epithelium. Productive tissue changes are very much in evidence around the small bronchioles. This is accompanied by considerable mononuclear infiltration. No productive tissue changes are noted in the walls of the alveoli surround ing the bronchioles. There are many necrotic areas .5 to 1 cm. in diameter (Fig. 55). The alveolar walls are very congested, the alveoli being filled with an exudate consisting of mononuclear phagocytes and a small number of polymorphs in a network of fibrin. In many instances, there is such a mononuclear exudation that the alveolar wall is not discernible. In some areas, the alveolar walls and the cells of the exudate are necrotic. The lymph nodes contain a large number of polymorphs in their sinuses.

Case #5 (Autopsy 3110), a male Guernsey, one month old which had a pneumonia of two weeks' duration before death. On the first day of sickness the temperature was 104°. The range of temperature during the course of the pneumonia was 104° - 106.4°. There were shallow and accelerated respirations, also a masal discharge and cough throughout the course of the pneumonia.

Gross examination:

Upon autopsy there is a pneumonia very similar to the previous cases. There is a consolidation of the cardiac and apical lobes of both lungs, characterized by a grayish-red consolidation with many areas of an acute nature. The anterior one-third of the diaphragmatic lobes are involved in an acute pneumonia. The lymph nodes are enlarged, edematous and have many petechial hemorrhages.

Microscopic examination:

The pleura is unaltered. The interlobular septa are edematous with a slight distention of the lymphatics. There is an acute bronchiolitis with some desquamation of the epithelium. In some areas, there is considerable cellular exudate in the lumina of the small bronchi and larger bronchioles. There are many areas of a recent bronchiolitis (Figs. 19 and 20).

A large number of the alveoli are filled with a cellular exudate consisting of mononuclear phagocytes and a few polymorphs. In the sections from the more recent areas of pneumonia, the alveoli and alveolar ducts contain a fibrinous exudate (Fig. 22). In these areas, the alveolar walls are very much congested. The lymph nodes show only a slight congestion and a few red blood cells in the sinuses.

Case #6 (Autopsy 3112), a female Holstein seven months old. There were clinical indications of pneumonia for three days prior to death.

Gross examination:

There is a very marked congestion of the cardiac and apical lobes of both lungs, which extends into the anterior two-thirds of the diaphragmatic lobes. The pneumonic areas are not as firm in consistency as they appear. The lobules are clearly outlined by their edematous interlobular septa. The mediastinal lobe is uniformly congested and similar in appearance and consistency to the other pneumonic lobes. Upon section, a thin bloody fluid exudes from the cut surface. Some of the large bronchi have a very congested mucosa. The mediastinal and bronchial lymph nodes are very edematous and hemorrhagic.

Microscopic examination:

The pleura is not thickened. The bronchi show a very acute bronchitis with much injury to their mucosae. There is considerable desquamation of the epithelium (Fig. 37). The lumina are filled with a serous exudate containing many desquamated epithelial cells (Fig. 38). The majority of the terminal and respiratory bronchicles are filled with a serous exudate, yet the epithelium does not show any marked injury. There is very much congestion of the alveolar walls (Fig. 21). About five per cent of the alveoli contain air and are very much distended. Most of the remaining alveoli are partially collapsed and filled with a serous exudate. A small number of the alveoli are filled with a fibrinous exudate (Figs. 25 and 26), while still others contain many blood cells. The lymph nodes are very

hemorrhagic, the lymph sinuses being filled with blood (Fig. 71).

Bacteriological examination:

Bacteriological examination reveals a staphylococcus, probably staphylococcus aureus. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #7 (Autopsy 3115), a male Holstein, four weeks old which had shown a pneumonia for ten days prior to death.

Gross examination:

There is a grayish-red consolidation of the apical, cardiac, and the anterior one-third of the diaphragmatic lobes of both lungs. The greater portion of the pneumonic areas are in a state of atelectasis with the exception of an emphysematous area in the right diaphragmatic lobe. This area is about 3 cm. in diameter. The pleural surfaces of the pneumonic lobes are studded with grayish-yellow areas 1 to 2 mm. in diameter. These areas are more numerous in the cardiac and apical lobes. Upon section, a purulent exudate exudes from the bronchi and bronchioles. The mediastinal and bronchial lymph nodes are enlarged and upon section, a thin yellowish fluid exudes from the cut surface.

Microscopic examination:

There is no thickening of the pleura or edema of the interlobular septa. Some of the lymphatics of the pleura

are distended with mononuclear phagocytes and a few lymphocytes. The lymph nodes show no marked changes. The mucosae of the bronchi are infiltrated with large numbers of macrophages. There is no desquamation of the epithelium in the larger bronchi. The terminal and especially the respiratory bronchioles show considerable productive tissue changes in their walls. The lumina contain a large number of mononuclear phagocytes and a few polymorphs. In those lobes where the pneumonia is of longer standing, the epithelium of the small bronchioles is desquamated. The peribronchial tissues show a large number of proleferating fibroblasts and mononuclear cells. In the anterior lobes, the alveolar walls are thickened by an increased number of fibroblasts. In approximately twenty-five per cent of the alveoli, the exudate consists of polymorphs and a few macrophages. In the remainder of the alveoli, the exudate is mostly mononuclear.

Bacteriological examination:

A gram negative short rod of the colon typhoid group, probably Bacillus coli communior, and a hemolytic streptococcus of the beta type resembling streptococcus pyogenes were isolated from this case. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #9 (Autopsy 3135), a female Holstein four months old, which had shown a pneumonia for twelve days prior to death.

Gross examination:

There is an early pneumonia involving the cardiac and

apical lobes of both the lungs and the anterior one-third of the right diaphragmatic, also two-thirds of the left diaphragmatic. In contrast with most of the cases, the pneumonia is more acute in the left lung than it is in the right lung. The anterior lobes of the right lung are studded by grayish-white areas indicating an abundant cellular exudate. A smaller number of similar areas are noted in the apical lobe of the left lung. The mediastinal lobe is entirely consolidated with grayish areas toward the hilus of the lobe. Upon section, the grayish areas become more numerous and a grayish-yellow exudate exudes from the bronchial tubes. The lymph nodes are enlarged and covered with petechial hemorrhages. Upon section, a clear watery fluid exudes from the cut surface.

Microscopic examination:

The pleura is thickened slightly over the pneumonic areas. In the diaphragmatic lobes which are more recently involved, there are only mild changes in the bronchi and bronchicles. Some show a slight mononuclear infiltration of the stroma, while a few more contain a small amount of serous exudate. There are some recent productive tissue changes in some of the small bronchicles (Figs. 49 and 50). There is much congestion of the alveolar walls accompanied by a moderate mononuclear exudate in the alveoli. The exudate is more abundant in the alveoli surrounding the respiratory bronchicles. In the cardiac lobes, where the pneumonia is of longer standing, the bronchi are hyperemic

with slight desquamation of the epithelium. There is decidedly more desquamation of the epithelium in the terminal bronchioles than in the bronchi or the respiratory bronchioles. The alveolar exudate varies from a serous exudate to one containing many mononuclear phagocytes. The apical lobes show a pneumonia of considerably longer standing than any of the other lobes. There is evidence of productive tissue changes around the bronchioles, especially the respiratory bronchioles. The exudate in the alveoli of the apical lobe has a tendency to be more cellular than in the other lobes. No marked changes were noted in the lymph nodes.

Bacteriological examination:

Hemolytic streptococci which resemble very closely streptococcus pyogenes were isolated from this case. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #10 (Autopsy 3146), a male Holstein, two and one-half months old, which had a pneumonia of twelve days' duration before death.

Gross examination:

The cardiac and apical lobes and the greater part of the pneumonic areas in the diaphragmatic lobes are in a state of grayish consolidation. The right diaphragmatic lobe shows areas in which one or two lobules are consolidated but are separated to a certain extent by normal lobules. The left diaphragmatic lobe is about one-fourth

pneumonic. There are many small circumscribed grayishyellow areas noted upon section of all the pneumonic lobes
suggestive of considerable cellular exudate. A large
amount of frothy, greenish-yellow purulent exudate exudes
from the small bronchi and bronchioles upon section of the
cardiac and apical lobes of either lung. The mediastinal
lobe is entirely consolidated and has a large number of
grayish-white areas on the pleural surface. The mediastinal and bronchial lymph nodes are swollen and hemorrhagic.

Microscopic examination:

The pleura is thickened. The interlobular septa are slightly edematous with some distention of the lymphatics. In the lobes most recently affected, as well as in the other lobes, there is a very marked at electasis. In those areas more recently involved, the bronchi show recent mononuclear infiltration of their mucosa, the alveoli are in a state of atelectasis and are very congested. Lobules that show a pneumonia of longer duration show a marked mononuclear infiltration of the mucosa of the bronchi with much desquamation of the epithelium. The terminal bronchioles reveal essentially the same changes. In some instances, the respiratory bronchioles appear entirely destroyed. In a small number of the bronchioles, there is evidence of productive tissue changes in the wall. There are some respiratory bronchioles which have a mononuclear infiltration of the peribronchial tissue (Fig. 47). The alveoli in this portion of the lung are

very congested and contain a large number of mononuclear phagocytes and a few polymorphs. In a few scattered areas, a serous exudate occurs (Figs. 23 and 24). The mediastinal and bronchial lymph nodes are also very congested.

Case #11 (Autopsy 5147), a male Holstein five months old which had a pneumonia of two weeks' duration before death.

Gross examination:

There is a marked atelectasis of all of the pneumonic lobes. It is especially noticeable in the areas of longer duration. The pneumonia involves all of the anterior lobes and approximately one-sixth of the diaphragmatic lobes. In the diaphragmatic lobes, recent red hepatization is noted. In the apical lobes, and to a lesser extent in the cardiac lobes, many grayish-yellow areas are seen on the pleural surface and upon section. The anterior lobes are considerably firmer in consistency than the pneumonic portions of the diaphragmatic lobes. The pneumonia is of longer duration in the right apical lobe than in the left apical. About three-fourths of the mediastinal lobe is consolidated. The mediastinal and bronchial lymph nodes are very enlarged and are a pale grayish-yellow color. Upon section, a yellowish turbid fluid exudes from the cut surface.

Microscopic examination:

There is considerable thickening of the pleura and in a few scattered areas, there is evidence of a fibrinous

pleurisy. The interlobular septa are edematous and the lymphatics are somewhat enlarged. The diaphragmatic lobes of both lungs present recent pneumonia with a marked atelectasis of the involved portions of the lobes. There is some desquamation of the epithelium in the bronchi and bronchioles of these lobes. In the cardiac lobes, there is evidence of chronicity, especially in the terminal and respiratory bronchioles, the walls of which show marked productive tissue changes. Polymorphonuclear leucocytes predominate in the alveolar exudate. In the apical lobes, there is some organization of the exudate in the small bronchioles and some productive tissue changes in the alveolar wall. These changes are accompanied by considerable atelectasis. The organized exudate in the bronchioles appears covered by cuboidal cells except at the point of attachment to the bronchial wall. For the most part, the alveoli are filled with a cellular exudate. The lymph sinuses of the mediastinal and bronchial lymph nodes are filled with polymorphonuclear leucocytes and some mononuclear phagocytes.

Bacteriological examination:

A hemolytic streptococcus of the beta type and a gram negative short rod of the colon typhoid group, probably Bacillus coli communior, were isolated from this case.

For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #12 (Autopsy 3150), a female Holstein, nine months

old which had a pneumonia of one week's duration prior to death.

Gross examination:

The cardiac and apical lobes and a small portion of the diaphragmatic lobes are involved in a very unevenly distributed bronchopneumonia. The dorsal borders of the apical and cardiac lobes are normal, pneumonia involving only the lower portions of the lobes. There are a few circumscribed grayish-yellow areas, 2 mm. in diameter, suggesting recent cellular exudate. These areas are present only in the pneumonic portion of the left apical lobe. The mediastinal lobe is entirely consolidated. The mediastinal and bronchial lymph nodes are a very light yellow with a few petechial hemorrhages on the surface. On section, a yellowish fluid exudes from the cut surface.

Microscopic examination:

The pleura is very slightly thickened. The interlobular septa do not show any marked changes. The microscopic changes in this case correspond to a certain extent with those of Case #6. There is a very acute bronchitis with much injury to the bronchial mucosa manifested by much desquamation of the epithelium, also very much congestion of the alveolar walls accompanied by marked serous exudation of the alveolar sac. In those lobules in which there is a slight congestion of the alveolar wall, there is some atelectasis of the alveoli. In the cardiac and apical lobes, there is a marked atelectasis and congestion of the alveolar walls accompanied by a slight cellular exudation into the alveoli. Several of the bronchi show some mononuclear infiltration in their mucosa. The lymph sinuses of the lymph nodes contain many polymorphonuclear leucocytes.

Bacteriological examination:

Staphylococcus aureus and a hemolytic streptococcus of the beta type were isolated from this case (Fig. 72).

For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #13 (Autopsy 3154), a male Holstein three months old, which had pneumonia of one week's duration prior to death.

Gross examination:

Neither the pleura nor the interlobular septa show any gross changes. The pneumonia involves the mediastinal, cardiac, and apical lobes, and approximately one-fifth of each diaphragmatic lobe. Immediately posterior to the areas of acute pneumonia of the anterior part of the diaphragmatic lobe, there are several large areas of emphysema. The cardiac and apical lobes are in a state of red hepatization and show no evidence of cellular exudation. The mediastinal lobe is much the same as the right cardiac. The left lung is in a more acute stage of pneumonia than the right, which is somewhat more patchy and not as evenly distributed as that of the right lung. The lymph nodes are swollen, edematous, and show evidence of an acute lymphadenitis.

Microscopic examination:

The pleura is unaltered as is the interlobular septa. The only acute bronchitis with severe injury to the bronchial mucosa is in the apical lobes. The diaphragmatic lobes show only an acute congestion of the alveolar wall with partial collapse (Fig. 30) and some with a small amount of mononuclear exudation into the air sacs. In the cardiac lobe, there is some serous exudate but more cellular exudate. There is comparatively more cellular exudate of the mononuclear type in the air sacs than in the cardiac and diaphragmatic lobes. The lymph nodes show hemorrhages into their lymph sinuses.

Bacteriological examination:

A staphylococcus very much like staphylococcus aureus was isolated from this case. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #14 (Autopsy 3166), a male Guernsey three months old which had pneumonia one week prior to death.

Gross examination:

There is a very acute pneumonia of the cardiac and apical lobes of both lungs. The pneumonia of the left diaphragmatic lobe is more extensive than that of the right diaphragmatic, only a small portion in the anterior part of the right lobe being affected. There are areas of very marked congestion and red hepatization which include from

	-	
-		
	•	
	· ·	
•		

three to four lobules. These are scattered throughout the consolidated portions of the various lobes. The mediastinal lobe is entirely consolidated. There are three consolidated areas 2 cm. by 4 to 5 cm. in the left diaphragmatic lobe showing a very marked congestion of, or hemorrhage into, the alveoli (Fig. 1). There are no gross changes in the mediastinal and bronchial lymph nodes.

Microscopic examination:

There is a marked distention of the lymphatics of the pleura. Those of the interlobular septa are distended only to a slight extent. There is also a moderate infiltration of mononuclear phagocytes into the mucosa of the bronchi accompanied by some congestion. The above pathology of the bronchi repeats itself in the terminal bronchioles, but with a more severe bronchiolitis which results in some desquamation of the epithelium. A number of the bronchioles contain a small amount of a cellular exudate consisting mostly of mononuclear phagocytes. In the anterior lobes, especially the right apical, there is a marked congestion of the alveolar walls accompanied by considerable cellular exudation into the alveolar sacs. There is no appreciable variation between the cardiac and diaphragmatic lobes. these lobes, there is only a marked congestion of the alveolar walls accompanied by a serous exudate in a small number of the alveoli. The lymph nodes contain a large number of polymorphonuclear leucocytes in their lymph sinuses.

• . • .

•

Bacteriological examination:

A hemolytic streptococcus of the beta type resembling streptococcus pyogenes and a gram negative short rod, probably Bacillus coli communior, of the colon typhoid group were isolated from this case. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #15 (Autopsy 3172), a male Holstein about ten weeks old, which had a pneumonia of eleven days' duration prior to death.

Gross examination:

The pneumonia involves the apical, cardiac, mediastinal, and diaphragmatic lobes of both lungs. The right cardiac and apical lobes are in a state of reddish consolidation which are studded by small gray areas 1 to 2 mm. in diameter. The anterior one-fifth of the right diaphragmatic lobe is congested. The interlobular septa appear thickened and perhaps edematous. Upon section of the apical or cardiac lobe, the grayish areas noted on the pleural surface appear more numerous. The cardiac and apical lobes of the left lung have more yellowish-gray areas on their pleural surfaces than the same lobes of the right lung. The anterior one-third of the left diaphragmatic lobe appears very much the same as the left apical and cardiac lobes. The mediastinal lobe is about two-thirds consolidated. The mediastinal and bronchial lymph nodes are very much enlarged and edematous.

Microscopic examination:

The pleura is thickened. The interlobular septa are edematous and contain lymphatics distended with mononuclear phagocytes. The apical lobes of both lungs show a pneumonia of considerably longer standing than that of the remaining lobes. There is an acute bronchitis in all the pneumonic lobes accompanied by much desquamation of the bronchial epithelium. The bronchitis is not as severe in the areas of shorter duration. The terminal and respiratory bronchicles show approximately the same pathology as the bronchi. Many of the respiratory bronchicles are filled with a cellular exudate which shows considerable necrosis in many of the bronchicles.

There is much congestion of the alveolar walls throughout all the sections. The alveoli in the anterior lobes are
filled with a cellular exudate consisting mostly of mononuclear phagocytes and a small number of polymorphonuclear
leucocytes. In the right diaphragmatic lobe where the
pneumonia is more acute, a number of alveoli are filled with
a fibrinous exudate. There are a number of widely scattered
alveoli in the cardiac lobe which are filled with a fibrinous
exudate which in some cases contains many polymorphonuclear
cells. The lymph nodes show marked changes.

Bacteriological examination:

Staphylococcus aureus and a non-hemolytic streptococcus were isolated from this case. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #16 (Autopsy 3200), a female Jersey ten weeks old which had a pneumonia of eighteen days' duration before death. The first symptoms noted were a dry cough and a nasal discharge. The daily temperatures are given in Chart III.

Gross examination:

The pneumonia involves the cardiac and apical lobes of both lungs, the entire diaphragmatic lobe of the left lung, and approximately four-fifths of the right diaphragmatic (Fig. 2). The pleural surfaces of the cardiac and apical lobes of both lungs and a greater part of the left diaphragmatic lobe are studded with grayish-yellow areas 1 to 3 mm. in diameter. In the right lung, these areas are present only in the anterior portion of the right diaphragmatic, the remaining part of the lobe appearing more acute. The mediastinal lobe is entirely consolidated. The cardiac and apical lobes appear more atelectatic than either of the diaphragmatic lobes. The mediastinal and bronchial lymph nodes are enlarged and upon section, a few petechial hemorrhages are noted on the cut surface.

Microscopic examination:

The pleura covering the cardiac and apical lobes shows some thickening. The interlobular septa are unchanged. The bronchi in any of the pneumonic areas of the lung show an acute bronchitis with much desquamation of the epithelium. Many bronchi show an abundant mononuclear infiltration of the mucosae. There is some evidence of chronicity in the

cardiac and apical lobes. A very large number of the terminal and respiratory bronchioles of the pneumonic areas in either lung show considerable necrosis of the cells of the exudate and walls of the bronchioles (Fig. 48). A number of elveoli show an increase of fibroblasts in the alveolar wall. In areas in which there is considerable thickening of the alveolar wall, the alveoli appear lined with cuboidal cells. The lymph nodes show no marked changes.

Bacteriological examination:

A hemolytic streptococcus of the beta type similar to streptococcus pyogenes and a short rod, probably Bacillus coli communior, were isolated. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #17 (Autopsy 3198), a male Holstein, one month old, which had a pneumonia of fifteen days' duration before death.

Gross examination:

of both lungs and a few scattered areas in the anterior portion of the diaphragmatic lobes. The pneumonia of the mediastinal lobe appears of the same duration as the right apical lobe. There is a grayish-yellow consolidation of the cardiac and apical lobes which appears at electatic.

Upon section of the cardiac and apical lobes, the apical lobe shows more evidence of cellular exudation. The

. · • • •

pneumonia in the left lung appears of longer duration than in the right lung. The scattered areas of pneumonia in the right diaphragmatic lobe are surrounded, in some instances, by a thickened interlobular septa. The mediastinal and bronchial lymph nodes are swollen and somewhat edematous.

Microscopic examination:

The pleura is slightly thickened. The interlobular septa are somewhat edematous and contain distended lymphatics. In most of the pneumonic areas there is a mononuclear infiltration of the mucosae of the bronchi which is accompanied by a varying amount of desquamation of the epithelium. Many of the respiratory bronchioles show a marked mononuclear infiltration of the stroma of the mucosa (Figs. 43 and 44). There are many areas throughout the pneumonic parts of the lungs in which there are considerable productive tissue changes in the alveolar walls. This is especially noticeable in the apical lobes of both lungs. There are numerous areas of hemorrhage in the peripheral portions of the mediastinal and bronchial lymph nodes.

Bacteriological examination:

Staphylococcus aureus and a gram negative short rod, probably Bacillus coli communior, were isolated. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #18 (Autopsy 3203), a female Jersey, ten weeks old which had a pneumonia of three weeks' duration prior

to death. At times during the three-week period, this animal showed signs of improvement and then relapses occurred. Daily temperatures are included in Chart III.

Gross examination:

The pneumonia involves the apical and cardiac lobes of the left and right lungs. The diaphragmatic lobe of the right lung is two-thirds pneumonic while the same lobe of the left lung is only one-third pneumonic. There is a marked atelectasis of the apical lobes, the pleural surfaces of which are studded by circumscribed yellowish-gray areas 1 to 3 mm. in diameter. These areas are more numerous in the left apical than in the right. The cardiac and diaphragmatic lobes of the left lung do not show as much evidence of cellular exudation as the left apical lobe, yet they show considerably more than the corresponding lobes of the right lung. The gross appearance of the mediastinal lobe is similar to that of the left cardiac. Upon section of any of the lobes of either lung, a purulent exudate exudes from the small bronchi and bronchioles. The mediastinal and bronchial lymph nodes are swollen, whitishyellow in appearance and upon section, a yellowish turbid fluid exudes from the cut surface

Microscopic examination:

The pleura covering both lungs is thickened (Fig. 27). In places, the interlobular septa are edematous and the lymphatics are distended with mononuclear phagocytes and a small number of lymphocytes. The bronchi of the lungs show

an acute bronchitis with a very congested mucosa and much desquamation of the epithelium. For the most part, the lumina of the bronchi contain an exudate of mononuclear phagocytes and polymorphs in which there are many desquamated epithelial cells. There is a marked congestion of the mucosa of the terminal bronchioles and some desquamation of their epithelium (Figs. 41 and 42). There is some evidence of productive tissue changes in the respiratory bronchioles of the apical lobes.

There is a marked congestion of the alveolar walls in all the sections studied. Considerable atelectasis is noted in the more anterior lobes. In one-third of the alveoli, the exudate is approximately equally divided between a serous and fibrinous exudate. In the remaining two-thirds, approximately seventy-five per cent of the alveoli are filled with varying amounts of a cellular exudate consisting of mononuclear phagocytes and a few polymorphs. The lymph nodes contain many polymorphs in their sinuses.

Bacteriological examination:

Staphylococcus aureus, a gram negative short rod, probably Bacillus coli communior, and a gram positive coccus, probably a micrococcus, were isolated. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #19 (Autopsy 3209), a female Jersey three months

old which had a pneumonia of eleven days' duration before death.

Gross examination:

There is a grayish-yellow consolidation of the apical and cardiac lobes of both lungs. The anterior one-third of the left diaphragmatic lobe appears approximately the same as the apical and cardiac lobes of the same lung. The anterior three-fourths of the right diaphragmatic lobe is in a state of grayish consolidation with the exception that along the posterior border of the consolidated portion there is a zone of acute congestion varying in width from 1 to 3 cm. The mediastinal lobe is very much the same as the apical lobe of the left lung. The left lung as a whole appears of longer duration than the right lung. There are a few shreds of fibrinous exudate on the pleura. The mediastinal and bronchial lymph nodes are very enlarged and edematous.

Microscopic examination:

The pleura is slightly thickened. Lymphatics of the interlobular septa are somewhat distended. In most of the pneumonic areas, there is a bronchitis. In the more acute areas, there is a mononuclear infiltration of the mucosa with desquamation of the epithelium (Figs. 39 and 40). In the areas of pneumonia of longer duration, there is considerable desquamation of the epithelium. The terminal bronchioles show a slight desquamation of the epithelium. The respiratory bronchioles do not show any appreciable injury to their mucosa but contain a cellular exudate in

• . • . •

many areas. The alveoli in the areas of longer duration are filled with a cellular exudate consisting mostly of polymorphonuclear cells. In a few areas of shorter duration, the exudate consists of fibrin and a few mononuclear phagocytes. The alveolar walls for the most part are congested. There are a few polymorphonuclear cells in the lymph sinuses.

Bacteriological examination:

Staphylococcus aureus and a gram negative short rod, probably Bacillus coli communior, were isolated. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #20 (Autopsy 3212), a male Holstein four months old which showed first symptoms of pneumonia forty-eight hours before death.

Gross examination:

The cardiac and apical lobes of both lungs are in a state of red hepatization. Approximately one-half of the left and right diaphragmatic lobes are consolidated. The mediastinal lobe and the left cardiac lobe have a few grayish-yellow areas on their pleural surfaces. These are the only gross lesions in either lung which suggest a cellular exudation. The mediastinal and bronchial lymph nodes are very much enlarged.

Microscopic examination:

The pleura is slightly thickened and contains many distended lymphatics. The interlobular septa are very

edematous and their lymphatics are very much distended. The pneumonia is evenly distributed throughout both lungs. The mucosae of the larger bronchi are very much congested but show only a slight mononuclear infiltration with no desquamation of the epithelium. A number of terminal and respiratory bronchicles contain a small amount of cellular exudate. The alveolar walls for the most part are very much congested and the alveoli contain a serous exudate. In a small number of widely scattered areas, there is a fibrinous or cellular exudate. The mediastinal and bronchial lymph nodes do not show any marked changes.

Bacteriological examination:

A hemolytic streptococcus of the beta type was isolated. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #21 (Autopsy 3213), a female Jersey six weeks old which had a pneumonia of four weeks' duration prior to death. The first day, a temperature of 106.2 was recorded. Twenty-four hours before the animal died, a temperature of 101.8 was recorded. The first symptoms noticed were dyspnea, accelerated respiration and a cough. The following day, gritting of the teeth and marked depression were noted. Pneumonia mixed bacterin was given daily the first three weeks. During the first few days of the pneumonia, the calf showed considerable improvement, then for about two weeks, its condition remained at a standstill. Three days

prior to death, a decided relapse was noticed.

Gross examination:

There is a grayish-red consolidation of the cardiac and apical lobes of both lungs, also about one-third of the left diaphragmatic lobe and approximately one-half of the right diaphragmatic lobe. The mediastinal lobe appears very much the same as the right diaphragmatic lobe. The pneumonia of the left lung appears of longer standing than the pneumonia in the right lung. There is considerable atelectasis of the right lung and considerable atelectasis in the cardiac and apical lobes of the left lung. The right apical lobe appears slightly atelectatic. The interlobular septa seem thickened and possibly edematous. The mediastinal and bronchial lymph nodes are very much enlarged and upon section, a yellowish turbid fluid exudes from the cut surface.

Microscopic examination:

The pleura is slightly thickened in some of the pneumonic areas. The interlobular septa are very edematous and contain slightly distended lymphatics. In the diaphragmatic lobes of both lungs, there is an acute pneumonia with much injury to the bronchial walls. The mucosa of the small bronchi and large bronchioles is very much congested with much desquamation of the epithelium. The terminal and respiratory bronchioles do not show the extent of injury to their mucosa that the larger bronchial tubes do. In some of the respiratory bronchioles, there is evidence of organization of the bronchial exudate (Figs. 53 and 54).

The bronchioles in the cardiac and apical lobes of both lungs show productive tissue changes in their stroma, indicating a bronchopneumonia at an early date. There is a mononuclear infiltration of the stroma of the mucosa in many of the terminal bronchioles (Fig.46). This is somewhat more noticeable in the cardiac and apical lobes of the left lung. The walls of the alveoli immediately adjacent to the terminal and respiratory bronchioles are very much congested. The alveoli contain a small number of mononuclear phagocytes and in some areas a fibrinous exudate. The mediastinal and bronchial lymph nodes contain a large number of polymorphonuclear cells in their lymph sinuses.

Bacteriological examination:

A staphylococcus and gram negative short rod, probably Bacillus coli communior, were isolated. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #22 (Autopsy 3220), a female Guernsey seven weeks old which had a pneumonia of three weeks' duration before the animal was killed, as death of the animal seemed very probable. A pneumonia mixed bacterin was given during the first two weeks of the pneumonia. Glucose was given once following the above treatment.

Gross examination:

There is a grayish-red consolidation of the cardiac and apical lobes of both lungs. The diaphragmatic lobes

of both lungs are approximately two-thirds consolidated. There is a small area on the ventral border of the right cardiac lobe which does not show any pneumonia (Fig. 3). The pneumonic portions of the left lung show considerably more evidence of cellular exudation than those of the right lung. The cardiac and apical lobes of both lungs show considerable atelectasis. The mediastinal lobe resembles the left cardiac lobe. The mediastinal and bronchial lymph nodes are very much enlarged and slightly hemorrhagic (Fig. 3).

Microscopic examination:

The pleura is thickened. The interlobular septa do not show any marked changes. The bronchi show an infiltration of mononuclear phagocytes of the mucosa (Figs. 31 and 32). In all the consolidated lobes, but to a greater extent in the cardiac and apical lobes, there is evidence of productive tissue changes in the stroma of the mucosa of the terminal and respiratory bronchioles. The mucosa of the above bronchioles are infiltrated with mononuclear phagocytes. Only a very small number show any desquamation of the epithelium. In a few scattered areas, the alveolar walls show some slight thickening. Those alveoli surrounding the terminal and respiratory bronchioles are filled with polymorphonuclear leucocytes. Those alveoli further from the bronchioles are congested and many contain a few mononuclear phagocytes. The lymph sinuses contain a few polymorphonuclear leucocytes.

Bacteriological examination:

Organisms not identified. For their morphological,

cultural, and biochemical characteristics see Tables I and II.

Case #23 (Autopsy 3232), a female Holstein two months old which had a pneumonia for two weeks prior to death.

Gross examination:

There is a pneumonia of the diaphragmatic, mediastinal, cardiac and apical lobes of both lungs (Fig. 4). The anterior portion of the right cardiac lobe and all of the right apical appear very atelectatic. About one-half of the right diaphragmatic lobe is in a state of reddish consolidation. The anterior one-third of the left diaphragmatic lobe is in a state of grayish-yellow consolidation. All of the pneumonic areas of the left lung appear very atelectatic. Upon section of the more acute areas in the left diaphragmatic, a grayish consolidation is the most predominant (Fig. 5). The mediastinal lobe is entirely consolidated and very similar to the lobes of the left lung. The interlobular septa appear thickened and edematous. The mediastinal and bronchial lymph nodes are enlarged and edematous.

Microscopic examination:

The pleura is thickened. The interlobular septa are edematous and their lymphatics are very much distended. The apical and cardiac lobes of both lungs show a severe bronchitis with much desquamation of the epithelium (Figs.33 and 34). In the right apical lobe, there is evidence of considerable productive tissue changes in the walls of the

respiratory and terminal bronchioles. The alveoli in the cardiac and apical lobes of both lungs are filled with a cellular exudate consisting mostly of mononuclear phagocytes and a small number of polymorphonuclear leucocytes. The areas of pneumonia in the anterior portion of the right diaphragmatic lobe do not show as many productive tissue changes as the apical and cardiac lobes. In this lobe, there is considerable congestion of the alveolar wall accompanied by an abundant serous exudate in the alveolar sacs. In a small number of alveoli, there is a fibrinous appearing exudate surrounded by cuboidal-like cells (Figs. 61 and 62). Many of the alveoli contain large numbers of giant cells (Fig. 64). There is some increase in the number of polymorphs in the lymph sinuses.

Bacteriological examination:

A streptococcus of the beta type was isolated. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #24 (Autopsy 3235), a male Holstein ten days old which had a pneumonia of six days' duration prior to death.

Gross examination:

A purulent fibrinous exudate covers the pleural surface of both lungs (Fig. 6). This is especially noticeable in the anterior lobes of both lungs in which the fibrinous exudate is .5 to 1 cm. in thickness. This exudate adheres to the chest wall in a number of places. There are 200 to

and apical lobes of both lungs are in a state of reddish-gray consolidation. Upon section, a yellowish-white fluid exudes from the bronchi and bronchioles. Nearly the entire diaphragmatic lobes of both lungs are somewhat congested but are consolidated only in their anterior portion which is covered by varying amounts of a fibrinous exudate. The mediastinal lobe appears the same as the right cardiac and apical lobes. The mediastinal and bronchial lymph nodes are only slightly enlarged.

Microscopic examination:

There is a very marked fibrinous pleurisy with much thickening of the pleura (Fig. 29). The interlobular septa are very edematous and contain many distended lymphatics. Both the interlobular septa and the lymphatics contain large numbers of gram positive organisms. In a number of acute areas, there is a zone bordering on the interlobular septa in which there is a marked cellular reaction. (Fig. 60). The cells of the exudate of the bronchi and bronchioles are necrotic and the exudate contains large numbers of organisms. In many areas, an entire lobule appears necrotic (Fig. 59). In the left apical lobe, there is an abscess just beneath the pleural surface approximately 5 mm. in diameter. several lobules, there is some hemorrhage into the alveolar sacs but for the most part, the alveoli are filled with a fibrinous or cellular exudate containing, in many instances, large numbers of organisms (Figs. 73 and 74). There are no

productive tissue changes in the walls of the alveoli. The interlobular septa surrounding necrotic lobules show considerable edema and thickening. The lymph sinuses do not show any marked changes.

Bacteriological examination:

A hemolytic streptococcus of the beta type and staphylococcus aureus were isolated. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #25 (Autopsy 3263), a male Holstein four months old which had a pneumonia of two weeks duration prior to death.

Gross examination:

The cardiac and apical lobes of both lungs are in a state of yellowish-gray consolidation. Many of the grayish areas are 3 or 4 mm. in diameter. In the apical lobes of both lungs, these small areas tend to converge into one large area. The anterior one-third of both the diaphragmatic lobes are consolidated and are very much the same as their corresponding apical and cardiac lobes. In most of the lobes, there are scattered areas of normal parenchyma (Figs. 7 and 8). In the posterior portion of the right diaphragmatic lobe, there are several large emphysematous areas (Fig. 7). The pneumonia in the left lung seems of slightly longer duration than the right although the difference is not pronounced. The mediastinal and bronchial lymph nodes are very swellen and edematous.

Microscopic examination:

The pleura is not thickened. The interlobular septa do not show any thickening nor are the lymphatics distended. Only a very little bronchitis is noted. In some areas, there is considerable cellular infiltration of the respiratory bronchioles which is accompanied by atelectasis of the adjacent alveoli (Figs. 35 and 45). The terminal and respiratory bronchioles of the involved portions of the left lung and the cardiac and apical lobes of the right lung show considerable productive tissue changes in their mucosa. There is much thickening of the alveolar walls due to productive tissue changes. For the most part, the alveoli are filled with polymorphonuclear leucocytes and in many areas appear lined with cuboidal cells (Fig. 68). Areas in which the alveolar walls show the most thickening are for the most part adjacent to terminal or respiratory bronchioles similarly affected. In the right cardiac lobe there is an abscess 1 cm. in diameter just below the pleural surface (Fig. 57). The lymph nodes appear some what edematous.

Bacteriological examination:

A gram negative rod, probably Bacillus coli communior, was isolated. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #26 (Autopsy 3266), a male Holstein three and one-half months old which had been reported as having pneumonia of a week's duration prior to death.

Gross examination:

There is a reddish-gray consolidation of the cardiac and apical lobes of both lungs. About one-third of the left diaphragmatic lobe is in a state of grayish consolidation while approximately two-thirds of the right diaphragmatic lobe is more of a reddish consolidation. The cardiac and apical lobes of both lungs and the consolidated portion of the left diaphragmatic lobe show considerable atelectasis. The mediastinal lobe is entirely consolidated and looks very much the same as the pneumonic portion of the left diaphragmatic lobe. The mediastinal and bronchial lymph nodes are very much congested.

Microscopic examination:

The pleura is unaltered. The interlobular septa are slightly thickened and in a number of areas contain distended lymphatics. In the pneumonic lobes, there is evidence of injury to the small bronchi and the terminal and respiratory bronchioles. There is an acute bronchitis of the small bronchi with much desquamation of the epithelium. The terminal bronchioles are practically the same. This is accompanied by a slight infiltration of mononuclear phagocytes into the mucosae of the involved bronchi and bronchioles. In several areas, there are marked productive tissue changes in the wall of the respiratory bronchioles (Fig. 52). The capillaries of the alveolar walls are very much congested and the alveoli contain many mononuclear cells. The lymph sinuses do not show any marked changes.

Bacteriological examination:

A staphylococcus very much like staphylococcus aureus was isolated from this case. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #27 (Autopsy 3286), a male Holstein five months old which was reported as having a pneumonia of five days' duration prior to death.

Gross examination:

The pneumonia involves the cardiac and apical lobes of both lungs. Approximately two-thirds of the left diaphragmatic lobe and about one-third of the right diaphragmatic lobe are consolidated. All the areas of consolidation are grayish-red in appearance. The mediastinal lobe shows the same gross appearance as the left cardiac. The extent of pneumonia is shown in Figs. 9 and 10.

There are a number of abscesses 4 or 5 mm. in diameter in the cardiac and apical lobes of both lungs, the majority of which are noted only upon section. The largest abscess observed in this case is in the lower posterior part of the left apical lobe (Fig. 9). Upon section, many grayish-white areas can be seen, suggesting abundant cellular exudate. A bloody purulent exudate exudes from the small bronchi when sectioned. The mediastinal and bronchial lymph nodes are slightly enlarged.

Microscopic examination:

The pleura is slightly thickened. There are productive

tissue changes in the walls of many of the bronchi and terminal bronchioles. This is accompanied by considerable cellular infiltration of the stroma of the mucosa. The predominating cell is the mononuclear phagocyte. An occasional polymorphonuclear leucocyte may be seen. Most of the bronchi and bronchioles contain in their lumina a cellular exudate of mononuclear phagocytes and some polymorphonuclear leucocytes. In the areas of longest duration, the alveolar walls are slightly thickened by an increase in number of fibroblasts. The polymorphonuclear leucocyte predominates in the cellular exudate in the alveolar sacs. The lymph sinuses of the mediastinal and bronchial lymph nodes contain many polymorphonuclear leucocytes.

Bacteriological examination:

A staphylococcus resembling staphylococcus aureus was isolated. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #28 (Autopsy 3290), a female Holstein six months old which was reported as having a pneumonia of two weeks' duration prior to death.

Gross examination:

There is a reddish-gray consolidation of the right apical and cardiac lobes. Approximately three-fourths of the left apical and cardiac lobes are in a state of reddish-gray consolidation. An area of normal lung can be seen along the dorsal border of the left apical and cardiac

lobes (Fig. 11). In the anterior portion of the cardiac lobe and nearly at the junction of the right cardiac and apical lobes, a lighter colored, somewhat constricted area can be seen (Fig. 11).

Upon section of the lobe at this point, many abscesses appear on the cut surface (Fig. 12). The diaphragmatic lobes of both lungs are approximately one-half consolidated. There are a few areas of a fibrinous pleurisy. The mediastinal and bronchial lymph nodes are very much enlarged and upon section, a yellowish fluid exudes from the cut surface.

Microscopic examination:

The pleura is thickened. The interlobular septa are very edematous and contain many distended lymphatics. There is an acute bronchitis accompanied by considerable desquamation of the epithelium. In the terminal and respiratory bronchioles of the cardiac and apical lobes, there is some evidence of productive tissue changes in their mucosa. alveoli contain an exudate made up of mononuclear phagocytes, polymorphonuclear leucocytes and in some areas considerable fibrin is noted. The alveolar walls are somewhat congested. The involvement of the alveoli appears to radiate from the nearest respiratory bronchiole (Fig. 36). In several small areas, there is much thickening of the alveolar wall due to productive tissue changes (Fig. 69). A large number of giant cells are present in the pneumonic areas. Several small abscesses are present in some areas. The lymph sinuses of the mediastinal and bronchial lymph nodes

contain some polymorphonuclear leucocytes.

Bacteriological examination:

A hemolytic streptococcus of the beta type, a staphylococcus, and a gram negative rod, probably Bacillus coli communior, were isolated. For their morphological, cultural, and biochemical characteristics see Table I and II.

Case #29 (Autopsy 3297), a female Guernsey seven weeks old which had a pneumonia of six days' duration before death. The symptoms noticed the first day were a marked dyspnea, and accelerated respiration. At this time, 300 cc. of citrated blood were given intravenously.

Gross examination:

There is a reddish-gray consolidation of the cardiac and apical lobes of both lungs. The right diaphragmatic lobe is approximately one-half consolidated while only a small portion of the left diaphragmatic lobe is affected (Fig. 13). The mediastinal lobe shows some slight congestion toward its hilus. There is more atelectasis of the cardiac and apical lobes than of either of the diaphragmatic lobes. The pneumonic portions of the left lung appear of slightly longer duration than the same areas of the right lung. The mediastinal and bronchial lymph nodes are very much congested but are not very much enlarged when compared with other cases.

Microscopic examination:

The pleura is thickened (Fig. 28). The interlobular

septa are very edematous and contain many distended lymphatics. Occasionally in these lymphatics and the surrounding edematous septa, there are large numbers of gram positive cocci which are very similar to those of Case #24. There is considerable atelectasis in all the pneumonic portions of the lung. There is a very severe and acute bronchitis with much injury to the mucosa. A severe bronchiolitis is noted in the terminal and respiratory bronchioles. There are many patchy-like areas, which appear necrotic and contain large numbers of organisms. In many of these areas, there is a small bronchiole in the center which is usually somewhat necrotic. In a few areas, an entire lobule appears to be necrotic. In some instances where this has taken place, the interlobular septa show many productive tissue changes. The lymph sinuses of the lymph nodes contain a small number of polymorphonuclear leucocytes.

Bacteriological examination:

A streptococcus of the beta type was isolated from this case. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #30 (Autopsy 3301), a female Holstein three weeks old which had a pneumonia of two weeks' duration before the animal died.

Gross examination:

There is a reddish consolidation of the right apical

and most of the right cardiac lobe. There is only a slight congestion of the left apical lobe but a grayish consolidation of the left cardiac lobe. Only a small portion of the left diaphragmatic lobe is consolidated (Fig. 14). The consolidated area in the left lung is of decidedly longer duration than the pneumonia in the right lung. The right diaphragmatic lobe is consolidated in the anterior one-third. There are several normal lobules in this consolidated portion. The mediastinal and bronchial lymph nodes are very much enlarged and upon section are very edematous.

Microscopic examination:

The pleura is unchanged. The interlobular septa are only slightly edematous with only a few distended lymphatics. The small bronchi in many areas do not show any marked changes. In approximately one-half of the terminal bronchioles, there is a severe bronchiolitis accompanied by much desquamation of the epithelium. The terminal and respiratory bronchioles contain a cellular exudate consisting mostly of mononuclear phagocytes, a very few polymorphonuclear leucocytes and some desquamated epithelial cells. The alveoli in most of the pneumonic areas are filled with a cellular exudate consisting of mononuclear phagocytes and some polymorphs. In the more acute areas, the alveolar exudate is nearly all fibrin with some serous exudate. There is considerable atelectasis in the cardiac and apical lobes. The lymph sinuses of the bronchial lymph nodes contain many poly-

• • • • morphonuclear leucocytes.

Bacteriological examination:

A hemolytic streptococcus and a gram negative short rod, probably Bacillus coli communior, were isolated. For their morphological, cultural, and biochemical characteristics see Tables I and II.

Case #31 (Autopsy 3314), a male Holstein four months old which had a pneumonia of eighteen days' duration prior to death.

Gross examination:

There is a grayish consolidation of the apical and cardiac lobes of both lungs and anterior one-third of the right and left diaphragmatic lobes. The pleural surface is studded by many small abscesses ranging in size from 1 to 2 mm· to 1 cm· (Fig. 15). These abscesses are present both on the dorsal and ventral surfaces of the lungs as shown in Figs. 15 and 16. There are numerous areas on the pleural surface which appear necrotic. In part of the consolidated portion of the diaphragmatic lobe, there are several areas of shorter duration. The entirely consolidated mediastinal lobe has many small abscesses on its pleural surface (Fig. 16). The mediastinal and bronchial lymph nodes are very much enlarged and upon section, a yellowish fluid exudes from the cut surface.

Microscopic examination:

The pleura is slightly thickened. The interlobular

septa contain a number of distended lymphatics (Fig. 70). In the cardiac and apical lobes of both lungs, there are considerable productive tissue changes in the walls of the terminal and respiratory bronchioles. This is also noted in the walls of the alveoli (Fig. 65). A large number of the small bronchi and bronchioles contain a cellular exudate consisting mostly of polymorphonuclear leucocytes and a few mononuclear phagocytes. In several small areas, there is an organization of the alveolar exudate (Fig. 66). Many alveoli show an increase in the number of fibroblasts in the alveolar wall (Fig. 67). In the diaphragmatic lobe of the right lung where the pneumonia is more acute, there is some congestion of the alveolar walls, also some serous exudate into the alveoli. There are many small encapsulated abscesses and necrotic areas in any of the pneumonic portions of either lung (Fig. 56). The lymph sinuses of the lymph nodes contain large numbers of polymorphonuclear leucocytes.


Bacteriological examination:

A staphylococcus and a gram negative rod, probably
Bacillus coli communior, were isolated. For their morphological, cultural, and biochemical characteristics see
Tables I and II.

DISCUSSION

The frequent occurrence of calf pneumonia in a large number of calves presented for autopsy was the motive for this work. The fact that the pneumonia was a bronchopneumonia seemed well established, yet a study of calf pneumonia from a pathological standpoint appeared incomplete.

In regard to the gross lesions, there were a number of variations which should be mentioned. Nocard (1) reports a pneumonia in calves with much abscessation and cavity formation in the pulmonary tissue. In the thirty cases considered in this study, the above conditions were not encountered with the exception that in seven cases there were numerous small abscesses without cavity formation (Figs. 9, 10, 12, 15, and 16). With a few exceptions, the cardiac and apical lobes are entirely consolidated (Table IV). There was considerable variation in the extent of consolidation noted in the diaphragmatic lobes of either lung. The gross picture of pneumonia in the left lung was consistently one of longer duration than the pneumonia of the right lung. The cardiac and apical lobes of the left lung and the pneumonia portion of the diaphragmatic lobe were, in cases of longer duration, a grayish-yellow color and appeared quite atelectatic as compared with the pneumonic lobes of the right lung of the same animal. The pneumonia in the left diaphragmatic lobe in many cases does not involve as much of the lobe as the usually more acute consolidated portion of the right lobe. However, in one case, the entire left



lung was consolidated and a small portion of the right diaphragmatic lobe was not pneumonic (Fig. 2). It is of interest to note the difference between Fig. 2 and Fig. 14. In the case illustrated in Fig. 2, it was surprising that the animal lived as long as it did while in Fig. 14 only a small portion of the lobes were consolidated. In another case, there is a marked fibrinous pleurisy, the extent of which is shown in Fig. 6. The remaining cases show varying degrees of pleurisy yet not as extensive as the case mentioned above. The posterior mediastinal and bronchial lymph nodes are very much enlarged in twenty-three of the thirty cases. When numerous petechial hemorrhages are present in the lymph nodes, they are as a rule more numerous in the mediastinal nodes than in the bronchial lymph nodes which were enlarged and yellowish-white in color.

Carpenter and Gilman (3) in their study were primarily concerned with the bacteriology of calf pneumonia although some work was reported on the pathology. The bacteriological findings in this investigation compare well with the work of Carpenter and Gilman. The organisms isolated most consistently are; namely, a streptococcus of the beta type, Bacillus coli communior, and a staphylococcus, usually staphylococcus aureus. Streptococci were isolated in thirteen cases. In four of these, the streptococcus is the only organism isolated. In five cases, it is associated with the Bacillus coli organism and in four cases with a

staphylococcus. In eleven of the twenty-four cases cultured Bacillus coli communior was isolated. In five cases, it was associated with a staphylococcus and in five cases it was isolated with a streptococcus. The staphylococcus was isolated in four cases associated with a streptococcus and in five cases with the Bacillus coli organism. In four cases, it was the only organism isolated. A streptococcus and staphylococcus were present in the majority of cases in which many necrotic and abscessed areas occurred.

There was considerably variation in the microscopic pathology in a number of the cases studied. Nineteen cases of the thirty showed a thickening of the pleura, thirteen of which were accompanied by an edematous interlobular septum which contained many distended lymphatics. thickening of the pleura was due to a fibrinous pleurisy and what appeared in many instances as productive changes in the pleura. Twenty-seven of the cases showed an acute bronchitis which ranged in severity from a moderate mononuclear infiltration of the mucosa to a marked congestion with much desquamation of the epithelium. Only one case showed any evidence of a chronic bronchitis. In most instances, the acute bronchitis was accompanied by much desquamation of the epithelium and congestion of the mucosa. In sixteen cases, there was evidence of a chronic bronchiolitis which was manifested by marked productive tissue changes (Figs. 51 and 52) in the stroma of the

mucosa and sometimes the peribronchial tissue. Fourteen of these cases were associated with an acute bronchitis. The above would indicate that the infectious agent probably affected the respiratory bronchioles first. lesions in the respiratory and terminal bronchioles were similar to those mentioned by Boyd (9) and MacCallum (10) in their descriptions of bronchopneumonia especially a streptococcal pneumonia. In a streptococcal pneumonia there is usually more injury to the small bronchioles and surrounding alveoli than to the bronchi. The pathology involving the alveoli varies considerably throughout the thirty cases studied. In eight cases of pneumonia there was a very marked atelectasis although in all cases, some evidence of atelectasis was usually present. Nine cases of the thirty show productive tissue changes of variable extent in the alveolar walls. In those pneumonias of any appreciable duration productive tissue changes were more in evidence in the apical and cardiac lobes. This condition seems slightly more pronounced in the left apical and cardiac lobes. The above conditions also hold true for the productive tissue changes in the small bronchioles. majority of alveolar walls were usually very much congested yet a few contained large numbers of polymorphonuclear leucocytes and some mononuclear cells. In those cases in which the pneumonia was of long duration, some alveoli will become lined with cuboidal cells (Fig. 68). The predominating alveolar exudate was cellular as shown in




Table IV. In five acute cases, the exudate was predominantly serous. It should be understood that serous and fibrinous exudate were present even though the cellular exudate predominates. In one case, there was an organization of the alveolar exudate (Fig. 66). In seven cases out of the thirty, there was much necrosis and abscessation.

SUMMARY

- 1. Thirty cases of calf pneumonia were studied in gross and microscopically for pathological changes. Twentyfour of these cases were studied also from a bacteriological standpoint.
- 2. Three distinct variations in the pathology of these cases are brought out, (1) an acute pneumonia which progresses very rapidly and the animal dies in several days without evidence of any productive tissue changes but usually with a serous exudate filling the alveolar sacs, (2) an acute pneumonia superimposed on a previous bronchiolitis or chronic bronchopneumonia in which a large part of both lungs are consolidated (Fig. 2). This is sometimes accompanied by a marked fibrinous pleurisy as in Fig. 29, and (3) the cases in which large numbers of small abscesses are present (Figs. 9, 12, 15, 16, and 56).
- 3. That the pneumonia is consistently bilaterial has been well established (Table IV).
- 4. That the pneumonia usually begins in the apical lobes has been brought out in gross descriptions and by the microscopic study.
- 5. Consistently the pneumonia of the left lung is of longer duration than the pneumonia of the right.
- 6. The suggestion is made that there is injury to the respiratory bronchioles prior to a bronchitis.

7. The isolation of a beta type streptococcus was made in thirteen of the cases which were cultured; the isolation of a Bacillus coli organism was made in eleven cases and the isolation of a staphylococcus in thirteen cases. Either of the first two organisms was considered as the probable causative agent from a bacteriological point of view.

Table I

Bacteriological Determinations

					-51									
St Mot ST Chn Coc Rod D L ML S I SL Hemo Hem	Gelatin Stab		Growth 7	No Growth 7 da		Papillate	Stratiform			Stratiform		Growth 7	Growth 7	Growth
Grm Mot ST Chn Coc Rod D L ML M S I SL Hemo			Grayish Thite	.Grayish . White		Grayish	Grayish Green	Sparse Beaded	Sparse Beaded	Grayish White		Sparse Beaded	• Sparse Beaded	• • •
Grm St Mot ST Chn:Coc Rod D L MI M S I SL - Mot SH Chn:Coc Rod D L MI M S I SL - Mot SH Chn:Coc Rod D L MI M S I SL - Mot ST Chn:Coc Rod B	Non		Non Heno	Non Hemo		Non Hemo	Non Hemo	• • •	• • •	Non Hemo		• • •	• • •	Non Hemo
Grm St Mot ST Chn Coc Rod D L ML M S I - Mot ST Chn Coc Rod D L ML M S I - Mot ST Chn Coc Sh	нешо			• • •		• • •	• • •	• Hemo	Hemo	• • •		нешо	Нешо	
Grm St Mot ST Chn Coc Rod D L ML M S - Non Sh Chn Coc Rod D L ML M S - Mot SH Chn Coc Rod D L ML M S - Mot SH Chn Coc Rod D L ML M S - Mot ST Chn Coc Rod D L ML M S - Mot ST Chn Coc Rod D L ML M S - Mot ST Chn Coc Rod D L ML M S - Mot ST Chn Coc Rod D L ML M S - Mot ST Chn Coc Rod D L ML M S - Mot ST Chn Coc Rod D L ML M B - Mot ST Chn Coc Rod D L ML R C - Mot ST Chn Coc Rod D L ML R C - Mot ST Chn Coc Rod D L ML R C - Mot ST Chn Coc Rod D R D R P R C - Mot ST Chn Coc Rod D R R R R R R R R R R R R R R R R R R	SL		,	•		•	9	+	+	•		+	#	+
Grm St Kot ST Chn Coc Rod D L ML M + Mot SH Chn Coc Rod D L ML M + Mot SH Chn Coc Rod D L ML M + Mot ST Chn Coc Rod D L ML M + Mot ST Chn Coc Rod D L ML M + Mot ST Chn Coc Rod D L ML M + Mot ST Chn Coc Rod D L ML M + Mot ST Chn Coc Rod D L ML M + Non Sh Chn Coc Rod D L ML R + Non Sh Chn Coc R H + + + + + + + + + + + + + + + + + +	Н	i i				Ф	€	1	1	Ф	i	1	1	+
Grm St Chn Coc Rod D L Hot ST Chn Coc Rod D L Hot ST Chn Coc Rod D L Hot ST Chn Coc Hot Sh Chn Coc Hot Sh Chn Coc Hot Sh Chn Coc		Case		+ •••	Case	• • •	• • •	+	• • • • +	• • •	Case	+	+	1 • • • • +
Grm St Chn Coc Rod D L Hot ST Chn Coc Rod D L Hot ST Chn Coc Rod D L Hot ST Chn Coc Hot Sh Chn Coc Hot Sh Chn Coc Hot Sh Chn Coc	MT		+ • •	+		e	•	+	• • • •	⊕		+	#	• • •
Grm St Mot ST Chn Coc Rod + Mot ST Chn Coc Rod + Mot SH Chn Coc - Mot SH Chn Coc - Mot ST Chn Coc - Mot ST Chn Coc - Mot ST Chn Coc Sh - Mot ST Chn Coc	• • • •		• • •	+		⊕		• • •	+	•••		+		•••
Grm St Mot ST Chn Coc Hot ST Chn Coc Mot SH Chn Coc Mot ST Chn Coc Non Sh Chn Coc Not ST Chn Coc Mot ST Chn Coc	D		+	+		•	•	+	+	⊕		+	+	+
Gra St Mot ST Non + Mot ST Non + Mot ST + Mot ST + Non + Non ST + Mot ST +	Rođ					Sn Rod	Sh Rod	• • •		Sh Rođ				• • •
Gra St Mot ST Non + Mot ST Non + Mot ST + Non + Non ST + Non	306			Soc				၁၀၀	300			၁၀င	300	300
Gra St Mot ST Non + Mot ST Non + Mot ST + Non + Non ST + Non	hn.		Shu	sh Jhn (કુલ	۲. بر	Sh.	Sh Shn	• • •		Sh n	Sh n	Sh Shn:
Gran Mot - Mot - Mot - Mot - Mot - Mot - Mot - Mot - Mot - Mot - Mot - Mot - Mot - Mot - Mot - Mot - Mot - Mot - Mot - Mot - Mot	• • • •			• • •		• • • •	•	• • •	• • •	•••		• • •	• • •	•••
			Non.			Mot	Mot							Non Mot
Cull No. 11 2 4 4 6 9 11 11 11 11 11 11 11 11 11 11 11 11 1	Grm St		+	+		1	•	+	+	1		+	+	+
	ተ :													
	No No					4	<u>س</u>] ²	7

I and the second		• • • • • • • • • • • • • • • • • • • •		
			•	
	• • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •		• • •
	• • • • • • • • •			
			1	
		• • • • • • • • • • • • • • •		
	• • • • • • •			
	• • • • • • • • • • • • • • • • • • • •			
	• • • • • • • • • • • • • • • • • • • •			• • •
1				• • • •
		- • • • • • • • • · · · · · · · · · · ·		
		the transfer to the second sec	• • •	• •
1				
			•	. •
			• • • • • • •	
		•••••	• • • • • •	
4				
ļ		• • • • • • • • • • • • • • • • • • • •		• • • •
				• • • •
•				
1				

 $(x_1, \dots, x_n) \in \{x_1, \dots, x_n\} \cap \{x_1, \dots, x_n$

	1	٠٠١		•••	. • • 1	۱••۹				ı•••ı	<u> </u>	• • •	• • •	۲۰۰۱	! 1	 •••	
	Gelatin	Stab		No Growth	Slight	: Slight	No Change		Surface Growth	No Growth		. Stratiform	• Crateriform	Crateriform		No Growth	No Growth
	Agar	Slant		Sparse Beaded	Grayish White	Grayish White	Sparse		whitish Yellow	Sparse Beaded		Yellowish Luxuriant	Grayish Thite	Grayish White		Sparse Beaded	Sparse Beaded
		Нешо			Non Hemo	Non Hemo	Non Hemo		иоN Нэто			Non	Non Hemo	Non Hemo		Non Hemo	
	• • •	Нешо		. Hemo		• • •			• • •	Нешо		• • •					Hemo
	•	SI		+	Ф	+	+		•	+		1	1			+	#
		н	1	1	⊕	١	1	જ	•	•	53	1	,		4	+	
		တ	#11#	+	æ	•••	1	#15	+	+	#13	+	• • •	• • •	e #14	+	***
		×	Case	١	⊕	+	+	Саѕе	#	+	Case	+	,	,	Case	1	+
	• • •	벌		1	•	+	• • •		+	-#		+	1	•		• • •	**
1		1		• • •		• • •	• • •		• • •	• • •		• • •	• • •			• • •	•••
	- •	A		+	•	+	+		• • •	+		+	+	+		+	•••
		Rod		• • •	Sh Rod	Rod :	• • •		• • •	•••		• • •	Sh Rod	Sh Rođ		•••	• • •
	• • •	Coc		၁၀၁			Coc		Coe	Coc		Coa				င္ပဝင	ဝဝဝ
]-	- 3	Chn		Sh Chn	•••	Sg	Pr		10:	••••		Sg	• • • •			Sh Chn	Sh
		Si		ST			ST		贸	ST						ST	ST
/- -	• _	Mot		Non Mot	Mot	Non Mot	Non Mot		Non	Non Mot		Non	Mot	Mot		Non Mot	Non Mot
	3	St		+	,	+	+		+	+		+	,			+	+
101	_	NO.		12	13	14	15		16	17*		18	19	20		21	22

able I Con

Stab Crateriform Crate riform Stratiform Stratiform Stratiform Stratiform Growth Papillate Gro wth Growth Growth Growth Gelatin S No No 0N No 8 Sparse Beaded Grayish Yellow Sparse Beaded Grayish Yellow Gray to White Grayish White Yellowish White White Yel lowish Grayish Grayish Grayish Grayish Non Hemo Non. Non Non Hemo Hemo. Non Es Hemo Non Hemo Hemo Non Hemo Нешо Нешо Eemo Нешо Non Нешо Non Non Non Non Hemo. : Hemo SI **⊕** + + Ф + Ф Н i ı **⊕** Φ + ı ⊕ #12 ø **#** #1 മ ı + • + + Ф + Case Case Case + **⊕** × + # **B** 1.1.1 # + + + ⊕ Rod: D + Ф + + • + **Sh** Rođ Sh Rod Rod Rod Rod Sh Rod Sh Rođ 00 C00 Coc . Coc ၁၀၀ Coc ပ္ပိုင္သ Chn Chn Sg .Sg :01 . C1 Sh P ES ST 띬 ST 띬 Mot: Mot Mot: Mot. Non Non Non Non Mot Mot Non Mot Non Mot Mot Non Mot Non Non Cul. Gra. St . • + No 33 23 **8**8 34 24 88 29 30 25 27 3

Table I Cont.

Stab Stratiform Stratiform Stratiform Stratiform Stratiform Stratiform Stratiform Stratiform No Growth Growth No Growth Gelatin ON. Sparse Beaded Sparse Beaded Sparse Beaded Grayish White Grayish White White Slant White Abund ant Abundant Grayish Gray ish Yellow Yellow Agar • • • Non Hemo Non Hemo Non Hemo Hemo Non Hemo Non Hemo Non Hemo Hemo Hemo Непо Non Hemo Non Non Non Non Нешо Hemo SL + Ф ₩ • H 1 **•** + ⊕ Н #19 #20 #18 #21 • + 1 • + • 02 1 Case Case Case Case • • ì + × • **⊕** 月 • • H + + • Ф • • + • + + Ф A • Sh Rod Sir Rod Rod Sh Rođ Rod Rod SP SP Coc Coc Coc Goo Coc ဝဝဝ Sh Chn Sh Chn . CJ 58 Sg Pr S P C SH ST SH ST R Mot. Mot Non Wot Non Mot Non Mot Non Non Non Mot Mot Mot Non Mot Mot E ı St 1 + + + + + No. 46 44 43 45 42 35 36 37 39 40 41

Table I Cont.

• • •			,	
• • •				
				Į.
				r
	• • • • • •	• • • • • • • •	• • •	
4				
	:			
• • • •		• • • • • • • • •		
• • •	• • • • •			
• • -		• • • • • •		• • • • • • • • • • • • • • •
• • •				
• • •				• • • • • • • • • • • •
_				
•				
1			0	4
•	• • • • • •			

	ŀ								,	•	•	•	•	•	•	•	
Cul.	Gra	Mot	ST	Свп	Go	• • • • •	Rod : 1	7 0	H H	*	• • • • • • • • • • • • • • • • • • •	н • • • •	SL	Нешо	NOII Hemo	Agar Slant	Gelatin Stab
47	+	Non Mot	SH	61			•••	•‡•					#		Non Hemo	Yellow	: Stratiform
										Case		#22 #					
8	+	Non				Ř	Rod		•		•••			•	Non Hemo	Grayish Yellow	. No Growth
49	1	Mot				 	Sh Bod	•	•		.:.			• • •	Non Hemo	Grayish Yellow	. No Growth
							,			Case		#23					
50	+	Non	ST	Sh	Goc		•••	*	•••	*	* • <u>;</u> •		+.	нешо		Sparse Beaded	No Growth
										Case		#24					
51	+	Non		Sh	COO		•••	;			*	1	+	. Hemo	••	Sparse Readed	No Growth
52	+	Non	.	80	1	١	Rod	• • •	+	ļ	 	+	+	•••	Non Hemo	Whiti sh	Papillate
53	+	Non	HS	12	Coc		•••	 • • •					+	• • •	Non Hemo	Yellow	Stratiform
										Case	1	#25					
54	1	Mot		SR		S C	Sh . Rod	•	•	1	•	•	⊕		Non Eeno	Grayish White	No Growth
55	+	Mot		Sg		N E	Sh Rođ				•••			• • •	Non Hemo	Yellow1sh	. Stratiform
										Case		#26					
56	+	#ON TO	₩ ₩	Sh	C.		•••	• ‡ •			• • •	•••	1	• • •	Non Hemo	Sparse Beaded	No Change
				- 1	1												

Stab Crateriform Stratiform Stratiform Stratiform Stratiform Papillate . No Growth No Growth Pap 111ate Growth Celatin S N Sparse Beaded Sparse Beaded Dirty Gray Slant White White Fhite Grayish Grayish Grayish Grayish · Yellow Yellow · Yellow Agar Non Hemo Non Hemo Non-Hemo Non Hemo Non Hemo Non Non Hemo Hemo Hemo Non Нешо Hemo. Hemo + SL н 1 #28 #29 #30 #31 427 + 1 **0** + + Case Case Case Case Case + × + + 以 + H + 1 + + + 1 + + A •• + + + 1 Rod Sh Rod Rod Sh Rođ Coo Sh Chn Coo Coc Coo Coc · 000 · Coc Coc Chn Sh Chn Chn Sg. SH C1 : 01 13: に Sg ST ST 띪 S TON TO Mot Non Non Mot Mot Non Non Non Non Mot Mot GT st + Tno No 60 63 65 66 629 57 64 61

Table 1 Cont.

• • •					• • • • •
• • •	••••	• • • • • •			• • • • •
		• • • • • • •	•••		
1			•••••••	• • •	
	• • • • •			· · ·	• • • • •
• • •	• • • • • •	• • • • • • •	• • • • • • • • •		• • • •
	· • • • • •			• •	
• • •	• • ,• • •	• • • • • •		• •	
• • •				• • •	
. 	• • • • •		· · · · · · · · · · · · · · · · · · ·	• •	• • • • •
• • •	• • • • •	• • • • • • •	• • • • • • • • •		• • • • •
• • •			• • • • • • • • • • •	• •	• • • • •
	• • • • •		• • • • • • • • • •	• • •	• • • •
			• • • • • • • • • • • •		
• • •	• • • • •		• • • • • • • • • • •	• • .	
	• • • •			• • •	
- · ·	• • • •				

.

. . . .

. . .

Salicin

Table 1 Cont.

1	• • •	•••	 •••	1
· Gel atin	Stab	· Crateriform	• Crateriform	No Growth
· Agar	Slant	Non Dirty Hemo Gray	Dirt	Grayish White
Non	mo Hemo	Non Hemo	Non Hemo	Non Hemo
• •	SI Hemo	•	0	1
	MI. N. S. I	0 • •	0 0 0	
-	널	•	0	+
	I A	•	0 0	+
	Rod	Rod	Rod	Rod
	Coc . Rod			
	Chn	Sg	Sg	Sg
• •	ST		• • •	•••
•	. Mot	. Mot	Mot	Non Mot
.0.	St	•		+
Can'l	No	67	68	69

Key to Abbreviations

Gram's stain reaction Motility	Streptococcus	Chain; Sh Short	Coccus form	Rod form	Dextrose	Lactose	Maltose	Manni te	Sucrose	Inulin
Grm. St. Mot.	ST.	Chn.	Coc	Rod.	.	i.	ML.	М.	ა	•

72. Culture #17 used in Fig.

Culture
Negative or no reaction
Positive or acid formation
Acid and gas
Partial Single In pairs Hemolytic Non-hemolytic Staphylococous Clusters SL.
Sg.
Pr.
Hemo. SH. Cl.

. $(x,y) = (\mathbf{v}_{i}, x_{i}, \mathbf{v}_{i}, \mathbf{v}_{i}, \mathbf{v}_{i}, \dots, \mathbf{v}_{i}, \mathbf{v}_{i}) = (\mathbf{v}_{i}, \mathbf{v}_{i}, \dots, \mathbf{v}_{i}, \mathbf{v}_{i}, \mathbf{v}_{i}, \dots, \mathbf{v}_{i}, \mathbf{v}_{i}, \dots, \mathbf{v}_{i}, \mathbf{v}_{i})$ and the second second

Table II

LITMUS MILK REACTIONS FOR CULTURES

CASE #6

Cultures	24 hours	48 hours	72 hours	5 days	7 days
ન જ	Acid but no curd No change The	Reduction, no curd Acid curd Acid curd Soft curd reactions were not consistent until	Acid curd Soft curd Sistent until after	Same Reduction r 48 hours.	Same Same
		CASE #7	47		
4	Acid, no curd	Acid and reduction	Same	Same	Same
ഗ	Acid, solid curd	Acid curd, 1/2 reduction	Digestion of curd	More digestion	Digestion almost
Ð	Slight peptoni-	Peptonization	Complete	Same	Same
4	Acid, no curd	No curd, slight	Same	Soft curd	Same
ω	Acid, no curd	reduction of aye Partial reduction and soft curd	More reduction, soft curd	Ѕвше	Same
		CASE #	6#		
9	No change No change	No change No change	Same Acid slight	Same Soft curd	Same Soft curd,
11	No change	Peptonization	More complete	Peptonization complete	siight reduction Same
		CASE #11	/1.1		
12	No change	No change	No change	Ѕапе	Same

Table II Cont.

Cultures	24 hours	48 hours	72 hours	5 days	7 days
13	Acid curd	Solid curd, 3/4	Acid curd, com-	Same	Same
14	Acid but no curd	Solid curd, 3/4	plete reduction Same	Same	Same
15	No change	reduction No change	Slightly acid, no curd	Ѕаще	Same
		CASE	#12		
16	Acid, no curd Acid curd	Acid, no curd Solid curd, 3/4 reduction (Culture #17 resembles	Same Acid curd, complete reduction very closely #13)	Same Same	Same S am e
		CASE	#13		
18 19 20	No change No change No change	Acid curd Acid curd Acid curd	Acid curd Acid curd Acid curd	Same Same Same	Sаше Sаше Sаше
		CASE	#14		
21	Acid curd soft	Solid curd, 2/3	Acid curd, more	Ѕаше	Same
88	Acid curd soft	Solid curd, 1/2	Acid curd, com-	Same	Seme
೮ ೪ 4	No change Acid curd (Culture	No change Acid curd #21 and #22 seem	No change Acid curd very similar in many	Same Same respects)	Same Same

ays 7 days	Same	Same	Same		curd Same		Seme	Same	4	#26 and #27)		Same	Same	Ѕаше		Same
5 days	Same	Same	Same	Solid	Solid Solid		Same	Same	0	cultures #	S MARK	Same	Seme	Same		Same
72 hours	Acid curd, com-	Acid curd, 2/3	Acid curd solid	curd	Acid curd soft Acid curd soft	E #16	Acid curd reduction	Complete peptoni-	Acid curd, com plete reducti	the reactions of #17	Apto curd	Acid curd		Acid curd	CASE #18	Acid curd, some
48 hours	Some reduction	Solid curd, 1/2	Solid curd, 1/3	No change	No change No change	CASE	Slightly acid	Peptonization	Solid curd, reduction	resembles very closely or casely or casely the case of	Acid գուժ	Acid curd		Acid curd	CAS	Acid curd solid
24 hours	No change	Acid curd soft	Acid ourd soft		No change No change		No change	No change	soft	(Culture #53 r	Actd enrd		Acid curd			Acid curd solid
Cultures	83 123	26	27	82	8 0 80 80 80 80 80 80 80 80 80 80 80 80 80		31	32	33		7.	35	36	37		39

CASE #15

Table II Cont.

•
44
₫
Ö
ŏ
_
H
Н
•
Н
2
ğ
شع

Cultures	24 hours	urs	48 hours	72 hours	5 days	7 days
40	No change	(Հոյ էոբ	Peptonization	Complete peptonization	Seme	Same
			CASE			
41	No change	Q	Peptonization	Complete	Ѕете	Same
4 8 8 8	Acid No change	9 .	Acid Peptonization	peptonization Acid curd Complete	Same Same	Same Same
	(Bot	(Both culture	#41 and #43 resemble	peptonization each other very clos	closely)	
			CASE	#20		
44	No change	Φ.	Peptonization	Complete	Ѕвше	Same
45	No change	(Both	Peptonization cultures resemble each	pepconization Complete peptonization h other very closely)	Same	Same
			CASE	#81		
4 6 47	Acid curd Acid curd	ਾਰ 'ਚ	Acid curd Solid acid curd	Same Same	Same Same	Same Same
			CASE	#22		
4 8 4 9	No change Acid curd	۵. نام	Rennet curd Solid acid curd	Acid curd Acid curd, com- plete reduction	Same	Sаше Sаше

	days	•		eurd 1		~ ~		•		4		_	
	7	Same		Soft Same Acid		Sаше Sаше		Same		Same		Seme	Same
	5 days	Seme		Soft curd Same		Acid curd Same		Complete peptonization		Ѕате		Complete	per contaction Complete peptonization
Case #23	72 hours	Comple te peptonization	CASE #24	Acid slight Acid curd Slightly acid	CASE #25	Acid curd Acid curd	CASE #26	Peptonization	CASE #27	Acid curd	CASE #28	Complete	peptonization Complete peptonization
	48 hours	Peptonization		No change Acid No change		No change Acid curd		Peptonization		Acid ourd		Pep tonization	Peptonization
ont.	24 hours	No change		No change Acid		No change No change		No change		No change		No change	No change
Table II Cont.	Cultures	20		51 52 53		ა გა		56		57		58	S S

			ırd		-71-	,		
7 days	Same Same		Soft curd		Same		Seme Seme	Same
5 days	Same Same		Soft curd Same		Same		Same Same	Same
72 hours	Acid curd Acid curd	CASE #29	Acid slight Acid curd	CASE #30	Acid curd Complete peptonization	CASE #31	Acid curd Acid curd	
48 hours	Acid curd Acid curd	70	No change Acid curd	70	Acid curd Peptonization	70	Acid curd Acid curd	
24 hours	Acid curd No change		No change Acid curd		Acid curd No change		No change Acid curd	Acid curd
Cultures	60		62 63		65 55		66 67	68

Table II Cont.

Chart III.



Ta

*

18

+ + + +

4/

.

		La constitución de la constituci
Case numbers	1 2 3 4 5 6 7 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28	29: 30: 31
Reported duration of pneumonia prior to death. (In days).	4 : 7 : 10 : 20 : 14 : 3 : 10 : 12 : 12 : 14 : 7 : 7 : 7 : 11 : 18 : 15 : 21 : 11 : 2 : 28 : 21 : 14 : 6 : 14 : 7 : 5 : 14 :	6: 14: 18
Bilateral or unilateral (+ or -).	<u> </u>	+ + + +
Left apical lobe (+ if entirely consolidated).	1/2: +: +: +: +: +: +: +: +: +: +: +: +: +:	
Left cardiac lobe (+ if entirely consolidated).	1/5: + : + : + : + : + : + : + : + : + : +	+:+:+
Left diaphragmatic lobe (+ if entirely consolidated).	- 1/4·1/5·1/3·1/3·2/3·1/3·1/4·1/6·1/5·1/5·2/3·1/3· + :1/8·1/3·1/3·2/3·1/3·2/3·1/3· ± :1/3·1/3·2/3·1/2·	
Mediastinal lobe (+ if entirely consolidated).	: - : - : + : + : + : + : + : + : + : +	1/4: + : +
Right apical lobe (+ if entirely consolidated).	1/6 + + + + + + + + + + + + + + + + + + +	+: +: +
Right cardiac lobe, anterior portion (+ if entirely consolidated).		+: 3/4: +
Right cardiac lobe, posterior portion (+ if entirely consolidated).	1-1+1+1+1+1+1+1+1+1+1+1+1+1+1+1+1+1+1+1	+: +: +
Right diaphragmatic lobe (+ if entirely consolidated).	: : : : : : : : : : : : : : : : : : :	1/2.1/3.1/3
Lymph nodes enlarged or edematous (+ or -).	- + + + + + + + + + + + + + + + + + + +	-: +: +
Pleura thickened (+ or -).	1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-	+ - ±
Interlobular septa edematous (+ or -).	1 _ 1 _ 1 _ 1 + 1 + 1 _ 1 _ 1 _ 1 _ 1 + 1 +	+: ±: ±
Lymphatic of septa distended)+ or -).	<u>; </u>	+: -: +
Bronchitis acute or chronic (A, C, or -; C* both A and C).	· A · A · A · A · A · A · A · A · A · A	Δ: -: Δ
Bronchiolitis acute or chronic (A, C, or -).		1: 1: 0
Productive tissue changes in alveolar wall (+ or -).	· · · · · · · · · · · · · · · · · · ·	-: -: ±
Marked atelectasis (+ or -).		+
Exudate S, serous F, fibrinous C, cellular (S, F, or C; F).	:-:s:c:c:r:s:c:c:c:c:c:c:c:c:c:c:c:c:c:c:c	c: c: c
Organization of exudate in the alveoli (+ or -).		: : :
Necrotic areas or abscesses (N, A, or -; A).	N N N N N A	N - A
Bacillus coli communior (+ when present).	* * * * * * * * * * * * * * * * * * * *	+ +
Streptococcus (Beta type) (+ when present).		+: +:
Staphylococcus (+ when present).		: : +
* A O N		

^{* 4} C N C, F, and A, mean that both conditions are present but that the numerator is slightly predominant. * Not examined bacteriologically.

. -. • •

Fig. 1. case 14. The lungs from 2 case of pneumonia of seven days duration. The congested areas are distinctly outlined due to their darker color. Note the lobular nature of the darker note congested areas in the anterior portion of the left dispiragmentic lobs. (1/2 actual size.

Fig. 1. Case 14. The lungs from a case of pneumonia of seven days' duration. The congested areas are distinctly outlined due to their darker color. Note the lobular nature of the darker more congested areas in the anterior portion of the left diaphragmatic lobe. 1/3 actual size.



Figure 1.

Fig. 2. Case 15. The large from a case of preventa of eighteen days' duration. Note that only a small amount of normal lung remains along the forcal posterior part of the right fixthree-matic lobe. 1/3 actual eige.

Fig. 2. Case 16. The lungs from a case of pneumonia of eighteen days' duration. Note that only a small amount of normal lung remains along the dorsal posterior part of the right diaphragmatic lobe. 1/3 actual size.



Figure 2.

Fig. 5. three No. Dee longs from a case of mannonia of the contact for the contact shape the cultrare enlarged and the contact of the characteristics and contact from a contact last the small eras of contactives and large three in the level part of the the carse of large 1.

Fig. 3. Case 22. The lungs from a case of pneumonia of three weeks' duration. Note the extreme enlargement of the bronchial and posterior mediastinal lymph nodes. Also note the small area of comparatively normal lung tissue in the lower part of the right cardiac lobe.

1/3 actual size.



Figure 3.

Fig. 4. On a construction of a social of a

Fig. 4. Case 23. The lungs from a case of pneumonia of two weeks' duration. Note the enlargement of the bronchial lymph node and the lobular appearance of the pneumonia along the borders of consolidated portions of the right and left diaphragmatic lobes. 1/3 actual size.



Figure 4.

Fig. 5. Case 22. A crocs season of the left disphrage water lobe of the cope case as shorn in Fig. 4.

Note the acute concested area mean the large broughish takes. The portion above this congested area is normal while the area below is of grayish white color, which upon close inspection shows a climitly mottled appearance.

1/3 actual size.

Fig. 5. Case 22. A cross section of the left diaphragmatic lobe of the same case as shown in Fig. 4.

Note the acute congested area near the large bronchial tubes. The portion above this congested area is normal while the area below is of grayish white color, which upon close inspection shows a slightly mottled appearance.

1/3 actual size.



Figure 5.

1

Sincapenq in which a resil a surface . As selected . 3 •rin.

Salinable to a cit . Will make teget mission

Brondance base suffer you want in amount increase

•evice lunces in the cit . Contains a contains a second contains.

Fig. 6. Case 24. The lungs from a case of pneumonia of six days' duration. Note the marked fibrinous pleurisy involving the cardiac and apical lobes of the right lung and scattered areas over both lungs. 1/3 actual size.




Figure 6.

Fir. 7. Used 18. Sured a cover of necessaria of two vectors values in Same short. Hotel is enimal of two vectors and sured in the second of the contract of the right of the contract of the right of the contract.

The fire leads of the contract of the right of the reference.

Fig. 7. Case 25. Lungs from a case of pneumonia of two weeks' duration. Note the enlarged bronchial lymph nodes and emphysematous areas in the posterior part of the right diaphragmatic lobe. 1/3 actual size.



Figure 7.

Fig. 8. Sea off. The court follows of the seas lung through the court of the seas lung through the contestence indules of the court season account. If the court season

Fig. 8. Case 25. The ventral aspect of the same lung shown in Fig. 7. Note the occasional lobules that appear normal. 1/3 actual size.




Figure 8.

rminodr winds bug flux to esson. • no esson • Q • minode to distribute and esson of the minode for the following flux of the finite flux of the finite flux of the flux of the






Figure 9.

Fig. 10. Once if. one captal necession to large in Fig. e. force the scrip ones area on the pleural wordens of the local condition.

1050. 1/3 act of size.






Figure 10.

Pinompean to also a orthogonal edit . The control of . If . if the state of the control of the c






Figure 11.

Fig. 13. Case 28. Cross machine from the right cardiac lobe of the last shown in Fig. 11.

This includes the lighter constricted erea mentioned under Fig. 11. Note the large number of abscesses in the cross sections.

1/3 setuch sins.






Figure 12.

Fig. 13. Cars Dr. A lung from a cost of gastronia of cix days darstron. 1/5 actual case.

Fig. 13. Case 29. A lung from a case of pneumonia of six days' duration. 1/3 actual size.



Figure 13.

Fig. 14. case 30. A lun; from a case of prouncils in which there we only a small amount of reddich-gray consolifation. 1/3 actual size,

Fig. 14. Case 30. A lung from a case of pneumonia in which there was only a small amount of reddish-gray consolidation. 1/3 actual size.




Figure 14.

Fig. 15. Case 31. A lung from a case of pneumonia of eighteen days' duration. Note the numerous small abscesses and the one larger one on the pleural surface of the right cardiac lobe. 1/3 actual size.



Figure 15.

Pig. 16. ora 31. The voltral and up on the hong sure. In This 18. Note the headmas show one age. 1/3 actes? Sime.

Fig. 16. Case 31. The ventral aspect of the lung shown in Fig. 15. Note the numerous abscesses. 1/3 actual size.



Figure 16.

- Fig. 17. 1808 1. redica of contains about the skowing strong productions a manufacture decomposition of the strong production of the strong produc
- Fire discount to bigher adjulfication of Wine in Fig. 17.

 caccine a normal respiratory breakings.

 7500.

- Fig. 17. Case 1. Section of normal lung showing a respiratory bronchiole. 120%.
- Fig. 18. Case 1. A higher magnification of Fig. 17 showing a normal respiratory bronchiole.

- -more legistrat a cairora meletar . 3 east . 31 . 217.

 **MOUL & cincular discount broads a disk eloide
 - Fig. 2. Cara 5. Section therity recent bronchepredatants with more extension into the sugmounding alveeler tissue. 130%.

- Fig. 19. Case 5. Section showing a terminal bronchiole with a recent bronchopneumonia. 100X.
- Fig. 20. Case 5. Section showing recent bronchopneumonia with more extension into the
 surrounding alveolar tissue. 110%.




Figure 19.




Figure 20.

Fig. 21. Case 6. An area showing scute connection of the siveclar walls. 150X.

Fig. 22. Case 5. Section showing recent fibrinous exudate in an alveolar duct of a case of

acute bronchopneumonia. 130X.

- Fig. 21. Case 6. An area showing acute congestion of the alveolar walls. 130X.
- Fig. 22. Case 5. Section showing recent fibrinous exudate in an alveolar duct of a case of acute bronchopneumonia. 130X.




Figure 21.




Figure 22.

Fig. 23. Case 10. Section showing soute conjection
the of the alveelar mall and core serous exudation into the alveelar sac. 15cK.

Fig. 24. Case 10. Same section as above but higher mannification. 575 K_{\odot}

- Fig. 23. Case 10. Section showing acute congestion of the alveolar wall and some serous exudation into the alveolar sac. 130X.
- Fig. 24. Case 10. Same section as above but higher magnification. 575X.




Figure 23.




Figure 24.

Fig. 10. Sim S. Septine and proving the world world so will be a septine of the world of the world so well to the world so well the world

fo tud evens as attions so a covered to the state of the

- Fig. 25. Case 6. Section showing an area in which practically all the alveoli contain a fibrinous exudate. 105X.
- Fig. 26. Case 6. Same section as above but of higher magnification. 575X.




Figure 25.




Figure 26.

Fig. 27. Case 18. Section showing a market thickening of the plears with atelectesis of some of the alveoli below the plears! surface. 120X.

Fig. 28. Case 31. Section showing similar condition of the pleura as above, but nove the mononuclear cells in the alventus below the pleura. SSCX.n.

- Fig. 27. Case 18. Section showing a marked thickening of the pleura with atelectasis of some of the alveoli below the pleural surface. 120X.
- Fig. 28. Case 31. Section showing similar condition of the pleura as above, but note the mononuclear cells in the alveolus below the pleura. 550X.




Figure 27.




Figure 28.

lig. 20. Care 16. Caction charing conniderable at a fact.

- Fig. 29. Case 24. Section showing a marked fibrinous pleurisy. Compare this with the gross picture (Fig. 6). 140X.
- Fig. 30. Case 13. Section showing considerable atelectasis. 160X.




Figure 29.




Figure 30.

Fig. 51. Case 82. Section of a bronchus with comparatively normal apithelium but slight infiltration of macrophages into the strome of the mucosa. 170X.

its, 52. Case 22. Same section as above but higher magnification. 600X.

- Fig. 31. Case 22. Section of a bronchus with comparatively normal epithelium but slight infiltration of macrophages into the stroma of the mucosa. 170X.
- Fig. 32. Case 22. Same section as above but higher magnification. 600X.




Figure 31.




Figure 32.

Fire our case the care sectors as above but him and the case of th

- Fig. 33. Case 23. Section showing an acute bron-chitis with much desquamation of the epithelium. Note the congested capillaries.
- Fig. 34. Case 23. Same section as above but higher magnification. 620X.




Figure 33.




Figure 34.

i

Fig. 85. Jase 26. ection -coling a nonclogulum moulus. Noto the selectants between the areas of broadcognechouse. 851.

Pig. 36. Case 23. Section dowing a recent pronchopneumonia. Note how the cellular infiltration appears to registe from the two bros-

- Fig. 35. Case 25. Section showing a bronchopneumonia. Note the atelectasis between the areas of bronchopneumonia. 85%.
- Fig. 36. Case 28. Section showing a recent bronchopneumonia. Note how the cellular infiltration appears to radiate from the two bronchioles.100X.




Figure 35.




Figure 36.



Fig. 57. Case 6. Section showing an scute bronchitis
with much desquenation of the spithelium.
Note the large number of epithelial cells
in the lumen of the bronchus. 105X.

Fig. 28. Case 6. Same section as above but higher magnification showing the desquamated epithelial cells. SECK.




Figure 38.

- Fig. 37. Case 6. Section showing an acute bronchitis with much desquamation of the epithelium.

 Note the large number of epithelial cells in the lumen of the bronchus. 105%.
- Fig. 38. Case 6. Same section as above but higher magnification showing the desquamated epithelial cells. 550X.




Figure 37.




Figure 38.




Fig. 39%, cisc 48.4 Section showing an early bronoffices with considerable inflictivation of mononuclear phagogytes into the strome of the mucosa. 150%.

Fig. 40. Case 19. same as above but higher magni-floation. 600X.




Figure 40.

- Fig. 39. Case 19. Section showing an early bronchitis with considerable infiltration of mononuclear phagocytes into the stroma of the mucosa. 150X.
- Fig. 40. Case 19. Same as above but higher magnification. 600X.




Figure 39.




Figure 40.

- Fig. 13 to the distribution of the state of the
- Fir. 40. Cace le. Suce section as shore. Note the open ever suce of pillumines and injury to the Arithmitistics. Collin.

- Fig. 41. Case 18. Section of a terminal bronchiole with much desquamation of the epithelium.

 130X.
- Fig. 42. Case 18. Same section as above. Note the congested capillaries and injury to the epithelium. 600X.




Figure 41.





Figure 42.



Fig. 43. Case 17. A respiratory bronchiole with much cellular infiltration into the stroma of the mucosa. 120X.

Fig. 44. Case 17. Same as above but higher magni-



F1 ure 44.

- Fig. 43. Case 17. A respiratory bronchiole with much cellular infiltration into the stroma of the mucosa. 120X.
- Fig. 44. Case 17. Same as above but higher magnification. 500X.




Figure 43.




Figure 44.

Fig. 45. Care 21. A terminal bronchiols with any red callular infiltration into the absolute of the mucosa; also note the designation of the critishium. 1907.

- Fig. 45. Case 25. Section of small bronchiole showing considerable lymphocytic infiltration of the peribronchial area. 110%.
- Fig. 46. Case 21. A terminal bronchiole with marked cellular infiltration into the stroma of the mucosa; also note the desquamation of the epithelium. 140X.




Figure 45.




Figure 46.

Fig. 47. Case 10. Section showing a small bronchicle with a zone of marked ceilmler infiltration of the perthronchial tissue. 150x.

Fig. 48. Case 16. Section showing a bronchiole with a necrotic wall and pecrosis of the caucate in the lumen. 150x.

- Fig. 47. Case 10. Section showing a small bronchiole with a zone of marked cellular infiltration of the peribronchial tissue. 150X.
- Fig. 48. Case 16. Section showing a bronchiole with a necrotic wall and necrosis of the cells of the exudate in the lumen. 150%.




Figure 47.




Figure 48.

ATITU MOTO ARBOTT TAIRCIS ADARBS (B) AR. (AF ATIS)

Ifare a in liter out at estat occurs

A.SI (aid) occurs

Fir. 50. Case 9. Sure section as approximate fire the confidention. P4CK.

- Fig. 49. Case 9. Section showing recent productive tissue changes in the wall of a small bronchiole. 105X.
- Fig. 50. Case 9. Same section as above but of higher magnification. 540X.




Figure 49.




Figure 50.

-Outloom (Giftsh of a very roughtsus) - 79 exc6 - -17 - rif a r hi also (10 filt as r fo s weight exit - infort -signiferor: Implicate a Ta Tic.

in. b2. Japa 23. Leation dicutor considerable property the substance with fibrosis in the winder of the consideration of the consideration of the consideration.

- Fig. 51. Case 27. Section showing marked productive tissue changes with fibrosis in the wall of a terminal bronchiole. 120X.
- Fig. 52. Case 26. Section showing considerable productive tissue changes with fibrosis in the wall of a respiratory bronchiole.




Figure 51.




Figure 52.



Fig. 53. Case 21. Section showing organization of the exudate in a respiratory bronchiola.

Fig. 54. Case 21. Same. Section las above but higher magnification. 550X.




Figure 54

- Fig. 53. Case 21. Section showing organization of the exudate in a respiratory bronchiole.
- Fig. 54. Case 21. Same section as above but higher magnification. 550X.




Figure 53.




Figure 54.

Fig. 55. Case w. Section showing macrosis of the entire plane of a lobele. Note the neorotic branchicola in the center of photomicrograph. 53 august 1203.

Fig. 56. Case 51. Section showing a small absorus.
120X.

Figure 56.

- Fig. 55. Case 4. Section showing necrosis of the entire plane of a lobule. Note the necrotic bronchiole in the center of photomicrograph. 120X.
- Fig. 56. Case 31. Section showing a small abscess. 120X.




Figure 55.




Figure 56.

Fig. 57. Case 24. Section showing the border of

Fig. 58. Case 4. Section showing the patchy appearance of a recent bronchopneumonia.

- Fig. 57. Case 24. Section showing the border of an abscess just below the pleural surface.
- Fig. 58. Case 4. Section showing the patchy appearance of a recent bronchopneumonia.

 120X.




Figure 57.




Figure 58.

15. 59. Case 24. Section showing neurosis of the entire plane of a lobule. Into the demotions and thickened interleading section lock.

Fig. 60. Cese 24. Section showing a marked accumulation of leucocytes along the border of an interlobular septum. Note the two small distended lymphatics. The elemetous septum is shown in the lower portion of the picture.

- Fig. 59. Case 24. Section showing necrosis of the entire plane of a lobule. Note the edematous and thickened interlobular septum.
- Fig. 60. Case 24. Section showing a marked accummulation of leucocytes along the border of an interlobular septum. Note the two small distended lymphatics. The edematous septum is shown in the lower portion of the picture.




Figure 59.



Figure 60.

Fig. 61. Case 23. Section showing small amounts of fibrin surrounded by suboical-like cells. 180%.

Fig. 62. Case 23. Same section as above but higher magnification. 525X.

- Fig. 61. Case 23. Section showing small amounts of fibrin surrounded by cuboidal-like cells. 130X.
- Fig. 62. Case 23. Same section as above but higher magnification. 525X.




Figure 61.




Figure 62.

Fig. 60. Case 3. Leadion country 1 let v no ber coler control in the character seconds.

Fig. 54. Case 23. Section showing slysolus contential reary mout calls. 600%.

- Fig. 63. Case 3. Section showing a large number of mononuclear phagocytes in an alveolar sac. 600X.
- Fig. 64. Case 23. Section showing alveolus containing many giant cells. 600X.




Figure 63.




Figure 64.



Fig. 65. Case 31. Section showing marked productive tissue changes in the alvectar well. 140X.

Fig. 66. Case 31. Same section as above but showing an organized exudate in alveolus. 650X.




Figure 66.

- Fig. 65. Case 31. Section showing marked productive tissue changes in the alveolar wall. 140X.
- Fig. 66. Case 31. Same section as above but showing an organized exudate in alveolus. 650X.




Figure 65.




Figure 66.

Fig. 67. Jase 31. Section showing an ingresse in the number of (ingredients in floor alveolar sail. 556%.

Fig. 63. Case 25. A section showing the alveolt



- Fig. 67. Case 31. Section showing an increase in the number of fibroblasts in the alveolar wall. 550X.
- Fig. 68. Case 25. A section showing the alveoli lined with cuboidal cells. 520X.




Figure 67.




Figure 68.

-Acces 10 to the term of the address of the state of the

Pir. 70. Case Ri. Am area shorter distantion of one of the article fit of the cast set is a single straint set is a single straint.

- Fig. 69. Case 23. Section of an alveolar wall showing considerable thickening due to productive tissue changes. 650X.
- Fig. 70. Case 31. An area showing distension of one of the lymphatics of an interlobular septum with fibrin and leucocytes. 130X.




Figure 69.




Figure 70.

Nir. 71. Gase 6. A section of a bronchial lymph mode showing the hemorphage into the lymph sinuses.

Fig. 72. Case 12. Photomicrograph of a smarr made from a 24 hour broth culture of culture #17.

400£

- Fig. 71. Case 6. A section of a bronchial lymph node showing hemorrhage into the lymph sinuses.
- Fig. 72. Case 12. Photomicrograph of a smear made from a 24 hour broth culture of culture #17.




Figure 71.




Figure 72.

- Fig. 70. date 3. lear Mar restent of local lind;
 for openeds. Four Was been all of
 organisms in jairs and a or dains in the
 slyeolar emudate. 600%.
- Fig. 74. The S4. Funding section strined the salar strine of the salar strine.

- Fig. 73. Case 24. Paraffin section of lung stained for bacteria. Note the large number of organisms in pairs and short chains in the alveolar exudate. 600%.
- Fig. 74. Case 24. Paraffin section stained the same as above showing a similar condition. 600X.




Figure 73.




Figure 74.

BIBLIOGRAPHY

- Nocard, E., A New Pasteurellose: White Scours and Lung Disease of Calves in Ireland. American Veterinary Review, Vol.XXV, 1901-02.
- 2. Hagen, W. A., The Etiology and Mode of Infection in White Scours of Calves. Correll Vet., Vol.VII, October 1917, p.263.
- 3. Carpenter, C. M. and Gilman, H. L., Studies in Pneumonia in Calves. Correll Vet., Vol.XI, 1921, p.111.
- 4. Smith, Theobald, Pneumonia associated with Bacillus
 Abortus (Bang) in Fetuses of New-born Calves. Jour. of
 Exp. Med., Vol.41, May 1935.
- 5. Smith, Theobald, A Pleomorphic Bacillus from Pneumonic Lungs of Calves Simulating Actinomyces. Jour. of Exp. Med., Vol.XXVII, p.323.
- 6. Hallman, E. T., Sholl, L. B., and Delez, A. L., Observations on the Pathology of Bacterium Abortus Infections.

 Mich. Agr. Exp. Sta. Technical Bulletin No. 93.
- 7. Clark, C. F., Calf Pneumonia. Report of the Division of Veterinary Science, Michigan State College, 1936.
- 8. Glynn, J. H. The Application of the Gram Stain to Paraffin Sections. Arch. of Path., Dec. 1935, Vol.XX, No. 6, p.896.
- 9. Boyd, W. A., Text Book of Pathology. Second Edition, 1934.
- 10. MacCallum, W. G., The Pathology of the Pneumonia in the United States Army Camps During the Winter of 1917-18, Monograph of the Rockefeller Inst. for Med. Res., No. 10, April 16, 1919.
- 11. Sisson, S., The Anatomy of the Domestic Animals. Second Edition, 1930.

ACKNOWLEDGMENTS

It is with pleasure that the writer wishes to express his sincere gratitude to Dr. E. T. Hallman for the encouragement and invaluable assistance extended to this project. He wishes also to express his thanks to Dr. I. F. Huddleson for his most helpful suggestions pertaining to the bacteriological phase of this work, and to Dr. C. F. Clark for the loan of data regarding temperatures and other clinical material used in a number of instances. He is likewise indebted to Dr. L. B. Sholl for his cooperation in the collection of much of the gross material.

ROOM USE ONLY

ROOM USE ONLY