#### A GRAVITATIONAL INVESTIGATION OF NIAGARAN REEFS IN SOUTHEASTERN MICHIGAN

Thesis for the Degree of Ph. D.
MICHIGAN STATE UNIVERSITY
Gery Gordon Servos
1965





This is to certify that the

thesis entitled

A Gravitational Investigation of Niagaran Reefs in Southeastern Michigan

presented by

Gary Gordon Servos

has been accepted towards fulfillment of the requirements for

Ph.D. degree in Geology

Major professor

Date February 4, 1965

The second secon

WE 23-1000 2 2 61 WILLIAM 1985 C. 25 61

NOV 8 14326

#### ABSTRACT

# A GRAVITATIONAL INVESTIGATION OF NIAGARAN REEFS IN SOUTHEASTERN MICHIGAN

#### by Gary Gordon Servos

Precision detailed gravity surveys can detect Niagaran reefs buried at depths up to at least 3500 feet in the Michigan Basin.

Three southeastern Michigan reefs, Marine City, Belle River Mills, and Berlin, are studied by the gravity method using graphical statistical and analytical techniques of interpretation. The characteristics and sources of the anomalies are studied and optimum methods of reef anomaly enhancement are suggested.

The gravity anomalies associated with reefs in the Michigan Basın are positive with a magnitude of about 0.2 milligals. The ratio of the width of the anomalies to the width of the reef varies from 2:1 to 1.3:1 depending on the technique used.

There are negative anomalies parallel and adjacent to the reef anomalies. In areas where bedrock topography is known, these negative anomalies coincide with bedrock channels. Theoretical calculations using the configuration of the bedrock channels show that the channels can produce a gravity effect with the magnitude of the observed negative anomalies.

The sources of the reef anomalies are lateral variations in the density of the dolomite formations overlying and surrounding the reef. The reef mass alone cannot cause the anomaly.

The graphical and statistical techniques isolate the anomalies, but many of the undesirable effects of shallow source anomalies are retained by these techniques. The analytical technique of downward continuation of an upward continued surface has a high resolution and eliminates the shallow source anomalies. The maximum resolution occurs when the upward continued surface is calculated at a height equal to the depth of the reef and then downward continued to a depth not greater than the reef. The second derivative of an upward continued surface has the highest resolution using Henderson's 1960 formula. The optimum enhancement occurs when the mesh interval used in these techniques is equal to one-half the reef depth.

# A GRAVITATIONAL INVESTIGATION OF NIAGARAN REEFS IN SOUTHEASTERN MICHIGAN

Ву

Gary Gordon Servos

#### A THESIS

Submitted to
Michigan State University
in partial fulfillment of the requirements
for the degree of

DOCTOR OF PHILOSOPHY

Department of Geology

1965

933259

COFYRIGHT

Br

GARY GORDON SERVOS

1965

#### ACKNOWLEDGMENTS

The author wishes to express his sincere thanks to the following individuals and organizations.

To Dr. W. J. Hinze for his sincere interest, patience, guidance, and helpful criticism during the study and preparation of this manuscript.

To Dr. C. E. Prouty, Dr. H. B. Stonehouse, and Dr. J. W. Trow for reviewing the manuscript and for their helpful suggestions.

To the McClure Oil Company, Alma, Michigan, particularly Mr. Harold McClure, for financial assistance in the gathering of the field data.

To Mr. D. W. Merritt for his assistance in preparing the data for the computer and for the use of several of his programs in interpretating the data.

To Messers R. Kellogg, W. Keith, J. Bradley, R. Ehlinger, R. Shaw, D. Hill, J. Norris, R. Cranson, and D. Walker whose hard work made the collection of the data possible.

To Michigan State University for free use of the C.D.C. 3600 computer.

## TABLE OF CONTENTS

																Page
ACKNO	DWL	EDGMENTS	•	•				٠	٠	¢	3	3	•	•	•	11
LIST	OF	TABLES		6	a '	3	¢	J	•		c	•	•	•	•	v
LIST	OF	FIGURES	٥	•	•	v	•	•	ć	c	٥	•	•	•	•	vi
Chapt	ter															
	I.	INTRODU	CT:	ION	٠	5	5	·	•	J	¢	•	•	•	•	1
		General Locatio							, of			S.	•	•	•	1
]	ΙΙ.	GRAVITY	. Si	<b>TU</b> D	Υ.	•				ç		•	•	•	•	6
		Field M Reducti				a	•		ن ټ	e e	•		•	•	•	6
I	ΙΙ.	GEOLOGY					٥	•	•	6	•	•	•	•	•	14
		Regiona Mich					e and	i St		ign	_	hy •	of •	3		14
]	IV.	INTERPF	RET	ATI	ONAI		IETHO	DS	o	•	•	•	c	۰	•	22
		General	. S1	tat	emer	nt		•	•	•	•	•	•	•	•	22
	٧.	GRAVITY	' Al	MOV.	ALIE	ES	FROM	I RE	EEFS	5.	•	•	•	•	•	42
. 1	/I.	INTERPF	RETA	ATI	ON N	ΙΞΊ	HOD					•	•	•	•	47
		Regiona Marine Belle F Berlin	Cit live	ty :	Reei Mill	f S ls	Study Reef	, St	udy	•	•	•	•	•	•	47 47 79 97
V	ΙΙ.	SUMMARY		c		•	ž	o	•		•	•	•	•	•	111
		Detecti Charact												•	•	111

																Page
	Sour Reco	ces mme	of nda	Re tio	ef ns	Anor	nal Fu	ies rth	er	Stu	idy	,		•		112 114
BIBLIOGRA																
APPENDIX.	•	v	•		•		•	c	•		•	•	•	•	•	120

### LIST OF TABLES

Table		Page
I.	Accuracy of Gravity Data	13
II.	Reef Gravity, Average Conditions and Results in Three Regions	43

### LIST OF FIGURES

Figure		Page
1.	Location of Areas	5
2.	General Stratigraphy of the Michigan Basin .	15
3.	Location of Reefs in the Michigan Basin	17
4.	General Stratigraphy of Niagaran and Salina Groups	18
5,	Structure of a Typical Reef	18
6.	Bedrock Topography of Marine City and Belle River Mills Areas	20
7.	Bedrock Lithology of Marine City and Belle River Mills Areas	21
7a.	Flow Chart of Interpretation Procedures	25
8.	Regional Gravity from Bouguer Anomaly Profile	24
9.	Template for Analytical Techniques	30
10.	Geometrical Elements for Talwani's Two-Dimensional Calculations.	35
11.	Geometrical Elements for Talwani's Three- Dimensional Calculations	38
12.	Portion of Regional Gravity Map of Michigan .	46
13.	Marine City Bouguer Gravity Anomaly Map	49
14.	Typical Bedrock Channel and Gravity Anomaly .	51
15.	Cross-Section of Units Used in Three-Dimensional Gravity Calculation and Resulting Gravity Anomaly	53
16.	Mg/Ca Density Ratios and Theoretical Gravity Anomaly.	55

Figure		Page
17.	Marine City Cross-Profile Residual Map	65
18.	Marine City Least Squares Residual 0-3rd Degree, 500 foot Station Interval	66
19.	Marine City Least Squares Residual 0-5th Degree, 500 foot Station Interval	67
20.	Marine City Least Squares Residual 0-3rd Degree, 1000 foot Station Interval	68
21.	Marine City Least Squares Residual 0-5th Degree, 1000 foot Station Interval	69
22.	Marine City Interpolated Bouguer Gravity Map.	70
23.	Marine City Bouguer Gravity Anomaly Upward Continued 2500'	71
24.	Boundaries of Analytical Techniques	72
25.	Marine City Bouguer Gravity Anomaly Upward Continued 2500' and Downward Continued 5000'	73
26.	Marine City Bouguer Gravity Anomaly Upward Continued 3000' and Downward Continued 5000'	74
27.	Marine City Bouguer Gravity Anomaly Upward Continued 5000' and Downward Continued 5000'.	75
28.	Marine City Second Derivative of 2500' of Upward Continued Surface	76
29.	Marine City Second Derivative of 3000' of Upward Continued Surface	77
30.	Marine City Second Derivative of 5000' of Upward Continued Surface	78
31.	Belle River Mills Bouguer Gravity Anomaly	85
32.	Belle River Mills Cross-Profile Residual	86
33.	Belle River Mills Least Squares Residual 0-3rd Degree, 500 foot Station Interval .	87

Figure		Page
34.	Belle River Mills Least Square Residual 0-5th Degree, 500 foot Station Interval .	88
35.	Belle River Mills Least Square Residual 0-3rd Degree, 1000 foot Station Interval.	89
36.	Belle River Mills Least Square Residual 0-5th Degree, 1000 foot Station Interval.	90
37.	Belle River Mills Interpolated Bouguer Gravity Anomaly	91
38.	Belle River Mills Bouguer Gravity Anomaly Upward Continued 2500'	92
39•	Belle River Mills Bouguer Gravity Anomaly Upward Continued 5000' and Downward Continued 5000'	93
40.	Belle River Mills Second Derivative of 2500' Upward Continued Surface	94
41.	Belle River Mills Second Derivative of 3000' Upward Continued Surface	95
42.	Belle River Mills Second Derivative of 5000' Upward Continued Surface	96
43.	Berlin Bouguer Gravity Anomaly	101
44.	Berlin Least Square Residual 0-3rd Degree, 500 foot Station Interval	102
45.	Berlin Least Squares Residual 0-5th Degree 500 foot Station Interval	103
46.	Berlin Least Squares Residual 0-3rd Degree, 1000 foot Station Interval	104
47.	Berlin Least Squares Residual 0-5th Degree, 1000 foot Station Interval	105
48.	Berlin Interpolated Bouguer Gravity Anomaly .	106
49.	Berlin Bouguer Gravity Anomaly Upward Continued 2500'	107
50.	Berlin Second Derivative of 2500' Upward Continued Surface	108

Figure						Page
51.	Berlin Second Continued	Derivative Surface .	_	•	•	109
52.		Derivative Surface	-	•	•	110

#### CHAPTER I

#### INTRODUCTION

#### General Statement

Exploration for oil and gas bearing Niagaran reefs has been emphasized in the Michigan Basin since 1960. This has resulted in the discovery of several reefs in southeastern Michigan in St. Clair and Macomb Counties. In addition, isolated reefs have been found at several locations around the basin and, thus, the prospect of locating additional reefs is excellent.

Exploration for the reefs has included both geological and geophysical techniques. Seismic investigations have had limited application, but the gravity method has been widely employed and is credited with making several major discoveries. Despite the wide use of gravity no detailed reports of case histories have been presented for reefs in the Michigan Basin.

The purpose of this investigation is to study the gravity anomalies of known reefs in the Michigan Basin and to establish the optimum methods of interpretation and isolation of the anomalies.

Although the results of gravity prospecting in the Michigan Basin have not been reported, several investigators

have discussed gravity prospecting for reefs. Yungul (1961) has produced the most comprehensive report on reef anomalies in several areas including the case histories of a reef in Ontario (Pohly, 1954). Geyer (1962), using theoretical approximations, concluded that reefs in the Michigan Basin could be found by gravity.

Three well defined reefs of different relief, areal extent, and depth of burial were selected for study. A precision detailed gravity survey was conducted over the three reefs during the summers of 1962 and 1963. A high station density was used to obtain all the details of the gravity anomalies. The three reefs chosen are Belle River Mills, Marine City, and Berlin.

The Belle River Mills reef is one of the larger reefs discovered. It is approximately 3000 feet wide and two miles in length. It is buried at a depth of 2300 feet below the surface and has a relief of 300 feet.

The Marine City reef is three miles south of the Belle River Mills reef and is buried at the same depth. It consists of two circular cone shaped pinnacles on a broad base. The pinnacles are one mile apart and have a relief of 250 and 300 feet. The Marine City reef contains less reef mass than the Belle River Mills reef.

The Berlin reef is located thirty miles west of the Marine City and Belle River Mills reefs and is buried at a

depth of 3500 feet. It has a width of 2500 feet and a length of one mile. The relief is about 200 feet.

Several graphical, statistical, and analytical techniques were used in the interpretation and isolation of the reefs. The techniques used include cross-profiles, least squares polynomial analysis, upward continuation, downward continuation and second derivatives. Two-dimensional and three-dimensional gravity formulas were used to calculate the theoretical gravity of the reefs. All computations were performed on a digital computer.

#### Location and Physiography of Areas


The areas of study, shown in Figure 1, are located in St. Clair and Macomb Counties, Michigan. The Marine City and Belle River Mills areas are in the extreme eastern part of St. Clair County along the St. Clair River. This area lies between latitude 42°40'00" N and 42°48'30" N and longitudes 82°30'00" W and 82°35'00" W. The Berlin area is located along the northern Macomb and southern St. Clair County line. This area lies between latitudes 42°52'00" N and 42°56'30" N and longitudes 82"53'45" W and 82°58'45" W.

Both the Marine City and Belle River Mills areas are situated in the Erie-Huron Lowland, a topographic subprovince in the Southern Peninsula of Michigan, (Newcombe, 1933). This sub-province is characterized by surficial glacial lake clays and lake bed sands. This area was

covered by several glacial lake substages of the Wisconsin glacial stage (Leverett, 1939). This area is generally flat with a maximum relief of fifty-five feet and the elevation gradients of five to ten feet per mile. The only exceptions are the several stream channels in the area which have a maximum relief of thirty feet.

The Berlin area is situated in the Thumb Upland, a topographic sub-province characterized by several moraines (Newcomb, 1933). The immediate area of the Berlin survey is flat with a maximum elevation difference of about forty-five feet. The elevation gradients also are five to ten feet per mile. Both areas are divided by a network of section line and intermediate roads. Generally the area consists of privately owned farms.

Primary drainage in the Marine City-Belle River Mills area is to the south and east. The major tributary is the Belle River. The Berlin area exhibits little or no drainage pattern and there are no major streams in the area.



LOCATION OF AREAS FIGURE 1

#### CHAPTER II

#### GRAVITY STUDY

#### Field Methods

The areal extent of the gravity surveys was selected to cover the reefs adequately and include the regional gravity of each area. The existing road network allowed good coverage, but in addition to the road traverses, cross-country traverses were placed in the immediate vicinity of the reefs at one-half mile intervals. These cross-country traverses were made in the areas whenever possible and in all cases at least one cross-country traverse was placed in the sections associated with the reefs.

A station spacing of five hundred feet was selected for the survey. The station elevations were established with a Zeiss self-leveling level and elevation control was obtained from three U. S. Geological Survey and several Michigan State Highway Department bench marks. All traverses were closed and cross-country lines were tied into established elevations. Horizontal distances were determined by stadia intervals and control was carried from road intersection to road intersection to eliminate cummulative error.

Two exploration type gravity meters were used in the survey. These were World Wide meter number 45 and Worden number 99. These meters had calibration constants of 0.10093 and 0.0995(1) mg. per scale division respectively. These constants were checked prior to the beginning of the survey each summer and midway through the summer. Both meters measure the relative acceleration of gravity to an accuracy of 0.01 mg.

The meters are subject to time variations or drift even though they are temperature compensated and barometrically controlled. These time variations are a result of temperature changes, tidal variations, spring fatigue, and handling of the meter. To determine these variations a check reading was made at a number of pre-established base stations at least once every hour during the course of the survey.

A primary base was established near the center of each area and sub-bases were established at strategic points by the base-looping method. The Belle River Mills and the Marine City areas are adjacent and the primary bases of each area were tied to allow correlation. The Berlin area was not tied to the other two surveys because it is thirty miles away.

Gravity readings were made every five hundred feet at the pre-established elevation. These readings were tied to the base station by the hourly drift check. In addition, a number of stations were reread at various times during the course of the survey to establish the accuracy of the readings. At each gravity station the meter was read until two values were obtained which did not differ by more than 0.2 scale divisions or 0.02 mg.

#### Reduction of Data

#### Introduction

Observed gravity, to be useful in geological studies must be corrected for station elevation, latitude, and terrain. These corrections are applied to each station.

The resulting values are called the Bouguer gravity anomaly.

The complete Bouguer gravity anomaly was calculated on a digital computer according to the following formula:

$$G_{cbg} = g_o - g_l + g_{e\&m} + g_t$$
 (1)

where

G<sub>cbg</sub> = complete Bouguer gravity anomaly

g = observed gravity

 $g_{\ell}$  = latitude correction

 $g_{e\&m}$  = combined elevation and mass correction

g<sub>+</sub> = terrain correction.

### Observed Gravity

The observed gravity values are obtained by correcting the meter readings in scale divisions for drift, and then multiplying the values by the calibration constant of the meter. The drift is determined from graphs of hourly

base-readings. When the drift exceeded one scale division per hour, the drift check readings were decreased to three-quarters of an hour.

#### Latitude Correction

The Earth's gravity increases from the equator to the poles. The latitude correction takes into account this change in gravity. Latitude corrections are made from a base latitude by multiplying the latitude distance of each station by a constant, K. Nettleton (1940) determined K to be 1.307 Sin20 mg. per mile, where 0 is the mean latitude of the survey area. In the survey areas of this study a base latitude of 42°40'00" N was used for the Marine City-Belle River Mills area, and a base latitude of 42°52'00" N for the Berlin area. The value of K is 0.0002474 mg. per foot for these areas.

The stations of the three surveys were placed on the Michigan Coordinate System, east zone, a plane-coordinate system established by the U. S. Coast and Geodetic Survey. This allows each station to have x (ordinate) and y (abcissa) coordinates which are on a rectangular grid system. In the process of reducing the data by the digital computer, the Michigan Coordinate System was rotated to correspond with latitude and the latitude correction was made in the normal manner.

#### Elevation and Mass Correction

The elevation or free-air correction and the mass or Bouguer correction are often combined and called the elevation and mass correction. The combined correction takes into account the vertical decrease of gravity with an increase in elevation and the increase in gravity due to the attraction of the material between the datum and individual stations. This correction is calculated from the combined formula:

$$g_{e\&m} = (0.09406 - 0.01276\rho)h$$
 (2) where

- h = elevation difference between the datum and
  the gravity station in feet
- ρ = density of surface material in g/cc.

The datum to which the gravity was reduced was chosen near the lowest elevation of the surveys.

#### Density Determination

The determination of the correct density for the near surface material in the survey areas is extremely important. A density value for glacial drift is often given in the literature as 1.8 g/cc. This value does not appear to be correct for near surface glacial drift in eastern Michigan. A density value of 2.1 g/cc was determined by the Nettleton density profile method at several locations in the vicinity of the survey areas. Klasner (1964)

determined a density of 2.15 g/cc for the near surface drift in western Michigan.

An incorrect density of 0.3 g/cc and relief of thirty feet will cause the gravity values to be in error by about 0.12 mg. In the search for reefs this value alone is close to three times the desired accuracy. In areas with no topography, a larger error in density can be tolerated. However, in areas with relief it is very important to determine the desnity with the highest precision possible.

#### Terrain Correction

The relief in the survey areas was low enough that no terrain corrections were needed. Where the local variations of the several stream valleys in the areas were encountered, stations were placed far enough away from the feature that the effect was negligible or the stations were not used. The placing of stations in these areas was determined from correction curves calculated by Hubbert (1948).

#### Summary of Accuracy

In making gravity reductions, there are several factors which cause errors in the Bouguer gravity anomaly. These are:

- 1. Errors in observed gravity
- 2. Errors in elevation
- 3. Errors in latitude
- 4. Errors in assumed density of the glacial drift.

To check the accuracy of the gravity readings for the Belle River Mills survey, 16 stations were reread during the survey. The standard deviation for the repeated stations is ± 0.01 mg.

The errors in elevation affect the elevation and mass correction. Thirty-six closures were made during the survey. The standard deviation of the closures is  $\pm 0.09$  feet. The maximum possible error in the elevation and mass correction caused by 0.09 foot error, using a density of 2.1 g/cc, is 0.006 mg.

The latitude measurements were made to an accuracy of fifty feet. They were measured on a base map with a scale of 1:12000. The maximum error from the latitude correction is 0.012 mg.

An incorrect density value of 0.1 g/cc with a maximum change in elevation between any two stations of 10 feet will cause an error of 0.01 mg. The formula for calculating this error is:

Error =  $0.00128\sigma h$ 

where

 $\sigma$  = error in density in units of 0.1 g/cc

h = maximum relief in feet.

The maximum error possible considering these sources is  $\pm 0.038$  mg.

Table 1 shows a comparison of the maximum errors for the Belle River Mills, Marine City, and Berlin surveys.

The errors for the Marine City and Berlin surveys were calculated in the same manner as those of the Belle River Mills survey.

TABLE 1
ACCURACY OF GRAVITY DATA

Number of Gravity Stations Reread	Standard Deviation of Reread Stations in mg.	Number of Elevation Closures	Standard Deviation of Closures in Feet	Maximum e&m Error Us Using Standard Devia- tion of Closures and Density of 2.1 g/cc.	Maximum Latitude Error in mg. Stations Accu- rate to 50 feet.	Maximum Error for 0.1 g/cc Density Error and Elevations Difference of 10 ft. between two stations.	Maximum Error Possible in mg.
16	0.1	36	0.09	0.006	0.012	0.01 ±0	0.038
27	0.13	30	0.06	0.004	0.012	0.01 ±0	0.041
30	0.14	8	0.11	0.007	0.012	0.01 ±0	0.043

#### CHAPTER III

#### GEOLOGY

# Regional Structure and Stratigraphy of Michigan Basin

The Michigan Basin is an oval intracratonic sedimentary basin enclosed by the Findlay, Algonquin, Kankakee and Wisconsin Arches, and the Canadian Shield. The development of the basin probably began in middle Silurian time (Ehlers and Kesling, 1962). During the middle Silurian (Niagaran Series) a system of reefs was formed on shelf areas surrounding the Michigan Basin. Slight rises in the areas bordering the Michigan Basin, accompanied by a barrier of Niagaran reefs, caused a restriction of the basin late in Silurian time. The Michigan Basin continued to receive sediments throughout the Paleozoic period.

The general stratigraphy of the basin is shown in Figure 2 (after the Michigan Geological Survey). The Niagaran and lower Cayugan Series of the Silurian, which are of major interest in this study, consists of the Niagaran Group (or Guelph) and the Salina Group respectively.

Lowenstam (1950) suggested that Niagaran reefs
possibly surround the entire Michigan Basin. Isolated reefs
which have been found around the margins of the basin

#### STRATIGRAPHIC SUCCESSION IN MICHIGAN

E		+	S	YSTEM	SERIES	ST	AGE			
	¥				RECENT	743	or Sade Conta Insurada	+	MICHIGAN DEPARTME	NT OF
	CENOZOIC	Q	u	ATERNARY	PLEISTOCENE	Glacaton Care	late Stade (R. Hu Stade	-1	GROUNGS Constit 1 (60)	
i	Ľ					Sangaron transface	vet trade			
•	,	_		0.0	CDOD 11011					
7	OU	061	1	TIME STRATIGRAL	CROP NOM	ROCK STRATIGRA	PHIC	7 -	SUBSURFACE N	ATK BA
_	T								ORMATION MEA	
FRA	Digital	TO TO TO		SERIES	GROUI	FORMATI	ON MEMBI	R		nes e i
	-	+			-		_		CONTRACTOR NO NO.	
č	2	2	5	ď				and on	-	
MESOZOIC	PACCIC	I VI	THE PARTY	KIMERIOGA	N			1	Ted Sed	
MES	ı		1	2				Subseries only?	TON	
-		÷	2	2	-	-	-	-		-
	PENNSYLVANIAN	Y	VANIAN	COMMANG	н	Crand Ever In	to the beautiful to the	7	Gold Row Fo.	
	YLVZ		AL IV		-		he dispus	-60 h		
	NNS	EABLY	DEMMENT	POTTSYSLE		Sagnew Ass	fee is		Suprav En Sur	
	N D	-			-	Bergot Li	-		Service Control	
	MISSISSIPPIAN	Z	MISSISSIDDIAN	MERAMICIAN	GANG M	Di Markeyan Am			Report to exact	
				OSASIAN		Marshall St	-		Total Island	
	MISS	EARLY	MISS	KNOERHOOKS		Colorador SA		Of the same	Caldways St	
	M	55	15	SIPPIAN-DEV	ONIAN U	red Charach St.		and the last of th	Marine M.	
		N.		CHAUTAUQUA	N	Annua M			haber in	Zana
		H		SINICAN	-	Squire Bay Lo Funder Bay Lo Futher Farm An Number From Fro Name Wille Dam Fro Name Wille Dam Fro	_	San Asian Asia	loans for in ph	
					Navey	Alpena Li	-		Append to 1	
	N.		Z	MAIR		Newton Coast La	Man	2/3/2/- +	and the state of t	
	DEVONIAN	MDDL	DEVONIAN			Parish Pair Fe.				
	ŝ	٤	Dey Dey			Gerakeu Fo Feron Foet for Backgor Charry I Bell Sh Rogers Cay to	-	THE RESERVE OF THE PARTY OF THE		
						Dunder to Anderdon fin			Section 1.	
				ULSTERIAN	DENOT RIVER	Locar An				T
				OLIVERIAN.		Anterotory for Software So Box Block for		-	refer the 2 for 2 or	9
					-	E box New York	-	A		_
ļ	1	LARLY	4			Corden Manual Sen			edge bland for	
					BASS ISLANDS	Resenting Dal Arrester Dal		of.		6
		š		CAYUGAN		St Specia Del nuo			-	
					SAUNA	Pe air Cheve St.				
		4	ŀ							5
14.4		100							2 Carbonal da	
CHIDIAN	1	CHILDIAN	2			Engadese Del		23	2 Evapores C	
				MAGARAN	MANSTOLE	Cordel Dut			3.4	+
						Cordel Dur Scheidright Dui Mandricks Dol Byron Dur	Short to	Charles of the		140
			ŀ		KANC NUI			The state of the s	Community Community	-
	APPLA	į.		ALEXANDRIAN	CATABACT	Cabot Head Sh.		-	der Head Sk. 1	CA
		Ť	t			Mannoule Sel By Hill Fe			mende Out 27	-
	ATS			CINCINNATIAN	ROWOND	Statemates Fin.	Spring Stor As No.		100 CH	
z						Mr Creek Sh			The same of the sa	and a
ğ		ORDOVICIAN	t			Green Quarry Re.	Official	Maria de la compación de la co	Colonia Colonia	+
DKDOVICIAN	MODEL	200		MOHAWKIAN	HINTON	Chandler falls file				100
5	Ľ	ő	L		BACK BYER	Bory falls for			Consession of	0 minut 1
	FARIT		-	CHAZYAN		-	1		Part St.	1
	Š			CANADIAN	PEARS DU ONN	Oneou Out	77		August Dat 155 • Referend So	PRI
ĺ	ľ		ľ		-	Transposition Fm.	noder to	Contract of the Contract of th	note by 2215	+
,	Z	,		ST CROSSAN	LAST SUPERIOR		It is to the book in		Side to a los	
		RIA				Municip Pa	Chape Said In		and the desired to the control of th	Ψ"
Compation	OLE	CAMBRIAN					-	~	-	-
1	EASEY MIDDLE	V				Jacobovilla Sa		V	derela la	
	ASE					(NA)		AMBRIAN ,	100 mm	


substantiate this (Figure 3). Reefs also are present in the Silurian outcrops on Drummond Island, Michigan (Manley, 1964). Generally the reefs developed around the basin are of the pinnacle type.

# Niagaran Structure and Stratigraphy of St. Clair County

The regional dip of the geological formations in this area is in the order of 30-40 feet per mile. This becomes an important factor in searching for reefs in the Michigan Basin as the gravity method is useless if the reef is below a limiting depth.

The general stratigraphy of the Niagaran and Salina groups in the St. Clair area is shown in Figure 4 (Alguire, 1962). There is a possibility that the carbonate sequences of the Niagaran and Salina Groups are regionally limestone and only locally dolomitized about the reefs (Jodry, 1965).

Typical reefs in the St. Clair County area have relief up to 300 feet with an areal extent usually less than one half mile in width and one mile in length. Basinward the reefs are more pinnacle in nature and have a smaller areal extent. The structure of a typical reef is shown by a cross-section through the Marine City reef (Figure 5, Alguire, 1962). This cross-section includes the stratigraphic units from the Clinton shale (Niagaran) through the B-Salt member of the Salina group. The entire geologic column from the reef facies upward show some degree of doming over the reef



LOCATION OF REEFS IN MICHIGAN FIGURE 3


# NIAGARAN AND SALINA STRATIGRAPHY

	T								
Formation	Unit	Description							
	G	DOLOMITE: brown, finely crystalline; shaly dolomite; some anhydrite.							
	F	SALT: in thick beds separated by beds of shale, shaly dolomite, gray and buff, and brown, crystalline dolomite; anhydrite nearly always present.							
	E	SHALE: with argillaceous, gray and buff dolomite.							
<b>₹</b>	D	SALT: nearly pure; thin partings of buff dolomite.							
SALINA	С	SHALE: gray, dolomitic.							
7S	В	SALT: thick salt beds with thin dolomite layers.							
	A-2	DOLOMITE: brown, brown gray, gray and dark gray, finely crystalline; some dark bituminous shale.							
		SALT: where salt is absent the base of A-2 is marked by anhydrite.							
	A-1	DOLOMITE: buff, brown, brown gray and dark gray, dense to medium crystalline; some dark bituminous shale.							
		ANHYDRITE: at base.							
1 ~ CL I	Guelph- ockport	DOLOMITE: tan, gray brown and brown, very finely to coarsely crystalline and vugular; often finely laminated near top.							
A G G R C	or iagaran	DOLOMITE: light and dark gray mottled, finely crystalline.							
IZ		DOLOMITE: light to blue gray, finely to coarsely crystalline.							

### AFTER ALGUIRE

# FIGURE 4

STRUCTURE AND STRATIGRAPHY OF MIDDLE SILURIAN NIAGARAN AND UPPER SILURIAN SALINA STRATA ACROSS A PART OF THE MARINE CITY REEF FIELD, ST. CLAIR COUNTY, MICHIGAN. VERTICAL SCALE 6.6x HORIZONTAL SCALE




(Ells, 1964). In most cases this doming of the sediments can be accounted for by compaction.

Details of the structural and stratigraphic development of reefs have been reported by many authors (Cummings and Shrock, 1928), (Lowenstam, 1950, 1957), (Textoris and Carozzi, 1964).


#### Bedrock Geology

The glacial drift deposits covering the St. Clair
County area consist primarily of lake bed clays and sands.
Isopach maps of the glacial drift (Brown, 1963) show variations in the drift thickness in the order of 200 feet.
Structure contours of the bedrock surface beneath the drift show numerous bedrock channels (Figure 6, Brown, 1963).
Klasner (1964) showed that gravity can be used successfully to determine the location of these bedrock channels. Bedrock channels with depths of 200 feet can cause negative gravity anomalies which will tend to mask the anomalies of the reefs.

According to Brown (1963) the bedrock lithology varies from older to younger beds to the west as illustrated in Figure 7. These changes in bedrock lithology are important in the theoretical gravity to determine the bedrock topographic effects, as variations in lithology change the density contrasts between the bedrock and glacial drift.



BEDROCK TOPOGRAPHY OF MARINE CITY AND BELLE RIVER MILLS AREAS C.I.=20' FIGURE 6



BEDROCK LITHOLOGY OF MARINE CITY AND BELLE RIVER MILLS AREAS FIGURE 7

### CHAPTER IV

### INTERPRETATIONAL METHODS

### General Statement

gravity measurementshave been used for a number of years as an exploration tool, but the confidence placed in the method has been limited. However, in the last decade confidence in the gravity method has increased markedly for several reasons. First, a better knowledge of geology by geophysicists and a better understanding of the application of geophysics by the geologist has contributed much to the change in attitude. Second, previous studies have defined the usefulness and validity of the gravity method for various geologic situations. Third, a better knowledge of the assumptions and variables used in gravity interpretation has been gained and applied in evaluating gravity data.

New and useful techniques such as density logging are and will be a great help in future gravity interpretations. The determination of the correct density contrast between geological formations is often difficult, but is extremely important in gravity interpretation. The gravity method depends on horizontal changes in density and, fortunately, geological conditions produce these density contrasts.

An inherent problem in gravity interpretation is the ambiguity of results. Skeels (1950) has shown that any gravity anormaly produced by a mass (x) can be reproduced by another mass (y) at a different depth. Studies in the last decade have helped to overcome the ambiguity by defining the types of anomalies produced by various geological features in different geological situations.

Regional gravity surveys are widely used, but detailed surveys have not been fully utilized in the search for small geologic features. Noise levels are constantly being lowered as detailed gravity survey are used more frequently. When the data is collected in the most accurate form and proper assumptions made in the interpretation, the detailed survey can be used with confidence.

Digital computers now allow many new and older interpretational techniques to be used which have been impractical due to involved and time consuming arithmetic. There has been much discussion and disagreement on graphical versus analytical and statistical techniques in interpretation. The graphical techniques are simple, but are subject to personal bias of the interpreter while the analytical techniques have a higher resolving power and are readily performed on the digital computer. The statistical methods also are easily adapted to the computer and are subject to less bias than the graphical methods. Each method has its use in gravity interpretation given the proper set of geological and geophysical conditions.

## Theory of Techniques Used for Interpretation

### General Statement

Graphical, statistical, and analytical techniques were used in the interpretation of the reef anomalies as one of the purposes of this study is to determine optimum methods of interpretation for a particular geologic situation. A flow diagram, shown in Figure 7a, illustrates the step by step procedure followed in the interpretation and the results of each particular method. The theory of these techniques is presented in the following sections.

### Cross-Profiles Graphical Technique

The cross-profile method was used to determine the relationship of the residual anomaly obtained using a graphical technique to those obtained using analytical and statistical techniques. This method consists of drawing a smooth curve of the estimated regional gravity gradient along a profile as shown in Figure 8.

### CROSS PROFILE METHOD





FIGURE 8



FLOW CHART OF INTERPRETATION PROCEDURES

This is done for all profiles of gravity values which intersect another profile. If the regional gradient selected is a good approximation, the point of intersection on two profiles will agree in value. If the values do not agree, then the approximated regional greadient is adjusted until the values at the intersection points agree and are geologically reasonable. When all values of intersection agree, it is assumed the regional gradient has been correctly determined. The approximated regional gravity is then removed from the Bouguer gravity values leaving the residual gravity.

This method has the advantage of being a simple method to obtain the residual and permits the knowledge of an area to be used. The results from this method are, however, subject to the personal bias of the interpreter which can be both an advantage and a disadvantage. The method is limited if the residual anomalies are low in magnitude or the regional gradients are complex.

# Least Squares Polynomial Statistical Analysis

This method consists of fitting a polynomial equation to a three-dimensional surface. In this case the surface consists of x, y coordinate points and a gravity value at each point giving the third dimension in the z direction. The polynomial equation is fitted to the surface to a degree where the regional gravity is adequately defined.

The portion of the surface which is not fitted exactly by the equation is considered the residual. The success of this approach depends on the validity of the assumption that the regional can be described accurately by a polynomial equation.

The least squares principle states that the coefficients of the polynomial equation must be such to make  $\sum_{\bf i} (\epsilon_{\bf i})^2$  a minimum, where  $\epsilon$  is the residual or the portion of the surface not fitted exactly by the polynomial equation.

The basic polynomial equation used is

 $\Delta \emptyset_{\text{Regional}} = \alpha_{00} + \alpha_{10} x + \alpha_{10} y + \cdots + \alpha_{pq} x^p y^q$  (4) where the  $\alpha$ 's are the coefficients of the equation and (p, q) is the degree of the equation. The equation can be solved easily by matrices on a digital computer and the resulting gravity value expression is subtracted from the Bouguer gravity to give the residual.

### Analytical Techniques

Upward and Downward Continuation

Often in the interpretation of gravity data it is desired to study the gravitational field on a plane above or below the original plane of observation. The upward continuation method results in moving the plane of observation further away from the source of anomalies. The effect of this method is to eliminate small sharp gradient anomalies from sources close to the plane of observation

further away from the source of anomalies. The effect of this method is to eliminate small sharp gradient anomalies from sources close to the plane of observation and to retain the broad lower gradient anomalies caused by masses at a greater depth. The downward continuation method allows the plane of observation to be moved to levels near the masses producing the anomalies. This results in an enhancement and increased resolution of the anomaly. Both upward and downward continuation have proven highly useful in gravity interpretation.

The following mathematical evaluation based on Henderson's (1960) development results in a practical approximation for both the upward continuation and downward continuation equations.

Basicaly, the problem is to compute the gravity value  $\Delta \emptyset(x, y, z)$  above and below the plane of observation.

The origin of the right-handed coordinate system is taken at the point where the field is to be computed, with the z-axis positive downward.

The integral solving the Dirichlet problem for a plane (the upward continuation integral) in polar coordinates is

$$\Delta \phi \ (-ma) = \int_{0}^{\infty} \frac{ma \, \Delta \phi(r) \, r \, dr}{(r^{2} + m^{2}q^{2})^{3/2}}$$

$$m = 1, 2, 3, \dots, n$$
(5)

where a is the mesh interval of the gravity values, m is the number of mesh units above the original plane, and

$$\Delta \overline{\mathbf{d}}(\mathbf{r}) = \frac{1}{2\pi} \qquad \Delta \phi(\mathbf{r}, \theta) d\theta. \tag{6}$$

The physical meaning of  $\Delta\phi$  on (r) is the average value of  $\Delta\phi$  circles of radius r about a point. Experimentally it has been found that radii of r=0, a,  $a\sqrt{2}$ ,  $a\sqrt{5}$ ,  $a\sqrt{8}$ ,  $a\sqrt{13}$ ,  $a\sqrt{25}$ ,  $a\sqrt{50}$ ,  $a\sqrt{136}$ ,  $a\sqrt{274}$  and  $a\sqrt{625}$  adequately sample the field. The number of mesh points falling on the circles having these radii are respectively 1, 4, 4, 8, 4, 8, 12, 12, 8, 8, and 12 as shown in Figure 9.


Next a Lagrange interpolation polynomial is fitted to  $\Delta \phi$  (0) and the n values  $\Delta \phi$  (-ma) computed from equation 5 to obtain the approximation formula,

$$\Delta \phi(\mathbf{s}) \approx \sum_{m=0}^{n} \frac{(-1)^{n} \mathbf{s} (\mathbf{s}+\mathbf{a})(\mathbf{s}+2\mathbf{a}) \cdots (\mathbf{s}+n\mathbf{a})}{\mathbf{a}^{n} (\mathbf{s}+m\mathbf{a}) (n-m)! m!} \Delta \phi(-m\mathbf{a}). \tag{7}$$

From experimental work it has been found that the approximations are good for n<5.

The accuracy of equation 7 depends on the accuracy of  $\Delta\phi(-ma)$  as computed from the upward continuation integral, equation 5. A numerical integration employing a mean value theorem over each integral  $r_1 \stackrel{\leq}{=} r_{1+1}^{\leq}$  is carried out. The integral in equation 5 is replaced by the sum

$$\Delta \phi(-ma) = \sum_{i=0}^{n=1} \left[ (r_{i+1} - r_i)^{-1} \int_{r_i}^{r_{i+1}} \Delta \phi(r) dr \right] ma \left\{ \left[ r_i^2 + (ma)^2 \right]^{-\frac{1}{2}} - \left[ r_{i+1}^2 + (ma)^2 \right]^{-\frac{1}{2}} \right\} + O\left( \frac{1}{r_n} \right)$$
(8)



TEMPLATE FOR ANALYTICAL METHODS
FIGURE 9

By setting a equal to unity and successively evaluating equation 8 for m = 1, 2, 3, 4, 5, the five sets of coeficients adopted for upward continuation are obtained. The working equation is given as

$$\Delta\phi(-\text{ma}) \approx \sum_{i=0}^{10} \Delta\overline{\phi} (r_i) \text{ K } (r_i,m)$$
 (9)

where K  $(\mathbf{r_i}, m)$  is the set of coefficients which operate on the center value and the ten ring-averages values,  $\Delta \overline{\phi}(\mathbf{r_i})$ , to give  $\Delta \phi$  at ma mesh units above the original plane.

For downward continuation, z is set equal to a, 2a, 3a, 4a, 5a, in equation 7. This results in the calculation of the gravity on these levels below z=0, using the same ten ring averages,  $\Delta \overline{\phi}$   $(r_1)$ . The working equation is given as

$$\Delta \phi(\mathbf{k}) \approx \sum_{i=0}^{10} \Delta \phi(\mathbf{r}_i) D(\mathbf{r}_i, \mathbf{k})$$
 (10)

where D  $(r_1, k)$  is the set of coefficients for computing  $\Delta \phi$  K mesh units below the original plane.

### Second Derivative

The second derivative method has proved useful for gravity interpretation. Its importance arises from the fact that the second derivative analysis tends to emphasize the smaller, shallower geologic features at the expense of larger, regional features. The subject of second derivatives

is treated by many authors, including Peters, (1949); Elkins (1951); Henderson and Zietz (1949); Rosenbach (1953), and Henderson (1960). Henderson's method was used in this study.

The comprehensive character of equation 5 for upward continuation is demonstrated in that it can be differentiated to give the first and second derivatives of the gravity field. The second derivative equation uses the same ten ring averages,  $\Delta \overline{\phi}$   $(r_i)$ .

Equation 7 can be written in the following equivalent determinantal form:

$$\Delta \phi(z) \approx -|V|^{-1} \begin{vmatrix} 0 & 1 & -z/a & (z/a)^2 \dots & (-z/a)^n \\ \Delta \phi(0) & 1 & 0 & 0 & \dots & 0 \\ \Delta \phi(-a) & 1 & 1 & 1 & \dots & 1 \\ \Delta \phi(-2a) & 1 & 2 & 2 & \dots & 2^n \\ & & & \ddots & & \ddots & & \\ & & & \ddots & & \ddots & & \\ & & & \ddots & & \ddots & & \\ \Delta \phi(-na) & 1 & n & n^2 & \dots & n^n \end{vmatrix}$$
 (11)

where (V) is the Vandermode determinant obtained by deleting the first two rows and columns of equation 11.

If n equals 5 in equation 11 and differentiating with respect to z the first derivative equation is obtained.

$$\frac{a(24)}{a^{2}} := (a(V))^{-1}$$

$$\begin{vmatrix}
0 & 0 & -1 & -2(z/a) & -3(z/a)^{2} & -4(z/a)^{3} & -5(z/a)^{4} \\
0 & 0 & 0 & 0 & 0 & 0 \\
0 & (-a) & 1 & 1 & 1 & 1 & 1 \\
0 & (-2a) & 1 & 2 & 2^{2} & 2^{3} & 2^{4} & 2^{5} \\
0 & (-3a) & 1 & 3 & 3^{2} & 3^{3} & 3^{4} & 3^{5} \\
0 & (-4a) & 1 & 4 & 4^{2} & 4^{3} & 4^{4} & 4^{5} \\
0 & (-5a) & 1 & 5 & 5^{2} & 5^{3} & 5^{4} & 5^{5}
\end{vmatrix}$$

$$(12)$$

If Z=O in equation 12, first vertical derivatives can be calculated on the plane of observation by the equation

To obtain the sets of coefficients for computing  $[\partial(\Delta\phi)/\partial z]_{z=k}$  using the ten ring averages,  $\Delta\overline{\phi}$   $(r_i)$ , setting a equal to unity, and putting the working upward continuation equation into equation 12, the first derivative working equation is obtained,

$$\frac{\partial (\Delta \phi)}{\partial \mathbf{g}} = k \quad i = 0 \quad D'(r_i, k), \qquad (14)$$

where D'  $(r_1,k)$  are the first vertical derivative coefficients for the appropriate K mesh units below the plane of observation.

Similiarly the second vertical derivative formula can be obtained by differentiating equation 12 with respect to Z.

$$\frac{\partial^{2}(\Delta\phi)}{\partial z^{2}} = -(\mathbf{a}^{2}|\mathbf{v}|)^{T} \begin{vmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ \Delta\phi(-2\mathbf{a}) & 1 & 0 & 0 & 0 & 0 \\ \Delta\phi(-3\mathbf{a}) & 1 & 1 & 1 & 1 & 1 \\ \Delta\phi(-3\mathbf{a}) & 1 & 2 & 2^{2} & 2^{3} & 2^{4} & 2^{5} \\ \Delta\phi(-4\mathbf{a}) & 1 & 3 & 3^{2} & 3^{3} & 3^{4} & 3^{5} \\ \Delta\phi(-4\mathbf{a}) & 1 & 4 & 4^{2} & 4^{3} & 4^{4} & 4^{5} \\ \Delta\phi(-5\mathbf{a}) & 1 & 5 & 5^{2} & 5^{3} & 5^{4} & 5^{5} \end{vmatrix}$$
(15)

The resulting working equation for the second vertical derivative is

$$\begin{bmatrix}
\frac{2}{\partial (\Delta \phi)} \\
\frac{\partial z}{\partial z}
\end{bmatrix}_{\mathbf{z}=\mathbf{k}} = \sum_{i=0}^{10} \Delta \overline{\phi} (\mathbf{r}_{i}) D''' (\mathbf{r}_{i}, \mathbf{k}) \tag{16}$$

where D'' ( $r_1$ , k) are the coefficients of the second derivative.


### Theoretical Gravity Formulas

Talwani's Gravity Computations for Two-Dimensional Bodies

In geological and geophysical studies, many of the structures of interest are horizontally linear and can be approximated by two-dimensional forms of analysis. The outer boundary of any two dimensional body can be approximated by a polygon. This approximation can be made as accurate as one wishes by making the number of sides of the polygon sufficiently large.

The vertical component of gravity due to the polygon can be obtained at any given point and there are no limits on the size or position of the body. The digital computer allows the complex expression for the vertical component of gravity to be solved easily and rapidly.

Given an n sided polygon ABCDE, Figure 10.



Geometrical elements for Talwani's 2-D calculations.

Let P be the point at which the gravitational attraction will be calculated. The point P lies in the xz plane as does the polygon. Let z be defined as positive downwards and  $\theta$  is measured from the positive x-axis downward.

The vertical component of gravity at the origin (P) from the two-dimensional polygon is

$$2G\rho \$zd\theta$$
 (17)

where G is the universal gravitational constant and P the volume density differential, with the line integral being taken around the outer boundry of the polygon (Hubbert, 1948). The above integral is then evaluated for the given polygon. The contribution of a side of the polygon, say, side BC can

be computed. The side CB is extended to meet the x-axis at q at an angle  $\phi_i$ . Let PQ =  $a_i$ , then

$$\mathbf{z} = \mathbf{x} \, \tan \theta \tag{18}$$

for any point R on side BC and

$$\mathbf{z} = (\mathbf{x} - \mathbf{a}_{\mathbf{j}}) \tan \phi_{\mathbf{j}}. \tag{19}$$

From equations 18 and 19

$$\mathbf{z} = \frac{\mathbf{a_i} \quad \tan\theta \tan\phi_i}{\tan\phi_i - \tan\theta} \tag{20}$$

now

$$\mathbf{g} d\theta = \int_{B}^{C} \frac{\mathbf{a}_{i} \tan \theta \tan \phi_{i}}{\tan \phi_{i} - \tan \theta} d\theta = \mathbf{g}_{i}. \quad (21)$$

The vertical component of gravity V can now be given as

$$V = 2Gp \sum_{i=1}^{m} \mathbf{z}_{i}$$
 (22)

with the summations being made over the n sides of the polygon. The integral for  $Z_i$  must now be solved.

For the general case it can by shown that

$$\mathbf{z}_{i} = \mathbf{a}_{i} \sin \phi_{i} \cos \phi_{i} \left[ \theta_{i} - \theta_{i+1} + \tan \phi_{i} \log \frac{\cos \theta_{i} (\tan \theta_{i} - \tan (\phi_{i})}{\cos \theta_{i+1} (\tan \theta_{i+1} - \tan \phi_{i})} \right]$$
(23)

where

$$\theta_{i} = \tan^{-1} \frac{z_{i}}{x_{i}}$$

$$\phi_{i} = \tan^{-1} \frac{z_{i+1}-z_{i}}{x_{i+1}-x_{i}}$$


and

$$a_i = x_{i+1} + x_{i+1} \frac{x_{i+1} - x_i}{x_{i-2_{i+1}}}$$

It should be noted that  $\theta i$ ,  $\theta_{i+1}$ ,  $\phi_i$ , and  $a_i$  can all be explicitly expressed in terms of the  $x_i$ 's and  $z_i$ 's. This is advantageous as the easiest way to define the outer edges of a body is to specify the coordinates of adjacent points or the vertices of the body.

Talwani's Gravity Computations for Three Dimensional Bodies of Arbitrary Shape

A three-dimensional body can be represented by contours on the body. By making the number of sides of a polygon sufficiently large, any irregular outline can be closely approximated. If each contour of the three-dimensional body is replaced by polygonal lamina, the gravitational attraction caused by the lamina can be calculated at any point external of the body. Thus, by interpolating between contour lines and numerical integration, the gravitational attraction of a three dimensional body can be computed.



Geometrical Elements for Talwani's 3-D Calculation

Figure 11 shows geometrically the elements involved in the gravity computations for a three-dimensional body. Point P represents the point at which the gravitational attraction of the three-dimensional mass will be calculated. Point P also is the origin of a left handed cartesian coordinate system with the z-axis positive downward. A contour on the surface of the body is represented by the polygonal lamina ABCDEF ... of infinitesimal thickness dz. Let Ag be the gravitational attraction at P from ABCDEF ....

Then

$$\Delta g = Vds$$
 (24)

where V is the attraction caused by ABCDEF · · · per unit thickness. V can be expressed by a surface integral, and this can be reduced to two line integrals along the boundary of ABCDEF ....

Now

$$V = G_{p}[\$d\Psi - \$E/(r + E)^{\frac{1}{2}}d\Psi], \qquad (25)$$

where G is the universal gravitational constant,  $\rho$  is the volume density differential of the lamina and z,  $\forall$  and rare the cylindrical coordinates used to define the boundary of ABCDEF ....

If P' is the projection of P on the plane ABCDEF .... then PP' equals z. The line r is the radius vector in the plane ABCDEF..., and Y is the angle which it makes with an arbitrary x-axis in this plane, where Y is positive in a clockwise sense from the positive x-axis.

If we evaluate the contribution of the two line integrals from side BC, the first integral gives

$$sd\psi = \psi_{i+1} - \psi_i \tag{26}$$

for  $\overline{P'C}$  and  $\overline{P'B}$ . The second integral can be evaluated by drawing P'J perpendicular from P' to BC. Let P'J equal  $P_i$  and  $\theta_i$  and  $\phi_i$  be respectively the angles which  $\overline{BP'}$  and  $\overline{CP'}$  make with BC. It can be seen from Figure 11 that

$$r = \frac{\text{Pi}}{\sin \left(\phi_i - \psi_{i+1} + \psi\right)} \tag{27}$$

Making this substitution, and noting that  $P_i$ ,  $\theta_i$ , and  $\psi_{i+1}$  are all constants, so that  $\psi$  is the only variable left, the integral can then be solved to give the value,

arc 
$$\sin \frac{z \cos \theta_{i}}{(P_{i}^{2}+z^{2})^{\frac{1}{2}}}$$
 - arc  $\sin \frac{z \cos \phi_{i}}{(P_{i}^{2}+z^{2})^{\frac{1}{2}}}$  (28)

Thus

$$V = G\rho \left[ \psi_{i+1} - \psi_{i} - \arcsin \frac{\mathbf{z} \cos \phi_{i}}{(P_{i}^{2} + \mathbf{z}^{2})^{\frac{1}{2}}} + \arcsin \frac{\mathbf{z} \cos \theta_{i}}{(P_{i}^{2} + \mathbf{z}^{2})^{\frac{1}{2}}} \right].$$
 (29)

The above expression is the attraction caused by the triangular lamina P'BC per unit thickness at point P. The total attraction is then obtained by summing V over the n sides of the polygon to obtain

$$V = G\rho \left[ \sum_{i=1}^{n} \left\{ \psi_{i+1} - \psi_{i} - \arcsin \frac{z \cos \phi_{i}}{(P_{i}^{2} + z^{2})^{\frac{1}{2}}} + \arcsin \frac{z \cos \theta_{i}}{(P_{i}^{2} + z^{2})^{\frac{1}{2}}} \right] \right]$$
(30)

The terms  $P_i$ ,  $\psi_i$ ,  $\psi_{i+1}$ ,  $\cos\theta_i$  and  $\cos\phi_i$  can be expressed in terms of  $x_i$ ,  $y_i$ , the coordinates of B, and  $x_{i+1}$ ,  $y_{i+1}$ , the coordinates of C. Thus V can be expressed in terms of coordinates of the vertices of the polygon ABCDEF....

The attraction caused by equation 30 is only for one lamina. The total attraction  $\Delta g$  from the mass can be evaluated by integrating equation 24 between z-top and z-bottom.

Thus

$$\Delta g$$
 total =  $V d g$ . (31)

### CHAPTER V

### GRAVITY ANOMALIES FROM REEFS

Yungul (1961) reviewed the problem of gravity exploration for reefs and answered some of the classical questions concerning reefs and gravity. Those questions which Yungul sought answers to are: "Do reefs create recognizable gravity anomalies?" and "If they do, what causes these anomalies?" At the time of his review, Yungul found literature concerning reefs and gravity for three geologic provinces.

His summary of the case histories prior to 1961 is shown in Table II which is shown on the following page.

A summary of the literature prior to 1961 as assembled by Yungul is as follows:

- 1. "Reefs frequently show recognizable gravity anomalies which may be (a) a simple high, (b) a high with a negative rim around it, (c) a low with a positive tendency in the center, or (d) a simple low. Most frequently it is a high with a negative rim around it, like a 'sombrero.'
- 2. When the anomaly is of the sombrero type or a simple high, its intensity is of the order of +0.5 mg. This is greater than an allowable density contrast between the reef mass and the enclosing sediments could produce. What is more, a gravity high may be present even when the reef density is less than that of the enclosing sediments. Clayton (1951) reports that at North Snyder the cores show the reef is less dense than the surrounding shales. Yet the anomaly is +0.6 mg.

TABLE II

REEF GRAVITY, AVERAGE CONDITIONS AND RESULTS IN THE THREE REGIONS (after Yungul, 1961)

Geography				-	Geology			Gre	Gravity		
	4								Re	Residual	
Region	to Reef	Height	Width					Regional			Zero
		of Reef (feet)	of Reef (miles)	Ree I Rock	Envelope Formation	Draping	Basal Structure		шВ	шg	Width (miles)
West Texas (Midland Basin)	0002	009	ب د.	caco3	Sandy Shale	у. Ф	Local high over a regional high	Flank, near ape x of a high 3mg/mile		+0.6 2.5	2.5
The Great Lakes Region	0002	300	1.2	٠.	Dolomitic beds, and salt beds thinning or wedging over	With salt Layer: no Without salt Layer: yes	c.	c.		+0.4 1.2	1.2
Alberta	2009	500	1.0	Dolomite	Calcareous Shale	Yes	6.	Flank	+0.25		1.5


- 3. "The existence of a negative rim has so far not been satisfactorily explained.
- 4. The gravity highs are too narrow to be generated by masses at specified reef depths.
- 5. When the anomaly is of the sombrero type, the width of the positive is of the order of the width of the reef, irrespective of the reef depth.
- 6. The local (residual) anomalies are frequently located near the apexes of regional gravity highs.
- 7. As far as west Texas is concerned, the reef bottoms are local positive reliefs. We have no such data for the other regions.
- 8. Thinning and draping in the shales over and around the reefs are more than that indicated by conventional calculations of differential compaction.
- 9. Either because of certain underlying causes, or merely because of world-wide abundance of shales, the surrounding regional formations are shales in most cases.
- 10. The equivalent porosities in the reefs are estimated to be of the order of 12 per cent. The main constituent of most reefs is calcium carbonate. Consequently, the density is about 2.5.
- 11. The reef heights are from about one hundred to about eight hundred feet, widths 1/2 to 3 miles, and the known depths 2,000 to 7,000 ft. The slopes on the flanks may be up to 40 degrees.
- 12. In view of the wide ranges of the observed reef geometries and anomaly types, it seems, at present, that there is no direct, evident relation between the reef mass specifications (depth, height, extent, density) and the gravity anomaly specifications (type, intensity, extent). Consequently, certain factors other than the reef mass play the major role."

Yunguls conclusions, after his study of sedimentation and compaction around the reefs, were:

- 1. Isolated reefs show gravity anomalies on the order of +0.5 mg, with a true negative ring around it.
- 2. No direct relation between reef specifications and gravity anomaly specification can be expected.
- 3. The importance of the reef is not the reef itself but the conditions it created following its development.
- 4. The gravity anomaly depends on what happened after the reef was covered and to some extent pre-reef formations.
- 5. The major cause of the anomaly is probably the lateral variations of sand content in the shales around and above the reef.

He also mentions that while regional surveys could be used to locate reef belts, the potential of the gravity survey at present, is to locate the isolated reefs in a known reef belt.

As stated earlier, one purpose of this study is to define the anomalies from reefs in the Michigan Basin Province. The geological conditions for the Michigan Basin are understandably different from the reef areas discussed by Yungul, thus reef anomalies in the basin also can be expected to show differences from the anomalies of other areas.



PORTION OF REGIONAL GRAVITY MAP
OF MICHIGAN C.I.= I mg

### CHAPTER VI

### INTERPRETATION METHOD

### Regional Gravity of Study Area

The Belle River Mills and Marine City survey areas are situated on the rim of a closed regional gravity minimum. This minimum, shown in Figure 12 (Hinze, 1963), is centered along the northern Macomb and Oakland County border. The Berlin area is situated closer to the center of the minimum than the other areas. The regional gravity of each of the survey areas agrees with the regional gravity pattern shown in Figure 12.

### Marine City Reef Study


### General Statement

The first reef selected for study is the Marine City reef. This reef is buried at a depth of about 2300 feet and has a base which is about one mile long and 3000 feet wide. Two pinnacles about one half mile apart are located on the broad base. These pinnacles have relief of 250 and 300 feet. The step by step procedures used in this study are shown by the flow chart (Figure 7a).

### Observed Bouguer Gravity Anomaly Map

As shown on the regional gravity map, Figure 12, the Marine City survey is located in an area where the regional gravity gradient changes from a north-northwest trend to a western trend. The Bouguer gravity map (Figure 13) shows the regional gradient increasing 3.2 mg from north to south.

There are several anomalous areas which are readily observed. The closed low in the north-central portion of the area is quite noticeable. At first this was believed to be a result of a deep bedrock channel. However, the size and low gravity gradients indicate a much deeper source. The other recognizable anomaly is a north-northeast trending gravity high associated with the Marine City reef bordered on both sides by linear gravity lows. The magnitude of this high determined by the cross profile method is about 0.3 mg. It has a width of about 1.2 miles. The gravity lows have a magnitude of -0.3 mg. The linear nature of the two lows immediately brings to mind the possible presence of bedrock channels. A check of the bedrock topography map (Figure 6), confirms the presence of the channels. sharpness and narrowness of the lows is a good indication that they are caused by features buried at a shallow depth. The outline of the reef mass is shown by dashed lines on all maps.



<u>Th</u>

Ef.

ir

are

li

(1)

st;

gr

in

301 573

gre

:01 ::1

)ec

ear kno

din

Ano

**11**3

### Theoretical Study

### Effects of Bedrock Geology


The bedrock topography in this study area is quite irregular as seen on Figure 6. There is no doubt that the linear gravity lows flanking the Marine City reef anomaly are coincident with the bedrock channels shown by Brown (1963).

Profiles were constructed perpendicular to the strike of the channels at several locations. The gravity effects from the channels were calculated using Talwani's gravity computation method for two-dimensional bodies. The results indicate that the gravity effect of the channels in the area can produce the -0.3 mg anomaly observed on the Bouguer gravity map. Figure 14 shows the profile of a typical bedrock channel in the area and the resulting gravity effect. A density contrast of -0.2 g/cc was assumed for the calculation. The width of the anomaly is about two miles.

From this portion of the study it was concluded that bedrock topographic effects cause significant anomalies which can mask the reef anomaly. When the bedrock topography is known these gravity effects can be calculated using the two-dimensional method and removed from the Bouguer gravity.

Another method of eliminating the bedrock effects will be discussed later.

# TYPICAL BEDROCK 51CHANNEL AND GRAVITY ANOMALY



Expected Gravity Effects from Reefs

The structure of the Marine City reef is well defined from a large number of wells drilled both on and off reef. Structural contour maps of units which exhibit definite lithologic changes were obtained from the Michigan State Geological Survey (Ells, 1964) for the Marine City reef. Using the known geological conditions around the reef, the gravity effects of the reef were calculated using Talwani's three-dimensional technique.

The density contrasts used in the approximation were obtained from two density logs in St. Clair County and from an unpublished density study of the department of Geology, Michigan State University, using well cuttings from the St. 'Clair County area.

The formational units which exhibited a density contrast were divided into sections which could be approximated by a three-dimensional body. The density contrasts of the bodies used for the approximation were both positive and negative. The resulting gravity effect, calculated on the plane of observation, is on the order of +0.006 mg or essentially zero. A geologic cross-section of the reef with the associated gravity anomaly profile is illustrated in Figure 15.

The gravity effects produced only by the thinning of salt over the reef also were calculated. A small positive anomaly of 0.08 mg was obtained. This indicates that the

CROSS-SECTION OF REEF USED IN THREE-DIMENSIONAL GRAVITY 'CALCULATIONS AND THEORETICAL ANOMALY FIGURE 15

salt thinning adds to the anomaly, but is not the primary factor causing the anomaly.

Several conclusions can be made from this part of the study. First, the anomaly is not caused by the geologic structure overlying the reef using average density values for the sediments. Second, the densities used are not representative or not enough information is available concerning the lateral variation of densities surrounding the reef.

A calculation of the theoretical gravity was made based upon a Mg/Ca ratio division of the stratigraphic units around and overlying the reef (Jodry, 1965). A theoretical anomaly of +0.23 mg was obtained using representative density values for the different lithologies as shown in Figure 16. The width of the anomaly is about two miles. These results confirm the earlier conclusion that there is insufficient knowledge of lateral density variations around the reefs.

A brief study was made to determine the depth at which the Marine City reef could be detected if all other factors were kept constant. A depth of burial of 4000 feet will cause an anomaly of about +0.1 mg. A detailed precision survey could detect an anomaly of this size.

The conclusions arrived at from the theoretical study are:

AND THEORY FICAL GHAVITY ANOMALY

# DENSITY VARIATIONS OVER MARINE CITY REEF AND THEORETICAL GRAVITY ANOMALY




FIGURE 16

- 1. The principal cause of the anomaly is not from the reef itself or the post-reef structure alone, but from lateral variations in densities around the reef associated with the post-reef structure.
- 2. The thinning of the salt over the reefs alone cannot produce the observed gravity anomaly.
- 3. Theoretically, the reefs can produce a gravity anomaly which can be detected by detailed surveys.
- 4. A reef with the same set of geologic conditions as the Marine City reef, but buried at twice the depth still can be detected by a detailed gravity survey.

#### Cross-Profile Method

The cross-profile residual is shown in Figure 17.

Each of the anomalous areas discussed on the observed

Bouguer map (Figure 13) are readily isolated by this

technique. The highest magnitude attained is +0.22 mg.

The lows flanking the high have magnitudes of -0.3 mg and

-0.2 mg and the low in the upper-central part of the area

has a residual magnitude of -0.3 mg. In addition, a +0.2

mg high is observed in the left center portion of the area.

In rechecking the observed Bouguer map, an indication of

this anomaly is apparent. This anomaly is associated with

the small China East reef. It was not known by the survey

party that the area included the China East reef and the

fact was not checked until the residual anomaly was isolated by the cross-profile method.

The width of the zero contour of the Marine City reef is 6500 feet. This is less than the value obtained from the theoretical calculations.

There is no question that the cross-profile technique isolates the reef, but the undesirable effects of the bedrock topography are still present.

#### Least Square Polynomial Analysis

A least squares approximation of the original data was made using a 3rd, 5th, 7th, 9th, and 1lth degree polynomial equations. The 3rd and 5th degree approximations isolate the reef anomaly with approximately the same amplitude as the theoretical anomaly. The 7th through 1lth degree equations approximate the original data too closely and no recognizable residual is left. The 3rd and 5th degree residuals are shown in Figures 18 and 19. The extreme edges of the maps were not contoured as anomalous effects are created in areas where station density is limited.

The anomalous areas previously discussed are apparent on both the 3rd and 5th degree residual. The 3rd degree residual does not show a completely closed zero contour around the reef anomaly. This is an indication that the 3rd degree polynomial may not have adequately approximated the regional surface. The 5th degree residual map indicates

a north-northeast trend to the positive anomaly which conforms with the trend of the reef. The magnitude attains a value of 0.19 mg, but the contours only show 0.10 mg due to the 0.10 mg contour interval. The width of the positive anomaly is 6000 feet. This agrees closely with the crossprofile residual, but is less than the theoretical anomaly. The anomaly from the China East reef also is isolated by this technique. The anomalies over the Marine City and China East reefs are the only areas on the map which attain a magnitude of about +0.20 mg.

The data used in the previous discussion were the original data at a 500 foot station spacing. The data were then divided into two groups based on alternate stations, resulting in two maps with a 1000 foot station spacing.

The 3rd and 5th degree polynomial approximations were calculated for both groups of 100 foot stations. The results show no appreciable difference from the results of the 500 foot stations. The 3rd and 5th degree residuals for the even number stations are shown in Figures 20 and 21.

It was concluded from the least squares polynomial analysis that:

1. The least squares analysis using random spaced data, will isolate the reef anomalies, but like the cross-profile method leaves the undesirable effects. 2. A 1000 foot station spacing is sufficient for detailed gravity surveys searching for reefs, however, there must be coverage in the immediate vicinity of the reef other than along the road network.

### Interpolation of Bouguer Gravity

In order to apply the upward continuation, downward continuation, and second derivative techniques, gravity values must be interpolated from the original random spaced data onto a rectangular grid or square mesh system. This can be done manually by overlaying a grid system on the Bouguer map and interpolating gravity values at the desired points or the process can be accomplished with the aid of a computer in a relatively short time. Figure 22 shows the results of interpolating the gravity values on a 500 foot grid system for the Marine City area. There is a slight lateral shift in some gravity values on this map due to the interpolation process, but it is small enough to be unimportant.

# Downward Continuation and Second Derivative on Original Surface

As described in the section on upward and downward continuation theory, the term  $\Delta \bar{\phi}$   $(r_i)$  is the average of the values computed on ten rings surrounding a center point. If all ten rings were used with a mesh interval of 500

feet, the diameter of the largest ring would be greater than the distance across the survey area. Therefore, a test was made using only seven rings. The results were compatible with the known reef anomaly and it was assumed that seven rings give the approximation desired. Using seven rings the radius of the largest ring is  $\sqrt{50}$  times the mesh interval.

By calculating the gravity values on a plane below the original observation plane the perceptibility of gravity values is increased. The Bouguer gravity values were calculated on planes m mesh intervals below the original plane where m=1, 2, 3, 4, 5 and the mesh interval equals 500 feet.

The size of the interpolated grid system is an important factor in the results as the mesh interval chosen depends on the lateral extent of the anomalies and the sharpness of the gradients. If the mesh interval is too small, oscillation will develop and if it is too large only regional anomalies will remain (Henderson, 1960). A mesh interval of about one-fourth the depth to the top of the mass causing the anomaly is usually satisfactory (Henderson, 1960; Vacquier, 1951; and Nettleton, 1940).

There was no correlation between any of the anomalies on the downward continued planes and the anomalies on the original plane. The anomalies on the downward continued planes showed an oscillation character indicating the mesh interval was too small.

#### Upward Continuation

depth are minimized by calculating the Bouguer gravity values on a plane above the original plane of observation. The gravity values were calculated on surfaces m mesh intervals above the original plane where m = 1, 2, 3, 4, 5 and the mesh interval equals 500 feet. The Bouguer gravity values on the plane five mesh intervals or 2500 feet above the original plane are shown in Figure 23. This method eliminates most of the small sharp anomalies and leaves the broad anomalies as desired. The anomaly over the Marine City reef is present but the anomaly from the China East reef is almost eliminated. The large low in the northern portion of the area is still present indicating a source other than a bedrock channel.

# Downward Continuation of Upward Continued Surfaces

The size of the rings used in calculating  $\Delta \bar{\phi}(r_1)$  cause an area to be lost around the periphery of the original area. When using a 500 feet mesh interval, the center point of the rings must be moved toward the center of the map a factor of seven mesh intervals before a complete set of values will be obtained as shown in Figure 24. This means a distance of seven mesh intervals is lost around the edge of the map when the upward continuation surface is calculated and an equal distance is lost from the upward continued surface when the downward continuation and second derivative values are calculated.

If the downward continuation and second derivatives techniques are performed on the upward continued surface, which is void of undesirable effects, the reef anomaly may be isolated. The upward continuation surface 2500 feet above the original surface was downward continued one through five mesh intervals maintaining the 500 foot mesh interval. The reef anomaly was not isolated by the downward continuation method, however, it was apparent that isolation of the reef was increasing for each successively lower plane. reef anomaly is successfully isolated on the plane 5000 feet below the 2500 foot upward continued surface as shown in Figure 25, however, the anomaly is narrower than obtained by previous methods. The maximum width is approximately 5000 feet. This is not unrealistic as the plane on which the Bouguer gravity is calculated is about equal to the depth of the reef. The magnitude of the Bouguer gravity values are relative to one another and are in milligals. anomalies attain a magnitude equal to the reef anomaly.

Nettleton (1954) compared the results of several derivative techniques using a mesh interval of one-fourth, one-half, and equal to the kepth of the source. The results indicate that while a mesh interval of one-fourth the depth of the source in most cases is optimum, a mesh interval of one-half the depth also is satisfactory. Because the 500 foot mesh interval appears to be too small for this study area, the mesh interval was changed to 1000 feet. When the

mesh interval equals 1000 feet, only five rings of the template are used. This was tested and the results indicated the approximation using five rings was adequate for this type of survey.

By changing the mesh interval on the upward continued surface from 500 feet to 1000 feet the downward continued planes can be calculated at intervals of 1000 feet. This effectively places the highest upward continued plane at 5000 feet above the original plane. By placing the planes of observation higher, the Bouguer gravity values show little change, except for a smoothing effect.

The downward continuation technique, using the 1000 foot mesh interval, applied to the 3000 and 5000 foot up-ward continuation surfaces successfully isolates the reef anomaly on the plane 5000 feet below each upward continued surface as shown in Figures 26 and 27.

The anomaly on the plane 5000 feet below the 3000 foot upward continued plane is narrow with a maximum width of 4000 feet. This is consistent with the previous downward continuation results. The magnitude of the closed anomaly is +0.8 mg.


The anomaly on the plane 5000 feet below the 5000 foot upward continued surface is unique in that it isolates not one but two anomalies which almost coincide with the two pinnacles of the Marine City reef. The plane on which the two anomalies are isolated is essentially the same as


the original plane. The magnitude of the closed anomalies is +0.3 mg which is near the value of the original observed anomaly. The widths of the anomalies are approximately 5000 feet.


The Marine City reef anomaly is thus isolated on planes 5000 feet below the upward continuated surfaces of 2500, 3000, and 5000 feet. It is concluded that by using a mesh interval of one-half the depth of the reef, the downward continuation method will isolate the reef anomalies from a surface upward continued a distance equal to the depth of the reef.


#### Second Derivatives


The second derivative technique of Henderson (1960) was applied to each upward continuation surface calculated. When the second derivative is calculated using seven rings and a 500 foot mesh interval, an oscillation effect results and no isolation of the reef anomaly occurs. When the derivatives are calculated using five rings and a 1000 foot mesh interval, the reef anomaly is isolated from the 2500, 3000, and 5000 foot upward continued surfaces (Figures 28, 29, and 30). The magnitudes of the reef anomaly produced by the second derivative techniques are not duplicated elsewhere on the map, however, the magnitude of the second derivative values were not evaluated quantitatively. The width of the anomalies is generally greater than those produced by the

















# BOUNDARIES OF ANALYTICAL METHODS









FIGURE 24














downward continuation technique below the surface, but less than the original observed and theoretical anomaly. They are on the order of 5000 feet wide. The anomaly isolated by the second derivative method from the 5000 foot upward continued surface (Figure 30) is actually two anomalies coinciding with the two pinnacles of the Marine City reef. This is consistant with the results of the downward continuation techniques.

It is concluded that the second derivative technique has a high resolving power and will successfully isolate the reef anomaly using a mesh interval of one-half the depth of the reef on upward continued surfaces calculated on planes approximately equal to the depth of the reef.

# Belle River Mills Reef Study

### General Statement

The second reef studied is the Belle River Mills reef. This reef is about three miles north of the Marine City reef and possesses several characteristics which are different from the Marine City reef. The two reefs are buried at approximately the same depth, but the Belle River Mills reef has a greater areal extent and is not a pinnacle type reef. The reef maintains a relief of 300 feet along its length of two miles. Its overall width is about 3000 feet. Compared to the Marine City reef, the Belle River Mills reef is greater in both areal extent and reef mass.

# Observed Bouguer Gravity Anomaly Map

The Belle River Mills area is situated in an area where the regional gravity gradient flattens out and changes from a northwest direction to a western direction as shown in Figure 12. The resulting pattern for the area is a saucer shaped depression with a maximum change in magnitude of only 0.9 mg (Figure 31). Three anomalous areas are readily apparent on the Bouguer gravity map. A gravity high with a magnitude of 0.3 mg trends northeast across the center of the area. Two closed gravity lows with magnitudes of -0.15 mg and -0.35 mg flank the gravity high. The northernmost part of the high is associated with the Belle River Mills reef. The southern portion of the high will be discussed later.

Comparing the Bouguer gravity map with the bedrock topography map (Figure 6), it can be seen that several bedrock features coincide with the gravity anomalies in the area. The linear low on the southeast side of the high does not correlate with any bedrock channels, but well control in this area is sparce and there is a good possibility that a bedrock channel exists. The trend of the low can be followed into the Marine City area to the south. A major bedrock channel does exist in the area bordering the southern edge of the Belle River Mills area and the northern edge of the Marine City area. The channel predicted for the Belle River Mills area could possible be a tributary of this bedrock

channel. For comparison purposes, the gravity stations on the southern border of the Belle River Mills area are coincident with those on the northern border of the Marine City area.

# Theoretical Study

# Effects of Bedrock Topography

The gravity low to the northwest of the reef anomaly coincides with a broad bedrock depression. The gravity effect from this depression was calculated using Talwani's two-dimensional method. The resulting theoretical anomaly of -0.15 mg indicates the depression can cause the observed anomaly.

### Expected Gravity Effects from Reef

The theoretical gravity effect of the Belle River Mills reef was calculated using Talwani's three-dimensional method. The average density contrasts were used because information on lateral density variations around the reef are not available. The calculated gravity anomaly is on the order of 0.006 mg. This is consistant with the anomaly calculated from the Marine City reef.

#### Results of Isolation Techniques

### General Statement

The isolation techniques used in the Belle River Mills study were essentially the same as employed in the Marine

City study. These are the cross-profile, upward continuation, downward continuation, and second derivative methods.

#### Cross-Profile Method

The cross-profile residual is shown in Figure 32. An anomaly is associated with the reef, but it does not isolate the reef satisfactorily. The width of the zero contour is nearly two miles and the magnitude is +0.2 mg.

#### Least Squares Polynomial Analysis

The same degree polynomial approximations calculated for the Marine City area were used for this area. Again the 3rd and 5th degree polynomial equations for both the 500 and 1000 foot station spacing isolate the reef anomaly. The residual maps are shown in Figures 33, 34, 35, and 36. For this area, however, the 3rd degree fit was better than the 5th degree fit. This may be explained by the fact that the regional gradient is very small. The 5th degree polynomials for both the 500 foot and 1000 foot station spacing tend to approximate the original surface too closely.

The magnitude of the reef anomaly from the least square polynomial analysis is +0.18 mg. This is almost identical with the value obtained for the Marine City reef. It had been expected that the Belle River Mills reef would cause an anomaly larger than the Marine City reef. The similarity of the magnitudes substantiate the previous conclusion that the anomalies are not caused by the reef, but

by post reef conditions. The southern high is also indicated with a magnitude of about +0.13 mg.

It is concluded from the comparison of the Belle River Mills and Marine City least squarea analysis that the degree of approximation depends on the nature of the regional gravity pattern and not the nature of the anomalies.

#### Interpolation of Bouguer Gravity

Several spurious effects are created in the interpolation of the Bouguer gravity for the Belle River Mills area (Figure 37). This is an inherent problem in interpolating the Bouguer gravity by the method adopted for the computer. These spurious effects are generally created in an east-west direction. The spurious effects on the Belle River Mills interpolated gravity map are near the edge of the reef anomaly but are not directly associated with the anomaly.

# Downward Continuation and Second Derivative on Original Surface

The results were identical with the Marine City study and confirmed the previous conclusions that a mesh interval of one-fourth the depth to the anomaly causing mass is too small.

#### Upward Continuation

The upward continued surface 2500 feet above the original plane of observation is shown in Figure 39. The reef anomaly remains with two minimum areas on either side. The anomaly on the northwest side is the anomaly discussed

previously which coincides with a bedrock depression. It was not removed in the upward continuation method indicating the anomaly is too broad.


# Downward Continuation of Upward Continued Surfaces

The same techniques applied to the Marine City study were used for the Belle River Mills area. The only combination which successfully isolates the reef anomaly is the downward continuation technique which calculated the values on a plane 5000 feet below the 5000 foot upward continued surface (Figure 39). A gravity high with the same magnitude of the reef anomaly also is isolated to the south of the reef anomaly.


The values calculated 5000 feet below the 2500 and 3000 foot upward continued surfaces failed to isolate the reef, but showed a general gravity high over the area. The only major difference between the Marine City study area and the Belle River Mills study area is the regional gravity gradient. It is possible that the regional gradient of an area can cause the downward continuation techniques to vary in their isolation capabilities.

#### Second Derivatives


The second derivative values were calculated for the upward continued surfaces of 2500, 3000, and 5000 feet. Each of the second derivative calculations successfully isolate

































4



the Belle River Mills reef anomaly (Figures 40, 41 and 42). This is consistant with the Marine City results.

The second derivative techniques also isolate the gravity high to the south of the reef anomaly. The magnitudes of the two anomalies are again about the same. There is a good possibility that another reef may exist to the south of the Belle River Mills reef.

The results of the second derivative method on the Belle River Mills area indicate that the isolation power of the derivative technique is greater than the downward continuation techniques.

## Berlin Reef Study

## General Statement

The Berlin reef is situated more basinward than the Marine City and Belle River Mills reefs and is buried at a depth of about 3500 feet. The reef has a width of about 2500 feet and a length of one mile. The major portion of the reef mass is salt filled, but there are several producing wells in the reef.

This study was made assuming that nothing was known about the conditions surrounding the reef. A theoretical study was not made and the cross profile method was not employed.

## Observed Bouguer Gravity Anomaly Map

The Berlin area, as shown on the regional gravity map (Figure 12) is located near the center of a closed regional gravity minimum. The Bouguer gravity map is shown in Figure 43. The nature of the regional gradient is readily observed on the Bouguer map by the semi-circle pattern of the contours. There are several anomalous areas on the Bouguer map, but the anomaly associated with the reef is not as obvious as in previous areas. Two linear lows flank a small positive trend which enters the area from the northeast.

Well control in this area is limited and no bedrock channels are indicated on the bedrock topography map (Brown, 1963). However, it is reasonable to assume that the linear gravity lows are reflections of bedrock channels.

#### Results of Isolation Techniques

#### Least Squares Polynomial Analysis

The results of the least squares polynomial analysis for the Berlin Survey are no different than the Marine City and Belle River Mills areas. The 3rd and 5th degree polynomial approximation for the 500 and 1000 foot station spacing isolate a positive anomaly over the reef. Negative areas are present on the west and east sides of the reef anomaly. The magnitude of the anomaly is again about +0.20 mg. This reaffirms the conclusion that the anomaly is not due to the

reef itself. The least square residual maps are shown in Figures 44, 45, 46 and 47.

### Interpolation of Bouguer Gravity

Again a spurious effect was created by the interpolation technique, but it is not in the reef area and, therefore, was disregarded. The interpolated Bouguer gravity map is shown in Figure 48.

#### Upward Continuation Technique

The results of the gravity values calculated on the plane 2500 feet above the original surface is shown in Figure 49. The 2500 foot plane does not show the effects of the linear gravity lows, indicating further that a possibility of bedrock channels exists. A high is still present in the vicinity of the Berlin reef.

#### Downward Continuation

The downward continuation techniques failed to isolate the reef anomaly. The gravity values downward continued 5000 feet below the 2500, 3000 and 5000 foot upward continued surface showed a tendency to isolate the reef anomaly, but only indicate a positive area in the reef vicinity.


#### Second Derivative


The second derivatives were calculated for the 2500, 3000 and 5000 foot upward continued surfaces. Each of the techniques successfully isolate the anomaly over the reef


(Figures 50, 51 and 52). For the values calculated from the 2500 and 3000 foot upward continued surfaces the closed contour with the highest value is essentially coincident with the reef pinnacle. The overall anomaly is about 5000 feet wide. The values calculated on the 5000 foot surface show the broad anomaly from the reef, but do not show the closure coincident with the reef pinnacle. This may indicate that the distance between the depth of the reef and the upward continued surface was too great and a portion of the anomaly was lost.


The conclusions reached from the Berlin study are:


- The second derivative techniques have a higher resolving power than the downward continuation techniques.
- 2. The upward continued surface should not be calculated higher than about 7500 feet above the expected reef depth.
- 3. The salt filling of the reef mass has no bearing on the magnitude of the observed anomaly.


























#### CHAPTER VII

#### SUMMARY

## Detection of Reef Anomalies

The results of this study show conclusively that detailed gravity surveys can detect the anomalies over reefs in Michigan. A known reef buried at 3500 feet was successfully detected and theoretical calculations indicate that reefs buried at depths up to 4000 feet probably can be detected.

## Characteristics of Reef Anomalies

Based on observations the characteristics of reef anomalies for the Michigan Basin are summarized as follows.

- 1. The gravity anomalies over reefs are positive with magnitudes of about +0.20 mg.
- 2. The width of the isolated anomalies varied with the isolation method. The anomalies isolated by the least squares method are about twice as wide as the reef mass. The downward continuation of an upward continued surface method gives a ratio of about 1.3 to 1 for the width of the anomaly to the width of the reef. The second derivative method gives a ratio from 1.3 to 1 for the 2500' upward continued surface up to 1.6 to 1 for the

5000' upward continued surface. The cross profile method on the Marine City reef gives a ratio of 2:1. The graphical and statistical methods of isolation indicate a larger anomaly width to reef width ratio than do the analytical methods. These results are consistant for the Marine City, Belle River Mills, and Berlin reefs.

- 3. The reef anomalies have adjacent parallel to subparallel negative anomalies associated with
  the positive anomalies. These can be accounted
  for by theoretical calculation of bedrock topographic effects, however, the coincidence of the
  negative anomalies indicate they are not fortuitous
  but may be controlled by structure in the sediments
  or they are coming from a deeper source.
- 4. The reef anomalies are not associated with any regional gravity pattern.

# Sources of Reef Anomalies

The source of the anomalies observed over reefs in the Michigan Basin are not the reef mass, but the conditions created after reef development. The probable source of the anomalies is the lateral variation in density overlying the reef. In the Michigan Basin these density contrasts are between dolomite and limestone and dolomite and salt.

# Optimum Methods for Isolation of Reef Anomalies in Michigan

A major purpose of this study is to establish optimum methods for the isolation of reef anomalies in this geologic province. The study suggests the following steps as an optimum method of isolation of reef anomalies in the Michigan Basin:

- 1. A precision detailed gravity survey is necessary. The error for field procedures and basic reduction of the data must be kept to a maximum of 0.05 mg.
- A station density of 1000 feet is sufficient in 2. searching for reefs by the gravity method. ever, the anomalies can change rapidly in a short distance. Thus, it is adviseable to place stations at a greater detail than the road networks will allow. If the survey is being used to detail a suspected reef, the immediate area of interest should contain complete data coverage. overall area of gravity coverage should extend at least three miles out on every side of the This is necessary to allow area of interest. the use of various isolation techniques and include the regional gravity of the area.
- 3. The cross profile and least squares polynomial analysis methods are not completely satisfactory

for isolating the reef anomalies. These methods include many anomalies from undesirable sources. It is not recommended that these methods by employed.

- 4. The upward continuation method is a useful method for eliminating the gravity effects of shallow sources. By calculating the upward continued surface on a plane greater than the suspected reef depth, using a mesh interval of one half this depth, the shallow sources are satisfactorily removed. Total separation between the suspected reef and the upward continued plane should not be greater than about 7500 feet.
- 5. The reef anomalies may be isolated by downward continuing the upward continuation surfaces to depths not exceeding the suspected reef depth.
- 6. Second derivative calculations of the upward continued surfaces have a high resolving power in isolating the reef anomalies. Henderson's (1960) formula seems to be optimum for this calculation.

## Recommendations for Further Study

The following recommendations are made for future study of reefs in the Michigan Basin by the gravity method.

1. A similar study should be performed over several reefs located around the periphery of the basin.

Geologic conditions of southeastern Michigan are not constant around the basin, thus, gravity anomalies may be different from those observed in this study.

- 2. A similar study performed over a reef where bedrock topographic features are not present. The study should include an investigation of the relationship of the parallel negative anomalies to the bedrock features and possible structural sources.
- 3. A regional study of the location of known reef areas in the Michigan Basin and their relationship to the regional structure of the basin.

BIBLICGRAPHY

#### BIBLICGRAPHY

- Alguire, S. L. (1962) Some geologic and economic aspects of Niagaran reefs in Eastern Michigan: Michigan Basin Geological Society Annual Field Conference, p. 30-38.
- Brown, R. E. (1963) Bedrock topography, lithology and glacial drift thickness of Lapeer and St. Clair Counties, Michigan; Masters Thesis, Michigan State University.
- Cohee, G. V. (1948) Thickness and lithology of Upper Ordovician and Lower and Middle Silurian rocks in the Michigan Basin: U. S. Geol. Survey, Oil and Gas Invest., Preliminary Chart 33.
- Cumings, E. R., and Shrock, R. S. (1928) Niagaran coral reefs of Indiana and adjacent states and their stratigraphic relationships: Geol. Soc. Amer. Bull. V. 39, p. 579-620.
- Ehlers, G. M., and Kesling, R. V. (1962) Silurian rocks of Michigan and their correlations: Michigan Basin Geological Society Annual Field Conference, p. 1-20.
- Elkins, Thomas A. (1950) The second derivative method of gravity interpretation: Geophysics, V. XVI, Jan., no. 1:29-50.
- Ells, G. D. (1964) Personal Communication.
- Geyer, R. A. (1962) Uses and abuses of gravity surveys in reef exploration: Michigan Basin Geological Society Annual Field Conference, p. 69-73.
- Grant, F. G. (1957) A problem in the analysis of geophysical data: Geophysics, Vol. 22, no. 3, p. 309-344.
- Henderson, R. G. (1960) A comprehensive system of automatic computation in magnetic and gravity interpretation: Geophysics, Vol. 25, no. 3, p. 569-585.
- Henderson, R. G., and Zietz, Isadore. (1949) The computation of the second vertical derivatives of geomagnetic fields: Geophysics, Vol. 14, p. 508-534.

- Hinze, W. J. (1963) Regional gravity and magnetic maps of the Southern Peninsula of Michigan: Michigan Dept. of Conservation, Geological Survey Division, Report of Investigation 1.
- Hubbert, M. K. (1948) Gravitation terrain effects of two-dimensional topographic features: Geophysics, Vol. 13, No. 2, p. 226-254.
- Jodry, R. L. (1965) Personal Communication.
- Klasner, J. S. (1964) A study of buried bedrock valleys near South Haven, Michigan by the gravity method:
  Masters Thesis, Michigan State University.
- Leverett, F. (1939) Correlation of beaches with moraines in the Huron and Erie basins: Amer. Jour. of Sci., Vol. 237, p. 456-475.
- Lowenstam, H. A. (1950) Niagaran reef of the Great Lakes area: Jour. Geology, Vol. 58, p. 430-487.
- Lowenstam, H. A. (1957) Niagaran reefs in the Great Lakes area: Geol. Soc. Amer., Mem. 67, Treatise on marine ecology and paleoecology, V. 2, Paleoecology chapt. 10, p. 215-248.
- Manely, T. R. (1964) The Niagaran rocks of Drummond Island, Michigan and the related rocks of the Michigan Basin: Phd. D. Thesis, Michigan State University.
- Nettleton, L. L. (1940) Geophysical prospecting for oil: McGraw-Hill, New York.
- Nettleton, L. L. (1954) Regional, residual, and structures: Geophysics, Vol. 19, p. 1-22.
- Newcombe, R. B. Oil and gas development in Michigan: Mich. Geol. Survey Pub. 37, Series 31, p. 156-157.
- Peters, L. J. (1949) The direct approach to magnetic interpretation and its practical application: Geophysics, Vol. 14, No. 3, p. 290-319.
- Pohly, R. A. (1954) Gravity case history: Dawn No. 156 pool, Ontario: Geophysics, Vol. 19, p. 95-103.
- Rosenbach, Otto. (1953) A contribution to the computation of the "second derivative" from gravity data: Geophysics, Vol. 18, No. 4, p. 894-912.

- Skeels, D. C. (1947) Ambiguity in gravity interpretation: Geophysics, Vol. 12, p. 43-56.
- Talwani, M., Worzel, J. Lamar, and Landisman, M. (1959)
  Rapid gravity computations for two-dimensional bodies with application to the Mendocino submarine fracture zone: Jour. Geophys, Res., Vol. 64, p. 49-59.
- Talwani, M., and Ewing, M. (1960) Rapid computation of gravitational attraction of three-dimensional bodies of arbitrary shape: Geophysics, Vol. 25, No. 1, p. 203-225.
- Textoris, D. A., and Carozzi, A. V. (1964) Petrogrophy and evolution of Niagaran (silurian reefs, Indiana: Bull. A. A. P. G., Vol. 48, No. 4, p. 395-426.
- Vacquire, V., Steenland, N. C., Henderson, R. G., and Zietz, I. (1951) Interpretation of aeromagnetic maps: Geol. Soc. Am. Mem. 47, p. 151.
- Yungul, S. H. (1961) Gravity prospecting for reefs: Effects of sedimentation and differential compaction: Geophysics, Vol. 26, No. 1, p. 45-56.

## APPENDIX

# GRAVITY DATA OF ST. CLAIR COUNTY REEF STUDY

Sample forms of data:

x and y -- 430000(feet)

x's and y's are coordinates based on Michigan Coordinate System, East Zone.

Elevation -- 605.50 feet

Observed Gravity -- 54.27 milligals

Meter Reading -- 542.7 scale divisions

Marine City and Belle River Mills Surveys:
World Wide Meter #45
Calibration Constant 0.10093 mg/S.D.

Berlin Survey
Worden Meter #99
Calibration Constant 0.0995(1)mg/S.D.

The following key is to be used for the data tables on the following pages:

_	Station				Observed	
Survey	No.	Х	У	Elevation	Gravity	Reading
					<del></del>	
1	2	3	4	5	6	7

# MARINE CITY SURVEY

1 2	2 1 2	5	6
MARINECYBASEA	803167440932	59900	5461
MARINECYBASEB	795148437793	60258	5398
MARINECYBASED	811036442610	58637	5557
MARINECYBASEE	810597449379	58653	5628
MARINECY 23-F	800029448826	60404	5504
MARINECY H-G	803387446310	59095	5573
MARINECY 33-H	800274443444	60059	5454
MARINECY001MC	803115441467	59629	5486
MARINECY002MC	803094441956	59332	5514
MARINECY003MC	803090442231	59281	5525
MARINECY004MC	803084442731	59484	5518
MARINECY005MC	803117443206	59377	5533
MARINECY006MC	803187443657	59337	5547
MARINECY007MC	803305444159	59129	5564
MARINECY008MC	803409444635	59046	5573
MARINECY009MC	803453445111	59046	5580
MARINECY010MC	803370445710	59085	5580
MARINECY012MC	803430446845	59022	5580
MARINECY013MC	803483447361	59022	5586
MARINECY014MC	803517447836	59323	5576
MARINECY015MC	803560448362	59395	5581
MARINECY016MC	803687448988	59226	5578
MARINECY017MC	803202448982	59381	5577
MARINECY019MC	802178448919	59688	5562
MARINECY020MC	801653448902	59949	5538
MARINECY021MC	801153448880	60115	5529
MARINECY022MC	800653448864	60175	5525
MARINECY024MC	800031448301	60237	<b>5</b> 511
MARINECY025MC	800062447791	60195	5507
MARINECY026MC	800089447281	60121	5508
MARINECY027MC	800116446727	60095	5508
MARINECY028MC	800149446117	60202	5488
MARINECY029MC	800180445653	60255	5479
MARINECY030MC	800202445118	60193	5478
MARINECY031MC	800225444544	60051	5478
MARINECY032MC	800243443929	60009	5473
MARINECY034MC	800291442920	59924	5447
MARINECY035MC	800318442395	59785	5458
MARINECY036MC	800355441870	59620	5477
MARINECY037MC	800377441346	59349	5496
MARINECY038MC	800409440796	59410	5499
MARINECY039MC	800934440803	59360	5501
MARINECY040MC	801443440835	59283	5508
MARINECY041MC	802008440867	5 <del>94</del> 71	5495
MARINECY042MC	802533440899	59539	5489

ı	2	3	4	5	6
MARINE	ECY043MC	80364	1442203	58696	5561
MARINE	ECY044MC	80415	1442145	58454	5574
MARINE	ECY046MC	80521	8442059	58500	5567
MARINE	ECY047MC	80564	3442000	58699	5549
MARINE	ECY049MC	80655	0442100	59273	5507
MARINE	ECY050MC	80707	75442200	59321	5502
MARINE	ECY051MC	80751	1442239	59246	5519
MARINE	ECY052MC	80798	9442320	58968	5529
MARINE	ECY053MC	80846	4442376	58817	5532
MARINE	ECY054MC	80891	3442432	58702	5527
MARINE	ECY055MC	80937	7442488	58747	5521
MARINE	ECY056MC	80983	37442544	58709	5534
MARINE	ECY057MC	81026	6442575	58817	5535
MARINE	ECY058MC	81073	86442606	58812	5546
	ECY059MC		4443110	58816	5550
	ECY060MC	81095	4443549	58803	5551
	ECY061MC	81094	3444009	58734	5554
	ECY062MC	81092	2444458	58743	5549
MARINE	ECY063MC	81090	6444908	58765	5548
	ECY064MC		35445383	58736	5560
	ECY065MC		9445833	58700	5573
	ECY067MC		32446767	58703	5588
	ECY068MC		27447117	58659	5598
	ECY069MC		5447682	58673	5601
	ECY070MC		3448146	58713	5607
	ECY071MC		2448606	58697	5616
	ECY072MC		7449380	58604	5633
	ECY073MC		23449359	58702	5626
	ECY074MC		3449338	58685	5625
	ECY075MC		3449312	58751	5626
	ECY076MC		73449281	58717	5621
	ECY078MC		4449220	58726	5616
	ECY079MC	_	4449204	58678	5620
	ECYOBOMC		5449183	58673	5620
	ECYOB2MC		0449121	58739	5623
	ECYOB3MC		1449090	58678	5628
	ECYOBAMC		1449060	58723	5631
	ECYO85MC		2449019	58786	5616
	ECYOB7MC		3440507	59845	5460
	ECYO88MC	_	4440033	59739 59869	5461
	ECY089MC	_	84439623	5 <del>9</del> 927	5444
	ECYO9OMC		5439172		5435
	ECY092MC ECY093MC		32438310	60192	5395
	ECYO94MC		3437854 3437428	60196 60198	5385 5373
	ECYO95MC		9436952	60191	5363
	ECYO96MC		0436490	60185	5349
	ECYO97MC		10436490 17435989	60243	5328
	ECYO98MC		8435510	60162	5322
	ECY099MC	,	4435065	60044	5338
	ECY100MC		5434601	59943	5340
	ECY101MC		6434156	59911	5332
******* * 1.46	LOTTOLFIC	50257	J-J-1.50	J / F & 1	JJJE

1	2	3		4	5	6
MARINECY	102MC	80268	743	3707	59697	5341
MARINECY	103MC	80269	343	3257	59677	5331
MARINECY	104MC	80258	443	2 <b>7</b> 95	59697	5321
MARINECY		80248			59656	5315
MARINECY		80237			59570	5317
MARINECY		80224			59384	5325
MARINECY		80210			59306	5321
MARINECY		80202			59253	5324
MARINECY	-	80199			59126	5327
MARINECY		80696			58515	5618
MARINECY		80739			58483	5612
MARINECY		80788			58516	5609
MARINECY	121MC	80833			58503	5610
MARINECY	122MC	80876	944	6560	58498	5605
MARINECY	123MC	80921	944	6581	58511	5603
MARINECY	124MC	80968	444	6602	58567	559 <b>6</b>
MARINECY	125MC	81014	844	6623	58659	5597
MARINECY	126MC	81066	644	9830	58701	5631
MARINECY	127MC	81063	445	0379	58704	5638
MARINECY	128MC	81060	345	0829	58691	5643
MARINECY	130MC	81056	645	1793	58692	5652
MARINECY	131MC	81053	545	2350	58765	5646
MARINECY	132MC	81050	345	2772	58737	5651
MARINECY	133MC	81047	245	3262	58729	5656
MARINECY	134MC	81045	645	3722	58778	5662
MARINECY	135MC	81041	945	4201	58793	5677
MARINECY	136MC	81039	845	4676	58794	5687
MARINECY		81037	445	4975	58875	5687
MARINECY	_	80379	644	9440	59105	<b>5</b> 593
MARINECY		80391			59060	5598
MARINECY		80403			59205	<b>5</b> 590
MARINECY		80420			59269	5588
MARINECY		80436			59303	5598
MARINECY		80451			59246	5610
MARINECY	_	80466		-	59265	5609
MARINECY	_	80482			59176	5621
MARINECY	_	80499			59178	5626
MARINECY		80510		_	59048	5638
MARINECY		80508			59249	5630
MARINECY		80506			59315	5636
MARINECY		80503			59402	5634
MARINECY		79952			60320	5507
MARINECY		79904			60539	5499
MARINECY		79856			60744	5495
MARINECY	_	79810			60774	5503
MARINECY	· · ·	79760			60765	5512
MARINECY		79995			60353	5503
MARINECY		79992			60377	5504
MARINECY MARINECY		79991			60389 60555	5512 5518
MARINECY		79985 79984			60719	5518 5521
DIRINECY		79976			60898	5521 5538
DIKINECT	104110	17710	<del></del> 5	J00 /	20090	2230

		306		
1	2	126 3 4	5	6
DIRINEC		789566440189	60436	5422
DIRINEC		802468438213	60193	5390
DIRINEC	_			
		801968438192	60096	5401
DIRINEC		801443438160	59918	5415
DIRINEC		800959438119	59730	5432
DIRINEC		800444438082	59574	5446
DIRINEC		799944438065	59527	5455
DIRINEC		799420438034	59568	5446
DIRINEC		798920438002	59685	5430
DIRINEC		798396437970	59745	5423
DIRINEC		797881437938	59903	5418
MARINEC		797375437950	60009	5416
DIRINEC		796867437880	60056	5425
DIRINEC		796347437858	60119	5424
DIRINEC		795847437827	60218	5416
DIRINEC	-	794648437751	60263	5395
DIRINEC	Y241MC	794124437714	60276	5384
DIRINEC	Y242MC	793619437693	60291	5388
DIRINEC	Y243MC	793124437666	60307	5388
DIRINEC	Y244MC	792620437630	60318	5387
DIRINEC	Y245MC	792115437608	60256	5387
DIRINEC	Y247MC	791615437581	60226	5391
DIRINEC	Y248MC	791085437549	60165	5392
DIRINEC	Y249MC	790576437518	60123	5385
DIRINEC	Y250MC	790141437487	60042	5393
DIRINEC	Y251MC	789802437458	60020	5390
DIRINEC	Y252MC	802120434969	59860	5345
DIRINEC	Y253MC	801668435163	59812	5346
DIRINEC	Y254MC	801232435207	59634	5361
DIRINEC	Y255MC	800759435056	59513	5366
DIRINEC	Y256MC	800285434960	59503	5376
DIRINEC	Y257MC	799784435029	59371	5394
DIRINEC	Y258MC	799267435157	59300	5410
DIRINEC	Y259MC	798796435301	59288	5427
DIRINEC		798266435269	59521	5415
DIRINEC	Y261MC	797781435237	59783	5400
DIRINEC		797282435211	59801	5402
DIRINEC	Y263MC	796792435184	59874	5399
DIRINEC		796267435158	59797	5411
DIRINEC		795732435136	59770	5414
DIRINEC		795283435105	59776	5415
DIRINEC	_	803518434372	58242	5432
DIRINEC		803996434169	58019	5430
DIRINEC		804468433975	58045	5426
DIRINEC		804951433756	58084	5417
DIRINEC		805404433547	58093	5406
DIRINEC		805866433368	58485	5372
DIRINEC		806334433159	58562	5355
DIRINEC		806797432960	58980	5317
DIRINEC		807265432762	59666	5263
DIRINEC		807727432553	59261	5294
DIRINEC		808210432349	58523	5338
- 1/114FC	. =			2233

	127		
1 2	3 4	5	6
DIRINECY279MC	808648432140	58301	5358
DIRINECY280MC	809150431941	58106	5373
DIRINECY281MC	809603431742	58031	5373
DIRINECY282MC	810006431558	58068	5372
D1RINECY283MC	810342431427	58186	5366
DIRINECY289MC	804993448556	58411	5635
D1RINECY290MC	804964448068	58 <b>354</b>	5638
DIRINECY291MC	804980447593	58442	5628
DIRINECY292MC	804986447151	58340	5631
D1RINECY293MC	804992446694	58330	5624
DIRINECY294MC	805117445907	58383	5621
D1RINECY295MC	805148445438	58306	5628
DIRINECY296MC	805154445003	58300	5632
DIRINECY297MC	805285444560	58247	5634
DIRINECY298MC	805331444065	58301	5622
MARINECY299MC	805600443250	58233	5613
DIRINECY300MC	805618443049	58316	5610
DIRINECY301MC	805605442624	58212	5600
DIRINECY302MC	807406448712	58504	5634
D1RINECY303MC	807412448247	58515	5624
DIRINECY304MC	807428447762	58491	5626
DIRINECY305MC	807475447263	58511	5618
DIRINECY306MC	807481446788	58420	5617
DIRINECY307MC	807442445938	58438	5607
DIRINECY308MC	807459445383	58374	5609
DIRINECY309MC	807471444838	58478	5598
DIRINECY310MC	807578444320	58593	5585 5540
DIRINECY311MC DIRINECY312MC	807596443765	58745	5568 5566
DIRINECY312MC	807628443228	58850	5556 5543
DIRINECY314MC	807660442679 808831442956	58 <b>92</b> 5 58466	5543 5569
DIRINECY314MC	808749443518	58492	5576
DIRINECY316MC	808742444030	58486	5583
DIRINECY317MC	808735444593	58487	5583
DIRINECY318MC	808753445093	58483	5591
DIRINECY319MC	808771445625	58451	5596
DIRINECY320MC	808739446175	58443	5605
DIRINECY322MC	808227445183	58485	5601
DIRINECY323MC	807728445142	58448	5607
DIRINECY324MC	807229445060	58360	5611
DIRINECY325MC	806679445003	58261	5622
DIRINECY326MC	806205444947	58417	5607
DIRINECY327MC	805680444925	58491	5617
DIRINECY328MC	808194443868	58584	5580
	0.700.4400.0		

807080443808 58657

806500443776 58445

806000443850 58332

805471443687 58318

804896443680 58342

804372443648 58218

803847443651 58264

803626438763 58362

5574

5597

5599

5612

5611

5616

5610

5529

DIRINECY330MC

DIRINECY331MC

MARINECY332MC

DIRINECY333MC

DIRINECY334MC

DIRINECY335MC

DIRINECY336MC

DIRINECY337MC

3

1

2

4

5

6

1	2		3 4	5、	6	. 7
DIRI	NECY408MC	799	5014585	73 6125	2 5612	· ———
DIRI	NECY409MC	799	4954590	63 6131	1 5624	
	3R I V5 1 5BR	810	2274604	92 5928	5 <b>5817</b>	
M28B	3RIV519BR	804	8554602	11 5981	1 5727	
M286	B MC 3061	794	9154421	77 6035	3 5444	
M288	B MC 3066	795	0774394	04 6042	B <b>5410</b>	
ST CL	AIR05531	794	10045969	5006208	3 (	005543
	AIR05534	794	1604582	3506194	0 (	005521
	AIR05537	794	2404568	0006184	в (	005500
ST CL	-AIR05540	794	2854554	0006184	0 (	005480
ST CL	AIR05543	794	35045429	5006173	9 (	005466
ST CL	AIR05546	794	39545279	5506141	5 (	005 <b>470</b>
ST CL	AIR05549	794	4504512	1506119	в (	005460
ST CL	A I R 05552	794	5104496	3006087	4 (	005458
	AIR05554	794	54544859	5506085	4 (	005447
	AIR05557	794	6504470	5006072	0 (	005452
	AIR05562	794	7954449	0006055	5 (	005436
	_AIR05578	795	4904597	4006204	5 (	005545
	A1R05581	796	9404598	1506186	7 (	005557
ST CL	_AIR05584	798	41045989	9506165	7 (	005582
	_AIR05586	799	4304599	5006158	9 (	005575
	_AIR05589	800	8854600	1506132	5 (	005 <b>587</b>
	AIR05592	802	3654600	9506110	1 (	005596
	AIR05595	803	8254601	7506012	6 (	005654
	_AIR05600	806	3304603	0005982	1 (	005675
	_AIR05603	807	8154603	9505908	4 (	005746
	_AIR05606	809	4104604	5505905	3 (	005775
	_AIR05739	795	7754543	2506154	0 (	005473
	AIR05742	797	1004544	0006151	0 (	005474
	_AIR05745		4004544			005497
	_AIR05748	799	3004545	2506121	7 (	005488
	_AIR05751		1504546			005506
	_AIR05754	802	4504547	0006022	4 (	005499
	_AIR05758	804	2004547	7505976	1	005558
	AIR05762		0004548			005612
	-AIR05765		4504548			005625
	AIR05768	808	9004549	5005896	6 (	005636
	_AIR05827		6504486			005443
ST C	AIR05830	797	0504487	0006080	3 (	005444

## BELLE RIVER MILLS SURVEY

1	2	3	3	4	5	6
M28BBR I	VBASEK	8098	34246	55826	59100	5910
M28BBRI				70709	61924	5841
M28BBRI				57546	61483	5812
M28BBRI				70952	60856	5922
M28BERI		-	_	76422	59457	6143
M28BBRI				76476	61329	6024
M28BBRI				57732	60617	5865
M28BBRI				71196	59797	5965
M288BRI	· - · · <del>-</del> - ·			5608	60422	5826
M28BBRI	_			76057	62105	5944
M28BBRI				76079	62054	5951
M28BBRI				76095	62004	5962
M28BBRI				76117	61897	5971
M2888R I				76138	61554	5994
M288BR I				76164	61196	
M288BR I						6018
	-			76185	61097	6027
M28BBRI				76233	61422	6007
M28BBRI				76255	61361	6016
M28BBRI				76281	61363	6015
M28BBRI				76308	61350	6018
M28BBRI				76329	61079	6038
M28BBRI				76345	60894	6054
M28BBRI				76371	60927	6055
M28BBRI				76393	60834	6064
M28BBRI				76414	60647	6075
M28BBRI				76440	60522	6081
M28BBRI				76448	60488	6087
M28BBR I				76469	60449	6090
M28BBRI				76495	59876	6121
M28BBRI				76532	59449	6150
M28BBRI				76073	59613	6121
M28BBRI		•		75578	59600	6105
M26BBRI	V026BR	8099	56147	75093	59497	6093
M28BBRI	V027BR	8095	57747	74623	59476	6079
M28BBRI	V028BR	8099	59847	74159	59530	6064
M28BBRI	V029BR	8096	51447	73674	59531	6046
M28BBRI	V030BR	8096	55047	73400	59520	6032
M28BBRI	V031BR	8096	66847	72649	59699	6008
M28BBRI	V032BR	8097	72547	72500	59953	5979
M28BBRI	V033BR	8097	70347	71835	59953	5968
M28BBRI	V034BR	8097	73447	71400	59928	5959
M28BBRI	V035BR	8097	73747	71200	59994	5949
M28BBRI	V036BR	8092	28747	71184	59677	5978
M28BERI	V037BR	8088	33747	71163	59581	5986
M28BBRI	V038BR	8083	38747	71147	58967	6029
M28BBRI	V039BR	8079	3847	71127	58960	6034
M28BBRI	V040BR	8074	8847	71106	59060	6033
M28BBRI	V041BR	8070	6347	71090	59092	6036
M28BBRI	V042BR	8066	51347	71069	59016	6044

1 2 3 4 5 6

			132			
1	2		3	4	5	6
M28BBRI	V097BR	803	317246	57714	60685	5862
M28BBRI			264246		60943	5841
M28BBRI			208346	-	61006	5837
M28BBRI			156946		61116	5829
M28BBR			21046		61356	5820
M28BBRI			013546		60779	5855
M28BBRI			944146		61642	5801
M28BBRI			910246		61847	5789
M28BBRI			906046		61872	5799
M28BBRI			903846	_	61880	5811
M28BBRI			900646		61908	5820
M28BBRI			39744		61947	5826
M28BBRI			3494		61596	5852
M28BBRI			038146		61700	5830
M28BBRI			041346		61558	5832
M28BBRI			043546		61532	5826
M28BBR			045246		61421	5826
M28BBR			051046	-	61485	5810
M28BBRI			45046		60887	5829
M28BBRI			72546		61106	5799
M28BBRI			225146		61085	5798
M28BBRI			280446		60805	5812
M28BBRI			339246		60643	5818
M28BBR			388446		60428	5829
M28BBRI			441046		60386	5829
M28BBRI			495946		60361	5828
M28BBRI			550946		59752	5862
M28BBR	V125BR	808	504446	55666	59674	5856
M28BBRI	V126BR	806	560946	55689	59454	5866
M28BBRI	V127BR	80	715846	55716	59361	5871
M28BBRI	V128BR	80	776846	55744	59234	5888
M28BBRI	V129BR	808	333346	55761	59075	5914
M28BBRI	V130BR	808	390846	55789	58977	5917
M28BBRI	V131BR	809	945746	55816	58736	5933
M28BBRI	V132BR	809	998246	55833	59247	5896
M28BBRI	V133BR	809	99674	76559	59421	6147
M28BBRI	V134BR	810	04764	76580	59440	6145
M28BBRI	V135BR	810	9714	76597	59441	6152
M2EBBR I			4424		59108	6172
M28BBRI			19034		59442	6144
M28BBRI			22404		59558	6158
M28BBRI			25444		59401	6188
M28BBRI			30604		59651	6174
M28BBRI			35354		59472	6189
M28BBRI			10124	-	59432	6185
M28BBRI			98254		59841	5951
M28BBRI			98504		59780	5941
M28BBRI			980646	-	59676	5932
M28BBRI			983846		59638	5929
M28BERI			985946		59575	5924
M28BBRI	A148BK	804	988646	00222	59510	5933

		133			
1	2	3	4	5	6
M28BBRI	V149BR	809907	467727	59406	5933
M28BBRI		809929		59407	5922
M28BBRI		809945		59305	5920
M28BBRI		809962		59261	5906
M28BBRI		810467		59011	5910
M28BBRI		810947		58997	5918
M28BBRI		811441		58924	5936
M28BBRI		811906	_	58832	5946
M28BBRI		812406		58743	5964
M28BBRI		812880		58690	5964
M28BBRI		813380		58643	5958
M28BBRI		813880		58599	5964
M28BBRI	V161BR	814430	466041	58520	5973
M28BBRI	V162BR	814979	466078	58542	5977
M28BBRI	V163BR	815479	466100	58614	5982
M28BBRI	V164BR	810261	471227	59770	5961
M28BBRI	V166BR	811286	471270	59847	5957
M28BBRI	V167BR	811796	471292	59270	6014
M28BBRI	V168BR	812336	471314	59107	6025
M28BBRI	V169BR	812825	471336	59139	6037
M28BBRI	V170BR	813335	471352	59158	6039
M28BBRI	V171BR	813875	471374	59171	6039
M28BBRI	V172BR	814395	471396	59149	6051
M28BBRI	V173BR	814934	471428	59251	6048
M28BBRI		815409	471439	59251	6057
M28BBRI	V175BR	809470	477032	59543	6155
M28BBRI		809454		59780	6164
M28BBRI		809432	478032	60030	6164
M28BBRI		809416		60226	6168
M28BBRI		809394		60359	6178
M28BBRI		809368		60427	6194
M28BBRI		809315	-	60416	<b>6</b> 222
M28BBRI		809298		60319	6240
M28BBRI		809277		60220	6269
M28BBRI		809271		60359	6274
M28BBRI		804014		61456	6021
M28BBRI		803976		61317	6059
M28BBRI		803944		61330	6076
M28BBRI		803923	_	61242	6103
M28BBRI		803906		61185	6122
M28BBRI		803875		60996	6152
M28BBRI		803847		61429	6152
M28BBRI		8038154		61347	6171
M28BBRI		8038004 7986414		61493	6174
M28BBRI M28BBRI		798641		62058 62115	5962 5970
M288BRI		798588		62024	597 <b>0</b> 5986
M28BBRI		798572	_	62099	599 <b>5</b>
M288BRI		795525		62111	6000
M28BBRI		798373		62200	5939
M28BBRI		797648		62220	5936
M28BBRI		797173		62250	5935
<del></del>					

<b>_</b>						
1	2	3	4	5	6	

M28BBRIV207BR	796159475923	62439	5921
M28BBRIV208BR	798443470662	61819	5847
M28BBRIV209BR	797843470629	61885	5839
M28BBRIV210BR	797453470609	61986	5835
M28BBRIV211BR	796969470583	62046	5833
M28BBRIV212BR	796454470551	62114	5830
M28BBRIV213BR	795979470525	62276	5826
M28BBRIV214BR	795495470504	62292	5821
M28BBRIV215BR	795035470483	62327	5820
M28BBRIV216BR	794520470451	62247	5822
M28BBRIV217BR	794021470419	62136	5833
M28BBRIV218BR	798579468694	61735	5808
M28BBRIV219BR	798118468728	61647	5811
M28BBRIV220BR	797644468652	61753	5804
M28BBRIV221BR	797165468560	61885	5796
M28BBRIV222BR	796670468539	61905	5794
M28BBRIV223BR	796196468518	61960	5793
M28BBRIV224BR	795696468496	62004	5792
M28BBRIV226BR	794706468453	61531	5822
M28BBRIV227BR	794207468427	61442	5832
M28BBRIV229BR	804588466122	60489	5833
M28BBRIV230BR	804571466662	60481	5849
M28BBRIV231BR	804539467162	60494	5856
M28BBRIV232BR	799998467222	59860	5904
M28BBRIV233BR	799991466962	59891	5893
M28BBRIV234BR	800029466773	60071	5875
M28BBRIV235BR	800107466544	60131	5870
M28BBRIV236BR	800125466309	60527	5836
M28BBRIV237BR	799953466102	61265	5782
M28BBRIV238BR	799632465742	61643	5750
M28BBRIV239BR	799922465371	61409	5755
M28BBRIV240BR	800212465035	61311	5750
M28BBRIV241BR	800670464756	61293	5745
M28BBRIV242BR	801146464697	61341	5745
M28BBRIV243BR	801596464678	60758	5789
M28BBRIV244BR	802024464489	60907	5774
M28BBRIV245BR	802428464204	61077	5757
M28BBRIV246BR	802831463919	60844	5769
M28BBRIV247BR	803161463548	60597	5775
M28BBRIV248BR	803476463203	60393	5776
M28BBRIV249BR	803884463003	60005	5788
M28BBRIV250BR	804368463029	60085	5784
M28BBRIV251BR	799213465312	61713	5732
M28BBR I V252BR	799229464837	61653	5724
M28BBRIV254BR	799282463868	61524	5720
M28BBRIV256BR	799329462933	61540	5693
M28BBRIV258BR	798214465259	61722	5728
M28BBR I V259BR	797709465217	61726	5729
M28BBRIV260BR	797214465196	61809	5728
M28BBRIV261BR	796715465164	61863	5722
M28BBRIV262BR	799229465987	61513	5776
M28BBRIV263BR	798777466156	61578	5776

		135			
1	2	3	4	5	6
	1V264BR		0466311	61504	5773
	V265BR		3466447	61414	5780
	V266BR		2466554	61680	5762
M28BBR	V267BR	796955	5466633	61960	5744
	1 V 2 6 8 B R	799274	4474013	61949	5914
M28BBR	(V269BR	799748	3474070	61896	5919
M26BBR	(V270BR	800202	2473811	61722	5928
	V272BR	801156	5473488	61532	<b>5930</b>
	(V273BR		473495	61054	5962
	1V274BR		5473576	60996	5973
	V275BR		4473633	60744	5992
	(V276BR		4473689	60771	5993
	(V277BR		3473745	60960	5988
	I V278BR		7473796	61038	5981
	I V279BR I V280BR		4465333 5464804	59124 59086	5896 5892
	1 V 2 8 1 B R		2464304	59140	5884
	1 V 2 8 2 B R		9463809	59156	5878
	V283BR		2463164	59211	5863
	V284BR		2474959	60529	6050
	V285BR		7474977	60622	6042
	V286BR		7474971	60722	6032
	V287BR		7474949	61090	6007
	V288BR		7474932	61064	6005
M28BBR	1 V289BR	804576	5473883	61116	5974
M28BBR	V290BR	805101	1473915	60688	6001
M28BBR	(V291BR	805535	5473945	60491	6009
M28BBR	V292BR	806050	473977	60306	6033
	(V293BR		5474007	60035	6057
	(V294BR		0472064	61842	5878
	(V295BR		4472121	62007	5869
	V296BR		4472102	61647	5895
	(V297BR		5472357	61718	5891
	V298BR		5472364	61418	5908
	1 V299BR		5472396	60954	5938
	IV300BR IV301BR		5472342 7472294	60859 60568	5945 5959
	V301BR IV302BR		3472204	60615	5965
	V302BR		3472211	60871	5949
	1V304BR		3472202	60852	5947
	V306BR		3472241	60621	5970
	V307BR		3472237	60669	5966
	1V308BR		3472229	60305	5986
M28BBR	1V309BR	803019	9471387	60488	5953
M28BBR	V310BR	803027	7471887	60516	5955
M2EBBR	(V311BR	802743	3472609	60670	5973
M28BBR	V312BR	802747	7473084	60667	5988
	(V313BR		3475667	61309	6003
	(V314BR		4475168	61134	6006
	(V316BR		2474168	60954	5989
	V317BR		9473644	61056	5962
M28BBR	(V318BR	801536	5473128	60992	5955

		136			
1	2	3	4	5	6
M28BBR I	V3198D	80155247	2680	60920	5949
M28BBR		80162447		61243	5910
M28BBR		80172547		61272	5902
M28BBR		80168747	_	61236	5887
M28BBR		80919347		59335	6036
M28BBR		80866947		59143	6055
M28BBR		80816847		59143	6060
M28BER		80767047		59057	6069
M28BBR		80717147		59004	6072
M28BBRI		80672247		59093	6063
M28BBR	V331BR	80827747		59059	6118
M28BBR	V332BR	80824947	4106	59014	6113
M28BBR	V333BR	80772347	4149	59031	6119
M28BBR1	V334BR	80720847	4167	59259	6100
M28BBR	V335BR	80685647		60320	6039
M28BBR	V336BR	79953646	9297	61777	5821
M28BBR	V337BR	80008646	9324	61617	5837
M28BBR	V338BR	80085446	9469	61338	5851
M28BBRI	V339BR	80133346	9490	61036	5874
M28BBRI	V340BR	80180846	9531	60902	5887
M28BBR	V341BR	80229346	9528	60845	5888
M28BBR1	V342BR	80278346	9559	60972	5882
M28BBR	V343BR	80338246	9617	60292	5932
M28BBRI	V344BR	80385746	9648	60692	5909
M28BBR1	V345BR	80433146	9679	60760	5907
M28BBRI	V346BR	80481646	9661	60481	5924
M28BBR1	V347BR	80529646	9692	60129	5947
M28BBR		80174747	0346	60929	5899
M28BBR		80174346	_	60733	5901
M28BBR		80175046		60752	5892
M28BBR1		80176646		60666	5890
M28BBR1		80175446		60876	5855
M28BBR		80314747		60267	5947
M28BBR		80325346		60392	5928
M28BBR		80336046		60124	5937
M2EBBR1		80341646		60052	5936
M28BBRI		80350746		60180	5919
M28BBR1		80357746		60630	5872
M28BBR		80445046		60389	5897
M28BBRI		80441846		60606	5894
M28BBR 1		80439746		60631	5904
M28BBRI M28BBRI		80435047 80434347		60804 60732	5915 5928
M28BBR 1		80625646		59020	6010
M28BBR 1		80679046		59005	6008
M28BBR 1		80725746		58935	6010
M28BERI		80775546		58761	6018
M28BBR1		80824046		58799	6012
M28BBR 1		80876546		59114	5989
M28BBR1		80845747		58928	6039
M288BR1		80842547		59027	6052
M28BBR1		80839347		59201	6058
		-5555541		•	-000

	- <b>J</b> ·	
1	345	6 7
M28BBRIV375BR	808372473057 59196	6064
M28BBRIV377BR	805925468275 59658	5931
M28BBRIV378BR	806424468307 59083	5959
M28BBRIV379BR	806914468338 59072	5955
M28BBRIV380BR	807398468404 59076	5956
M28BBRIV381BR	807843468775 58740	5991
M28BBR!V382BR	808068468788 58769	5992
M28BBRIV383BR	808718468837 58914	5988
M28BBRIV384BR	809175468950 59250	5955
M288BRIV385BR	809668468799 59476	5936
M28BBR I V386BR	807095470591 58993	6029
M28BBRIV387BR	807101470091 59007	6019
M28BBRIV389BR	807188469202 58872	5996
M28BBRIV390BR	807354468744 58940	5972
M28BBRIV392BR	807529467946 59345	5923
M28BBRIV393BR	807686467448 59213	5926
M28BBRIV394BR	807439467195 59025	5926
M28BBRIV395BR	807469466845 58970	5922
M28BBRIV396BR	807524466396 59152	5901
M28BBRIV411BR	799452460416 61465	5648
M28BBRIV412BR	799421460899 61415	5669
M28BBRIV413BR	799404461389 61459	5679
M28BBRIV414BR	799383461864 61526	5685
M28BBRIV415BR	799352462364 61582	5689
M28BBRIV500BR	795199475885 62447	5925
M28BBRIV501BR	794250475848 62485	5923
M28BBRIV502BR	798502479479 62128	6045
M28BBRIV503BR	798468480178 62261	6058
728BBRIV510BR	795750465100 62078	5723
M28BBRIV511BR	794816465054 62154	5701
M28BBRIV513BR	810150462205 59278	5846
M28BBRIV514BR	810192461220 59361	5830
M28BBRIV516BR	804745462919 60034	5790
M28BBRIV517BR	804782461985 59814	5777
M28BBRIV518BR		
		5760 5814
M28BBRIV521BR	806694463010 59395	5814
M28BBRIV522BR	807693463058 59321 808658463100 58819	5833 5875
M28BBRIV523BR	811640482500060695	5875
ST CLAIR05288		006239
ST CLAIR05291	810630482050060144	006254
ST CLAIR05355	808220481920060938	006181
ST CLAIR05358	806735481845061208	006148
ST CLAIR05361	805255481720061597	006119
ST CLAIR05366	802770481615061596	006113
ST CLAIR05369	801355481515061808	006085
ST CLAIR05372	800000481440062262	006054
ST CLAIR05374	799050481370062465	006035
ST CLAIR05378	797385481340062508	006030
ST CLAIR05381	795945481260062687	006017
ST CLAIR05386	793555481153063443	005978
ST CLAIR05473	792940480100063275	005958
ST CLAIR05476	793005478780062925	005932

	13	8			
1 2	3	4	5	6	7
ST CLAIR05479	793100	477330	062615		005910
ST CLAIR05482		475785			005869
ST CLAIR05485	793260	474320	062145		005868
ST CLAIR05488	793325	472940	062385		005818
ST CLAIR05491	793420	471480	062051		005809
ST CLAIR05493	793480	470395	060368		005896
ST CLAIR05495	793530	469370	062251		005753
ST CLAIR05497	793580	468430	061437		005783
ST CLAIR05500	793725	466390	062287		005677
ST CLAIR05520	793800	465025	062220		005647
ST CLAIR05522	793870	464045	062136		005627
ST CLAIR05526	793955	462085	062185		005 <b>57</b> 9
ST CLAIR05529	794045	460670	062092		005560
M28BBRIV515BR	810227	7460492	59285	581	7
M28BBRIV519BR	804855	5460211	59811	572	7
ST CLAIR05531		459650			005543
ST CLAIR05578	795490	459740	062045		005 <b>545</b>
ST CLAIR05581	_	459815	•		005 <b>557</b>
ST CLAIR05584	798410	459895	061657		005582
ST CLAIR05586	799430	459950	061589		005575
ST CLAIR05589		460015	-		005587
ST CLAIR05592	802365	460095	061101		005596
ST CLAIR05595		5460175	-		0056 <b>54</b>
ST CLAIR05600		460300			005675
ST CLAIR05603		460395			005746
ST CLAIR05606	809410	460465	059053		005775

## BERLIN SURVEY

1	2		3	4	5	6
OOBERL	IN00001	693	30951	1545	81222	4496
	100002		28751		81675	
	1N00003		27051		81730	
	IN00004		24851		81739	
	100005		23151		81701	4536
	100006		21551		81700	4558
	IN00007		69051		81239	
	80000N		20051		81166	
	N00009		68551		81174	4577
	01000N		16151		80629	
OOBERL			56651		80083	
	N00012		58351		79874	4622
	IN00013		59951		79552	4624
	N00014		61651		79276	
	IN00015		62351		79289	
	N00016		64551		79261	4596
	IN00017		16051		79167	
	N00018		67051		79429	
	N00020		69951		80446	
OOBERL			80451		81456	
	100022		31951		81586	
	N00023		79851		81511	4481
	N00024		31851	_	81380	4493
	N00025		82851		81352	
	N00026		28851		81256	
	N00027		81751		81104	
	1N00028		26751		81034	4522
	N00029		25551	_	81072	
OOBERL			22751		81056	
	N00032		21051		80907	
	1N00033		18251		81116	
	N00034		70351		81249	
	N00036		65851		81321	4583
	N00037		19351		81097	
	N00038		65951		81074	4593
	100039		19451		81260	4583
	IN00040		68451		81466	
OOBERL			33551		80445	_
	N00042		34151		79962	
	IN00043		37650		79534	4541
	N00044		37650		79403	
	N00045		39350		79155	
	N00046		20950		78725	
	IN00048		23850		78578	
	IN00049		25550		78504	4486
	N00050		26750		78484	4466
	N00051		28350		78436	
	N00052		76850		78309	
	N00053		24350	_	78239	
	N00054		71850		78340	
	_				_	

		140			
1	2	3	4	5	6
OOBERLIN	100055	691158	505856	78613	4450
OOBERLIN		690878	505847	78561	4453
OOBERLIN	NO0057	690857	506317	78667	4463
OOBERLIN	100058	690830	506842	78660	4481
OOBERLIN	100059	690794	507341	78630	4498
OOBERLIN	100060	690752	507846	78625	452 <b>6</b>
OOBERLIN	N00061	690720	508366	78596	4546
OOBERLIN		690705		78758	4559
OOBERL IN		690698		78947	4564
OOBERL IN		690681		79089	4573
OOBERLIN		690675		79238	4585
OOBERLIN		690658		79246	4590
OOBERLIN	-	693767	_	78626	4440
OOEERLIN		694267		79019	4417
OOBERLIN		694792		79168	4408
OOBERLIN		695292		79192	4406
OOBERLIN		695802		79313	4403
OOBERLIN		696312	-	79399	4401
OOBERLIN		696792		79400	4416
OOBERLIN		697291		79381	4421
OOBERLIN		697806		79362	4437
OOBERLIN		699341	-	78886	4476
00BERLIN		699941 700480		79598 79937	4436
OOBERLIN		701015		79804	4421 4433
OOBERLIN		701015		79714	4447
OOBERLIN		702089		79556	4466
OOBERLIN		702589		79459	4482
OOBERLIN		703164		79375	4498
OOBERLIN		703689		79443	4499
OOBERLIN		704264		79384	4513
OOBERLIN		704824		79193	4546
OOBERLIN		690554		80370	4601
OOBERLIN		690532	515089	80785	4597
OOBERLIN		690506	515603	81344	4581
OOBERLIN	100092	690483	516153	81312	4611
OOBERLIN	100093	690462	516683	81613	4616
OOBERLIN	100094	690445	517213	81859	4617
OOBERLIN	100095	690423	517747	81990	4626
OOBERLIN		690401		82293	4625
OOBERLIN		690384		81944	4664
OOBERL IN		690351		81972	4678
OOBERLIN		690344		81991	4705
OOBERLIN	-	690337		81679	4745
OOBERLIN		690325		81339	4785
OOBERLIN		690313		81223	4818
OOBERLIN		690302		80753	4878
OOBERLIN		690295		80527	4917
OOBERLIN		690283		80563	4932
OOBERLIN		690266		80305	4962
OOBERLIN		690254 690237		80212 80274	4979
OOBERL IN	400 t 00	070E3 F	JE4000	302 /4	4999

		Τ,	† <u>T</u>			
1	2	3		4	5	6
OOBERLI	N00109	69320	0251	4719	81700	4572
OOBERL I	N00110	69318	3151	5233	81654	4591
OOBERL I	N00111	69316	5451	5763	81730	4609
OOBERL I	N00112	69314	751	6283	81849	4616
OOBERL I	N00113	69312	2551	6803	81970	4621
00BERL I	N00114	6931	351	<b>7</b> 323	81943	4642
OOBERL I	N00115	69308	3751	7847	81682	4683
OOBERL I	N00116	69306	5551	8372	81320	4721
OOBERL I	N00117	69304	4851	8882	81055	4756
OOBERLI		6930	1451	9597	81252	4769
OOBERLI		6930	1351	9997	81043	4802
OOBERL I		69299	9252	0521	80631	4854
OOBERL I		69298	3052	1046	80629	4873
OOBERL I		6929	5652	2071	80512	4910
OOBERLI		69293	3952	2606	80346	4947
OOBERLI	N00125	69292	2852	3116	80183	4981
00BERL I	N00126	6929	1152	3626	80102	5004
OOBERL I	N00127	69289	9452	4130	79886	5041
OOBERL I	N00128	6928	7552	4805	79755	5077
OOBERL I	N00129	69716	5551	4915	81018	4619
OOBERL I	N00130	69714	4851	5435	80830	4650
OOBERL I	N00131	69713	3251	5905	80628	4680
OOBERL I	N00132	6971	1651	6410	80455	4717
OOEERL I	N00133	69709	9451	6925	80180	4754
OOBERL I	N00134	6970	7751	7440	79894	4789
OOBERLI	N00135	69709	5651	7949	79736	4820
OOBERL I	N00136	69703	3951	8424	79998	4821
OOBERLI	N00137	69702	2351	8899	79984	4852
OOBERL I	N00139	69699	9351	9684	79651	4900
OOBERL I		6977(			80608	4641
OOBERL I		6982			80024	
OOBERL I		6987			79691	4709
OOBERLI		6992	1251	4447	79620	
OOBERL I		6997			79619	
OOBERLI	N00145	7002			79927	4709
OOBERLI		70074			80039	
OOBERL I		70129			80020	
OOBERLI	-	70178			80003	
OOBERL I	N00149	7023	0051	4558	79967	4703
OOBERL I		70283			79977	-
OOBERL I		7038	7051	4613	79851	4741
OOBERL I	N00153	70438			79804	
OOBERL I		70489			79657	
OOBERL I	- <del>-</del>	70533	3951	4657	79464	
OOBERL I		70582			79302	
OOBERL I		70632			78886	
OOBERLI		6975			79646	
OOBERL I		6980			79559	_
OOBERL I		69839			79417	
OOBERLI		6983			79332	
OOBERLI		69833			79262	
00BERL I	N00163	6983	1852	1176	79282	5014

		142			
ı	2	3	4	5	6
OOBERI	IN00164	698297	521671	79305	5036
	IN00166	698244		79333	5084
	1N00167	698218		79532	5091
	IN00168	698191		79396	5121
	IN00169	698170		79238	5151
	IN00170	698148		79256	5171
	IN00171	697727		80566	4559
	IN00172	698207		80100	4589
	IN00173	698697		79902	4603
	IN00174	699186	511792	79885	4603
OOBERL	IN00175	699606	511807	80016	4595
OOBERL	IN00176	699822	511335	80063	4570
OOEERL	IN00177	699988	510867	80410	4528
OOBERL	IN00178	700155	510379	80064	4537
OOBERL	IN00179	700291	509916	80197	4520
00BERL	IN00180	700372	509442	80137	4510
OOBERL	IN00181	700366	509167	80098	4505
00BERL	IN00183	701375	509195	79787	4532
OOBERL	IN00184	701875	509212	79549	4554
OOBERL	IN00185	702425	509224	79504	4564
OOBERL	IN00186	702875	509230	79208	4584
OOBERL	IN00187	703385		79208	4592
OOBERL	IN00188	703900	509253	79027	4612
	IN00189	704425		79086	4622
	IN00190	704930		79135	4621
	IN00191	705435		78991	4639
	IN00192	705935		78800	4654
	IN00194	706924		78068	4700
	IN00195	707424		78087	4704
	IN00196	690036		79758	4639
	IN00198	689027		79081	4722
	IN00199	688637		79063	4727
	IN00200	687987		79540	4693
	IN00201	687487		79091	4724
	1N00202	686958		78950	4734
	IN00203	686458		78962	4737
	IN00204	685968		78995	4733
	IN00205 IN00206	685468 684964		78972 79255	4738
		684464		79172	4718
	IN00207 IN00208	683949		79098	4724 4741
	IN00208	683439		79283	4733
	IN00209	682929		79055	4745
	IN00210	682539		79097	4759
	IN00212	682039		79383	4753
	IN00213	681564		79587	4759
	IN00217	690195		78655	4565
	IN00218	689685		78551	4570
	IN00219	689181		78577	4570
	IN00221	688166		79000	4549
	IN00222	687666		78932	4555
	IN00223	687146		78868	4562

		14)	3		
1	2	3	4	5	6
OOBERLI	N00224	68663	7508612	78892	4567
OOBERL I			7508585	78976	4559
OOBERLI			2508556	79150	4554
OOBERLI			3508536	79172	4568
OOBERL I			8508519	79062	4583
OOBERL I		68403	3508503	78848	4601
OOBERLI		68351	8508481	78595	4616
OOBERL I	N00231	68301	8508460	78585	4621
OOBERL I	N00232	68250	4508448	78505	4627
OOBERL I	N00233	69058	8505474	78455	4449
OOBERLI	N00234	69058	4505053	78474	4433
OOBERL I	N00235		5504543	78519	4415
OOBERL I			2504437	78528	4412
OOBERL I			2504425	78566	4414
OOBERL I		-	2504404	78613	4410
OOBERL I			2504387	78600	4425
OOBERL I			4504220	78572	4430
OOBERLI			9503849	78624	4408
OOBERLI			6503339	78533	4401
OOBERLI			7502854	78472	4391
00BERL I			9502359	78506	4375
OOBERLI			5501859	78529	4352
OOBERLI			2501360	78587	4325
OOBERL I			9500830	78550	4292
OOBERL I			3505581	79259	4435
OOBERL I			9505061	79140	4427
OOBERL I			1504547	78904	4427
OOBERL I			3504012 6503397	78695 78265	4423
00BERL I			8502912	78171	4430 4420
OOBERL I			4502412	78001	4420
OOBERL I			6501913	78102	4397
OOBERL I			7501403	78188	4372
OOBERL I			4500888	77576	4397
00BERL I			6500378	78169	4336
OOBERL I			4499389	78260	4295
00BERL I			5498889	78095	4286
OOBERLI			9505414	78412	4434
OOBERLI			7502901	78271	4388
OOBERLI			8502426	78404	4361
OOBERL I			2501411	78647	4309
OOBERL I		69350	8500901	78532	4297
00BERL I	N00274	69352	0500402	78524	4283
OOBERL I	N00275	69354	6499902	78395	4275
OOBERL I	N00276	69357	8499382	78415	4255
OOBERL I	N00277	69359	5498877	78463	4236
OOBERL I		69361	6498403	78405	4221
OOBERL I			1511377	79167	4591
OOBERL I			6511345	78814	4602
OOBERL I			5506526	79376	4435
OOBERLI			7507525	79298	4481
OOBERL I	N00286	69721	1507985	79283	4500

		144			
1	2	3	4	5	6
OOBERL I	N00287	6972155	508471	79462	4507
OOBERL I		6972139		79603	4513
OOBERL I	N00289	6971305	509610	79797	4525
OOBERL I	N00290	6970889	510089	79980	4532
OOBERL 1	N00291	6974565	510244	80032	4539
OOBERL I		6973135	511242	80479	4543
OOBERL I		690902		81505	4625
OOBERL I		6914225		81930	4598
OOBERL I		6923815		81699	4629
OOBERL I		692875		81697	4635
OOBERL I		6938789		79043	4542
00BERL I		6943289		79122	4544
OOBERL I		6947879		79152	4542
OOBERL I		6952525		79364	4531
OOBERL I		6957279		79414	4528
OOBERL I		6961879 6967029		79463 79471	4524 4523
00EERLI 00BERLI		6929295	-	78796	452 <i>3</i> 4552
OOBERL I		6924545		78458	4580
OOBERL I		6920045		78551	4577
OOBERL I		6915295		78723	4565
OOBERL I		6952355		81005	4582
OOBERL I		6952565		81817	4514
OOBERLI		6952535		81926	4498
OOBERL I		6952795	_	81807	4495
OOBERL I		6951339		81692	4491
OOBERL I		6951405	511099	81237	4483
OOBERLI	N00325	6952665	510625	80255	453 <b>5</b>
OOBERL I	N00326	6952885	510135	79809	4545
OOBERL I	N00327	6953299	509661	79643	4542
BERLIN	00328	6878189		79350	4690
BERLIN	00329	6878345		79039	4692
BERLIN	00330	6878505	512588	78856	4694
BERLIN	00331	6880509		78800	4567
BERLIN	00332	6880469		78791	4566
BERLIN	00334	6876879		78841	4540
BERLIN	00335	6877129		78917	4530
BERLIN	00336	6877245	_	78891	4514
BERLIN	00337	6877399		78674	4515
BERLIN	00338	6878339		78645	4442
BERLIN	00339	6878129		78638	4457
BERLIN	00340	6877969		78568	4474
BERLIN	00341 N00342	6877669 6933619		78541 79693	4492 4544
00BERL I	1400342	0933013	110220	1 3033	4344

MICHIGAN STATE UNIV. LIBRARIES
31293101993636