



This is to certify that the

dissertation entitled

COMPARATIVE ADAPTATIONS TO FOLIVORY IN PRIMATES

presented by

Mark Stephen Edwards

has been accepted towards fulfillment of the requirements for

Ph.D. degree in Animal Science

Suam & Willsey
Major professor

Date Argus 22,1995

LIBRARY Michigan State University

PLACE IN RETURN BOX to remove this checkout from your record.
TO AVOID FINES return on or before date due.

DATE DUE	DATE DUE	DATE DUE

MSU Is An Affirmative Action/Equal Opportunity Institution

COMPARATIVE ADAPTATIONS TO FOLIVORY IN PRIMATES

Volume I

second focuses on digestion By salities of saveral captive

Dally San Mark Stephen Edwards

A DISSERTATION

Submitted to
Michigan State University
for the degree of

DOCTOR OF PHILOSOPHY

Department of Animal Science

ABSTRACT

COMPARATIVE ADAPTATIONS TO FOLIVORY IN PRIMATES

Ву

Mark Stephen Edwards

This study is divided into two major sections. The first examines feeding habits and nutrient composition of natural diets of free-ranging red howlers (Alouatta seniculus). The second focuses on digestive capabilities of several captive primate species.

Daily activity, feeding behavior, food items selected, and number of bites of food were recorded during monthly observations over a 12-month period of two groups of free-ranging red howlers. After 5 months of observation, one group was transferred to an on-site holding facility. These animals were used as models to determine weights of an individual bite for specific plant parts consumed by adult animals (10,29%). Representative plant samples were collected monthly for nutrient analyses. Daily activities included resting (75.4%), feeding (12.6%), moving (7.5%), and other behaviors (4.0%). Feeding time was distributed over 85 plant species and parts, including leaves (52.9%), fruits (31.2%), flowers (13.1%), and other items (2.0%). Ninety-one plant parts were analyzed for proximate, fiber, and mineral composition. Combining observations of bite counts, bite weights, and

nutrient analyses, a nutrient profile of the natural diet was developed. Daily fresh weight and dry matter intake were estimated to be 544.0 g and 151.0 g, respectively. Daily gross energy intake was estimated to be 711.2 kcal. Nutrient concentrations in diet dry matter were estimated to be 13.4% crude protein, 51.5 % NDF, 37.8% ADF, and 19.0% cellulose.

Based upon estimated fiber concentrations in the wild diet, a higher fiber primate diet was developed for captive feeding trials. Two test diets with different levels of fiber (15% ADF, 30% ADF) were fed to primates with simple qastrointestinal tracts (ruffed lemurs), hindgut fermentation (howlers), and foregut fermentation (colobines). Apparent digestibility (%) of dry matter (DM), gross energy (GE), and fiber components (NDF, ADF, C, HC) were measured in the three gut types. Rates of digesta transit (TT1) and retention times (Rait) were assessed using acetate beads, Co-EDTA, and Crmordanted fiber. Animals with simple gastrointestinal tracts had lowest digestibilities (%) of DM (51.0, 41.7), GE (47.0, 39.7), and NDF (20.4, 20.7) for 15ADF and 30ADF, respectively. Hindgut fermenters exhibited greater digestive capabilities for (%) DM (69.3, 61.7), GE (68.5, 61.5), and NDF (44.8, 47.4) for 15ADF and 30ADF, respectively. Colobine primates were most efficient in digesting (%) DM (81.2, 76.6), GE (80.9, 75.6), and NDF (77.1, 74.7) in 15ADF and 30ADF, respectively. TT1 was fastest in simple gut species and slowest in foregut fermenters. Rate was shortest in simple gut species and longest in foregut fermenters. No significant differences between marker types were detected. Development of a higher fiber (>25% ADF) primate diet is warranted based upon these results.

DEDICATION

This dissertation is devicated to my parents, Silism
Patrick and Susanna arranges and or wife. Sichette. Their
unwavering engine!

Copyright by Mark Stephen Edwards 1995

DEDICATION SECOND SECON

This dissertation is dedicated to my parents, William Patrick and Suzanne Edwards, and my wife, Michelle. Their unwavering support, love, and sacrifice throughout my life have created endless opportunities for me.

ACKNOWLEDGEMENTS

As with any project of this magnitude, the number of people who helped toward the completion of this work, to which I owe a great deal, is great. My appreciative thanks go out to my graduate committee for their support and guidance throughout the duration of my program. Dr. Duane E. Ullrey, my major advisor, has provided inspiration, encouragement, patience, and most of all, a myriad of opportunities that most students only dream. Additionally, Dr. Ullrey and his wife, Susanne, provided me with "student housing" during the later "traveling" portions of my project, which was more than generous. It was a special privilege to work with the other members of my committee: Drs. Sue Crissey, Richard Dukelow, Olav Oftedal, Norm Sauer, and John Shelle. Each of them contributed their special talents to my academic and professional career. Additional thanks should be extended to Dr. Richard Aulerich, for his review and comments of this dissertation.

Special thanks go to Drs. Oftedal and Crissey, with whom I worked, as a field assistant for the Department of Zoological Research of the National Zoological Park, before my enrollment at Michigan State University. Their confidence and friendship were a guiding presence throughout the duration of the Red Howler Project in Venezuela. The memory of their

visits to the field study site is the stuff of which legends are made. The generosity of Thomas and Cecelia Blohm in allowing us to conduct the studies at Hato Masaguaral will never be forgotten. The initial portion of this study was funded by the International Environmental Studies Program of the Smithsonian Institution. Access to those funds were due, in part, to the cooperation of Dr. "Rudy" Rudran. The assistance and comradery of Kristen Silvius, Julie Currier, Virginia and David Edwards were essential to completion of the field work.

The captive studies would not have been possible without the cooperation of the Cleveland MetroParks Zoo and the Zoological Society of San Diego. The staff of both institutions welcomed me, and made me feel as a part of their team. My impressions of family life and the enormous responsibilities it includes were forever altered by my time living with the Lewandowski family. Dr. Albert Lewandowski, and his wife, Karen, welcomed me into their home, during my stay in Cleveland. I could never thank them enough for their generosity.

The opportunity to work with such a variety of folivorous primates at each zoo was made possible through the courtesy of a number of individuals including: Dr. Don Janssen, Karen Killmar, Dr. Albert Lewandowski, Tom Silva, and Allen Sironen. Their participation in this project is only a small

demonstration to their commitment to the health and preservation of these species through captive propagation.

Working with the animal keepers, who tolerated and assisted with the collection of the captive animal data made the time spent away from home so much more enjoyable. I am fortunate to have many of those people as my friends to this day. The participation of Connie Carson, Elaine Chu, Ann Dahl, Joe Kustra, Rachél Rogers, Jennifer Sanders, Tad Schofner, David Smith, Michele Stancer should not go without mention.

The test diets which were fed during the captive studies were donated through the generosity of Jim Goetz and Marion Zoological, Inc. This type of support from the sector of the feed industry which supplies foods for our captive wildlife is essential to expand our knowledge base.

The large volume of analytical work which was need to complete this project could not have been done without the assistance of Dr. Pao Ku and Sharon DeBar. Their instruction, guidance, and friendship as we worked through the seemingly endless quantity of samples was essential therapy. The additional support of Jennifer Showers, Bethany Lutz, and Karen Miller made it possible to meet our "deadlines" before the demolition of the Comparative Animal Nutrition Lab in Anthony Hall.

The advice and assistance of Dr. Mike Allen concerning

digesta marker synthesis, analysis, and related calculations was invaluable.

The friendships which develop among graduate students are critical for interpersonal support during those discouraging periods of self doubt in graduate school. My time spent in East Lansing will always be an important part of my life, due in part to my fellow graduate students, particularly Joni Bernard, Stacey Doumit, Josep Garcia, Cammie Heleski, Kim Howard, Scott Kramer, Daniel Nelson and Mike Schlegel.

To my wife, Michelle, I am most grateful. Her extraordinary patience, understanding and love enabled me to take advantage of incredible opportunities which led to the completion of this degree.

TABLE OF CONTENTS

	vi	
Acknowledgemen	nts vii	
	sxiii	
List of Figure	es xxi	
CHAPTER I. N	UTRITIONAL STRATEGIES OF HERBIVORES	
Plar Mamm Prim Brie	roduction	
CIMILIEN 11.	FEEDING HABITS AND NATURAL DIETARY NUTRIENT COMPOSITION OF FREE-RANGING RED HOWLER MONKEYS (Alouatta seniculus)	
Mate St Cl Ve Pl Be Fl In Nu Resu Be Pl	Aground	7 7 50

Plant Sample Collection and Analysis Intake Estimation Nutrient Composition of Consumed Diet	100 106 106 109 115
HAPTER III. DIGESTIVE CAPABILITIES AND DIGESTA TRANSIT LEAF-EATING PRIMATES	IN
Feeding Habits of Folivorous Primates	117 121 127
General Test Diet Formulation Animals, Housing, and Sample Collection Chemical Analysis of Feed and Feces Rate of Digesta Passage Measurements	134 135 136 140
Marker Administration and Sampling Marker Synthesis Analysis of Markers	143 144 147
Results Body Weight Changes Dry Matter Intake Apparent Dry Matter Digestibility Energy Digestibility and Energy Intake Fiber Component Digestibility Markers Discussion	149 152 152 154 154 158
General Varecia variegata spp. Alouatta spp. Colobines	175 187 188
Literature Cited	
Appendix A	
Appendix B	259

LIST OF TABLES

CHAPTER I	
Table 1-1.	Classification of some mammals based on gastrointestinal anatomy
Table 2-1.	Measurements of body length, and surface area, and volume of stomach, small intestine, cecum, and colon in howler and colobine
	monkeys
CHAPTER II	
Table 2-1.	Summary of total annual precipitation for Corozo Pando, a village 8 km south of Hato Masaguaral
Table 2-2.	Demographic composition of two red howler monkey study groups
Table 2-3.	Example of a single month's observation schedule for MSE group
Table 2-4.	Example of field notes used for recording behavioral observations of focal animal in LM1 group
Table 2-5.	Selected nutrient composition of a "higher" fiber primate diet fed to captive red howlers in Venezuela
Table 2-6.	Distribution (%) of daily activities of three adult (10,299) red howlers (group MSE) 66
Table 2-7.	Distribution of daily activities of three adult (10,299) red howlers (group LM1) expressed on an annual, dry season, and wet season basis
Table 2-8.	Food plant species consumed by red howlers

	(group LM1), ranked by total annual feeding time
Table 2-9.	Food plant species consumed by red howlers (group LM1), ranked by total dry season feeding time
Table 2-10.	Food plant species consumed by red howlers (group LM1), ranked by total wet season feeding time
Table 2-11.	Average fresh weight of plant parts consumed by red howlers (Alouatta seniculus) in the central llanos region of Venezuela 90
Table 2-12.	Mean bite weight (g) of plants consumed by red howlers (Alouatta seniculus) in the central llanos region of Venezuela 93
Table 2-13.	Mean body weight of free-ranging red howlers (Alouatta seniculus) in the central llanos region of Venezuela96
Table 2-14.	Body weight and daily feed intake of nutritionally complete, "higher" fiber primate diet by captive red howlers (Alouatta seniculus)96
Table 2-15.	Calculated proximate and fiber composition of the mean annual diet selected by three adult (10,299) red howlers (group IM1)97
Table 2-16.	Calculated macro- and trace mineral composition of the mean annual diet selected by three adult (10,299) red howlers (group LM1)
Table 2-17.	Proximate nutrient composition (dry basis) of total diet as consumed by red howlers (group LM1) in the central llanos region of Venezuela calculated based on time spent feeding98
Table 2-18.	Macro and trace mineral composition (dry basis) of tot al diet as consumed by red howlers (group LM1) in the central llanos region of Venezuela calculated based on time spent feeding

Table 2-19.	gastrointestinal tract of red howlers (mm)	11
CHAPTER III		
Table 3-1.	Diet fed to two black and white colobus during pretrial and collection periods to determine digestibility	25
Table 3-2.	Select nutrient composition of higher fiber "cubed" diet fed to silvered leaf monkeys . 12	6
Table 3-3.	Diet fed to a group of six (one male, five females) proboscis monkeys at the New York Zoological Park	8
Table 3-4.	Apparent dry matter, fiber and energy digestion coefficients of various diets by leaf-eating primates and humans as determined in previous studies	9
Table 3-5.	Transit time (TT_1) , 50% recovery, and mean retention time (R_{git}) of various markers fed to leaf-eating primates	1
Table 3-6.	Classification of markers used to evaluate rates of digesta passage	2
Table 3-7.	Experimental feeding trials and data collected	4
Table 3-8.	Composition (percentage by weight, air-dry basis) of the lower and higher fiber diets 137	7
Table 3-9.	Estimated nutrient composition (dry matter basis) of the test diets	3
Table 3-10.	Scientific and common names of species included in the study	,
Cable 3-11.	Mean body weight (BW) and mean dry matter intake (DMI) of primate study subjects fed diets 15ADF and 30ADF grouped by gastrointestinal tract (GIT) type	
able 3-12.	Mean body weight (BW) and mean dry matter intake (DMI) of primate study subjects fed diets 15ADF and 30ADF grouped by species/	

	subspecies
Table 3-13.	Mean apparent dry matter digestibility (DMD), digestible energy (DE) of diets 15ADF and 30ADF, and digestible energy intake (DEI) for primate study subjects grouped by gastrointestinal tract (GIT) type
Table 3-14.	Mean apparent dry matter digestibility (DMD), digestible energy (DE) of diets 15ADF and 30ADF, and digestible energy intake
	(DEI) for primate study subjects grouped by species/subspecies
Table 3-15.	Mean digestibility (%) of various fiber components (NDF, ADF, C, HC) in diets 15ADF and 30ADF for primate study subjects grouped by gastrointestinal tract (GIT) type 15
Table 3-16.	Mean digestibility (%) of various fiber components (NDF, ADF, C, HC) in diets 15ADF and 30ADF for primate study subjects grouped by species/subspecies
Table 3-17.	Mean transit time (TT ₁) of various markers, acetate bead, Co-EDTA, and Cr-mordanted fiber, in diets 15ADF and 30ADF for primate study subjects grouped by gastrointestinal tract (GIT) type
Table 3-18.	Mean transit time (TT ₁) of various markers, acetate bead, Co-EDTA, and CT-mordanted fiber, in diets 15ADF and 30ADF for primate study subjects grouped by species/ subspecies
Table 3-19.	Mean retention time $(R_{\rm yit})$ of various markers, acetate bead, Co-EDTA, and Cr-mordanted fiber, in diets 15ADF and 30ADF for primate study subjects grouped by gastrointestinal tract (GIT) type
Table 3-20.	Mean retention time $(R_{\rm git})$ of various markers, acetate bead, Co-EDTA, and Cr-mordanted fiber, in diets 15ADF and 30ADF for primate study subjects grouped by species/ subspecies
Table 3-21.	Digestible energy (DE) and dry matter intake (DMI) of two nonspecialized herbivores of

	Similar body weight
Table 3-22.	Transit time (TT_1 , h) and total mean retention time (R_{pit} , h) of particulate markers for two nonspecialized herbivores of similar body weight 168
Table 3-23.	Apparent digestibility (%) of plant cell wall (NDF), hemicellulose (HC), and cellulose (C) by howler monkeys (Alouatta spp.) 171
Table 3-24.	Digestibility (%) of cellulose (C) and hemicellulose (HC) by various nonruminant animals
Table 3-25.	Fermentation contents (FC) (rumen or saccus + cecum + colon) of species with foregut fermentation, expressed as a percentage of BW
Table 3-26.	Features of gut dimensions in colobine monkeys
Table 3-27.	Digestibility (%) of dry matter and neutral- detergent fiber (NDF) by species with foregut fermentation
Table 3-28.	Body weight (BW), estimated daily energy production from microbial fermentation (MFE) of digesta based upon calculations by Bauchop and Martucci (1968), estimated basal metabolic rate (BMR) and percent of MFE used for BMR by colobines
Table 3-29.	Rates of digesta passage (TT_1 ,h) and mean retention time (R_{glt} , h) determined in two langur species fed similar higher fiber (15% ADF) diets
Table 3-30.	Comparative passage rates of solid and liquid digesta in species with simple GIT, hindgut, and foregut fermentation 190
APPENDIX A	
Table A1-A7.	Proximate and fiber composition (dry basis) of plants consumed by red howler monkeys (Alouatta seniculus seniculus) in the central llanos of Venzuela

Table A8-A1	4. Macro mineral composition (dry basis) of plants consumed by red howler monkeys (Alouatta seniculus seniculus) in the central llanos of Venzuela
Table A15-A	21. Trace mineral composition (dry basis) of plants consumed by red howler monkeys (Alouatta seniculus seniculus) in the central llanos of Venzuela
APPENDIX B	
AFFENDIA D	
Table B1.	Body weight (BW) of all study subjects prior to, during, and following feeding trials
Table B2.	Body weight (BW) and dry matter intake (DMI) of ruffed lemurs (Varecia spp.) fed diet 15ADF and apparent digestibility of dry matter (DM) and fiber fractions in that
Table BIZ.	diet
Table B3.	Body weight (BW) and dry matter intake (DMI) of ruffed lemurs (Varecia spp.) fed diet
	30ADF and apparent digestibility of dry matter (DM) and fiber fractions in that diet
Table B4.	Body weight (BW) of ruffed lemurs (<i>Varecia</i> spp.) fed diets 15ADF and 30ADF and the
	apparent digestibility of dry matter (DM) and gross energy in those diets 263
Table B5.	Body weight (BW) and dry matter intake (DMI) of howler monkeys (Alouatta spp.) fed diet 15ADF and apparent digestibility of dry
	matter (DM) and fiber fractions in that diet
Table B6.	Body weight (BW) and dry matter intake (DMI) of howler monkeys (Alouatta spp.) fed diet 30ADF and apparent digestibility of dry matter (DM) and fiber fractions in that diet
Table B7.	Body weight (BW) of howler monkeys (Alouatta spp.) fed diets 15ADF and 30ADF and the

	apparent digestibility of dry matter (DM) and gross energy in those diets 266
Table B8.	Body weight (BW) and dry matter intake (DMI) of colobines fed diet 15ADF and apparent digestibility of dry matter (DM) and fiber fractions in that diet
Table B9.	Body weight (BW) and dry matter intake (DMI) of colobines fed diet 30ADF and apparent digestibility of dry matter (DM) and fiber fractions in that diet
Table B10.	Body weight (BW) of colobines fed diets 15ADF and 30ADF and the apparent digestibility of dry matter (DM) and gross energy in those diets
Table B11.	Digesta transit time (TT_1) and total retention time ($R_{\rm git}$) of ruffed lemurs ($Varecia$ spp.) fed diet 15ADF
Table B12.	Digesta transit time (TT,) and total retention time ($R_{\rm git}$) in hours of ruffed lemurs ($Varecia$ spp.) fed diet 30ADF 271
Table B13.	Digesta transit time (TT ₁) and total retention time ($R_{\rm git}$) in hours of howler monkeys (Alouatta spp.) fed diet 15ADF 272
Table B14.	Digesta transit time (TT $_1$) and total retention time (R $_{\rm git}$) in hours of howler monkeys (Alouatta spp.) fed diet 30ADF 273
Table B15.	Digesta transit time (TT ₁) and total retention time (R_{git}) in hours of colobine primates fed diet 15ADF
Table B16.	Digesta transit time (TT_1) and total retention time (R_{git}) in hours of colobine primates fed diet 30ADF
Table B17.	Body weight (BW) and dry matter intake (DMI) of two cercopithecine monkeys fed diet 15ADF and apparent digestibility of dry matter (DM) and fiber fractions in that diet 276
Table B18.	Body weight (BW) and dry matter intake (DMI) of two cercopithecine monkeys fed diet 30ADF and apparent digestibility of dry matter

	(DH) and liber fractions in that diet	276
Table B19.	Body weight (BW) of two cercopithecine monkeys fed diets 15ADF and 30ADF and the apparent digestibility of dry matter and gross energy in those diets	277
	Colobine primate gastrointestinal trect: an adult black and white colobus (Colobus gueres), body length = 50 cm. ce-drawn from Stevana (1988), Sodia equals 20cm	

LIST OF FIGURES

CHAPTER I		
Figure 1-1.	Colobine primate gastrointestinal tract: an adult black and white colobus (Colobus guereza), body length = 50 cm, re-drawn from Stevens(1988). Scale equals 20cm	17
Figure 1-2.	Colobine primate gastrointestinal tract: an adult (3 yr) female northern douc langur (Pygathrix n. nemaeus), body length = 52.0 cm, body weight = 6.09 kg, drawn directly from a photograph (Edwards, 1993). Scale equals 11 cm	18
Figure 1-3.	Cebid primate gastrointestinal tract: an adult, red howler (Alouatta seniculus) drawn from measurements and descriptions given by	20
CHAPTER II		
Figure 2-1.	Political map of Venezuela with location of field study site (Hato Masaguaral)	43
Figure 2-2.	Map of Hato Masaguaral with location of study area in box. Scale equals 1 km	44
Figure 2-3.	Average maximum and minimum temperatures (Estacion Biologica, SVCN) (Troth, 1979)	47
Figure 2-4.	Study site and estimates of study group (MSE, LM1) territories	50
Figure 2-5.	Major plant parts (leaf, fruit, flower) consumed by red howlers (group MSE), listed as percent of total feeding time	57
Figure 2-6.	Specific plant parts consumed by red howlers (group MSE), listed as percent of total	

	reeding time 68	
Figure 2-7.	Major plant parts (leaf, fruit, flower) consumed by red howlers (group LM1), listed as percent of total annual feeding time 70	
Figure 2-8.	Major plant parts (leaf, fruit, flower) consumed by red howlers (group LM1), listed as percent of total dry season feeding time 71	
Figure 2-9.	Major plant parts (leaf, fruit, flower) consumed by red howlers (group LM1), listed as percent of total wet season feeding time 72	
Figure 2-10.	Specific plant parts consumed by red howlers (group LM1), listed as percent of total annual feeding time	
Figure 2-11	Specific plant parts consumed by red howlers (group LM1), listed as percent of total dry season feeding time	
Figure 2-12	Specific plant parts consumed by red howlers (group LM1), listed as percent of total wet season feeding time	
Figure 2-13	Major plant parts (leaf, fruit, flower) consumed by red howlers (group LM1), listed as percent of calculated mass intake 101	
Figure 2-14.	Specific plant parts consumed by red howlers (group LM1). listed as percent of calculated mass intake	

Chapter I

Nutritional Strategies of Herbivores

Introduction

The co-evolution of plants and the animals that eat them has led to adaptations of both the consumers and the plants. Mammals that consume plant materials rich in structural carbohydrates are commonly referred to as herbivores. Eisenberg (1978) distinguished between species that consume grasses and forbs with little or no woody tissue, which were termed herbivores, and species that specialize on a diet of arboreal leaves, which he termed folivores. The ability of animals to use the structural carbohydrates in plants as food is dependent upon the capacity of gastrointestinal microbes to degrade them and of the ability of the host to use the end-products of microbial fermentation and the microorganisms themselves (Van Soest, 1982).

Plant Material as a Food Source

Structural polymers are the most abundant sources of energy in plants (Parra, 1978). However, vegetative plant

parts are relatively low in digestible nutrient concentrations (Langer, 1988). The structural carbohydrates which make up plant cell walls, include cellulose, hemicelluloses, and related compounds (lignins) that are resistant to digestion by mammalian enzymes. Availability of the energy within these compounds depends on an ability to cleave glycosidic bonds within plant carbohydrates and between plant carbohydrates and other substances (Van Soest, 1982). Mammalian digestive enzymes can hydrolyze α 1-4 and α 1-6 glycosidic linkages in starches and sugars. However, enzymes required to hydrolyze the β 1-4 glycosidic linkages between individual glucose units of cellulose and hemicellulose are lacking in the digestive secretions of almost all animals (Moir, 1968; Van Soest, 1977).

Cellulose is the most abundant carbohydrate in the world, but is not found independently in plant cell walls. Rather, it is combined with lignins, hemicellulose, cutin, and minerals. The availability of cellulose as a source of energy for herbivores varies from total digestibility to complete indigestibility, depending upon the extent of lignification and the presence of other interfering compounds (Van Soest, 1982). Studies of the relationship of cellulose to lignin have produced alternative views concerning the nutritive value of cellulose. One view suggests that the nutritive value of cellulose is regulated by its ratio with lignin. An opposing

view is that there are two types of cellulose, one of which is lignified and resistant to digestion and another that is free of the protective properties of lignin (Van Soest, 1982).

Hemicellulose is a highly complex mixture of polysaccharides with β 1-4 linkages in the main xylan core polymer. Hemicellulose is more closely associated with the lignin fraction of the plant cell wall than other polysaccharides (Sullivan, 1966). Like cellulose, the digestibility of hemicellulose is negatively correlated with the extent of lignification. However, the digestion of hemicellulose by ruminants is not accomplished in the same manner as is digestion of cellulose. Nonruminant herbivores typically digest relatively more hemicellulose than cellulose, whereas ruminants usually digest approximately equal quantities of both carbohydrates (Van Soest, 1994).

After cellulose, lignin is the most abundant of all natural polymers (Nimz, 1974) and is comprised of high molecular weight phenylpropanoid units. Lignin is not a single substance but a collection of monomeric units of different dimensions. As a result, the characteristics of lignin fractions are variable among plant species and among plants at different stages of maturity. Grass lignins are considerably more soluble in alkali than are lignins from either woody plants or forage legumes (Van Soest and Wine, 1967). The structural rigidity that lignins provide to plants

may be one of their most important benefits, but their presence tends to lower the value of plants as food.

The degradation of lignin is dependent upon oxidation and is resistant to anaerobic microbial fermentation in the gastrointestinal tract. As a consequence, lignin is a very important factor in reducing the digestibility of plant fiber by herbivores (Van Soest, 1967; Deinum, 1971; Parra et al., 1972). The precise manner in which lignin limits digestibility is the subject of numerous theories and is poorly understood.

Crude lignin is the collective analytical term for the noncarbohydrate fraction of plants that is resistant to enzymatic and acidic hydrolysis. There is no routine analytical procedure that will determine true lignin (Van Soest, 1982).

Other components of the plant cell wall include pectins, xylans, and gums. Pectin is a polysaccharide, high in galacturonic acid content, which has cementing properties. Although included with other soluble fractions in the measure of dietary fiber used in human nutrition, pectin is dissolved by neutral detergent (ND) and thus is not included with other plant cell wall components in the neutral detergent fiber (NDF) fraction or residue.

Cutin is a structural, aliphatic polymer component of the plant cuticle. Suberin, a mixed aromatic-aliphatic polymer, is found between the plant cell wall and cytoplasm. The cell walls of browse plants may contain up to 15% of a cutin-suberin complex (Robbins and Moen, 1975). Although these two compounds are partially digestible (Brown and Kolattukudy, 1978), they may reduce the digestibility of more readily available cell wall polysaccharides through physical obstruction.

Secondary plant metabolites comprise a broad group of compounds which exhibit a range of characteristics from bitter taste to toxicity. They include, but are not limited to, tannins, flavonoids, terpenes and essential oils, alkaloids, and coumarins. Few of these compounds have any primary metabolic functions within the plant, hence their classification as "secondary compounds".

Tannins are low molecular weight, polymerized, soluble phenolic compounds. Although tannins are only one of many groups of compounds that are responsible for feeding aversion to plants, they appear to be the most important (Swain, 1978). Four groups of tannins are defined based upon chemical structure, molecular weight, water solubility, and tannin action: hydrolyzable and condensed tannins (proanthocyanidins), oxytannins, and a group of miscellaneous tannins referred to as β -tannins (Swain, 1977; Swain, 1978). Tannins are capable of binding almost all soluble proteins and macromolecules such as nucleic acids and polysaccharides. The

result is the precipitation of cellular proteins and dramatic reductions in digestive enzyme activity, sometimes to zero (Rhoades and Cates, 1976; Lohan et al., 1981). Tannins found in browse plants consumed by ruminants may also inhibit cellulase activity of microbes (Mandels and Reese, 1963; Robbins et al., 1975).

Evidence supporting the role of tannins as a defensive mechanism of plants against herbivory has been explored (Levin, 1971; Levin, 1976; Harbourne, 1977; Swain, 1977). Feeding studies, including those with Colobus spp. (Oates et al., 1977) and Gorilla spp. (Harbourne, 1977), suggest that tannin concentrations greater than 2% may deter feeding.

The terpenoid class represents one of the largest and biologically most important classes of natural products (Mabry and Gill, 1979), which include steroids and saponins. Saponins are a widespread group of secondary plant metabolites characterized by their "soaplike", foam-forming properties. This action of saponins has been suggested to contribute to ruminant bloat. Saponins may alter the surface tension of the ruminal contents, leading to entrapment of gaseous microbial end-products as a froth (McCandlish, 1937; Quin, 1943; Applebaum and Birk, 1979). Nonruminant species exhibit different responses to saponins, with food aversion as a primary response, associated with their bitter taste (Cheeke, 1976).

A group of aromatic secondary plant metabolites known as flavonoid pigments, contribute to the color of plants. The toxicity of common flavonoids to insects has been established with studies of the tobacco bud worm (Heliothis virescens) (Shaver and Lukefahr, 1969). However, common flavonoids are apparently harmless to humans (Harborne, 1979).

The alkaloids are structurally a very diverse group of compounds (Robinson, 1979) that can exhibit pharmacological activity and inhibit digestion.

Mammalian Adaptations to Herbivory

Mammalian herbivores have developed a wide variety of physical adaptations which promote, either through symbiotic microbial fermentation or mechanical action, the destruction of the structural and chemical defenses of the plants upon which they feed. Van Soest (1982) classified several herbivorous mammals based upon their gastrointestinal anatomy (Table 1-1).

The presence of bacteria, and frequently protozoa, in the digestive tract of most animals is common. Plants which contain a high concentration of structural carbohydrates can be digested with the assistance of these microorganisms. The benefit of these organisms to the host animal, and the end-products produced, is directly related to the structure of the digestive tract and the chemical composition of foods

Table 1-1. Classification of some mammals based on gastrointestinal tract anatomy (Van Soest, 1982).

Class	Species	Dietary habit
Pregastric fermenters	i y -	ory out out ore ori ore
Ruminants	Cattle, sheep	Grazing herbivore
	Deer, antelope	Selective herbivore
Nonruminant	Colobine monkey,	Selective herbivore
	hamster, vole	
	Kangaroo,	Grazing and selective
TO SERVICE STREET	hippopotamus	herbivore
Hindgut fermenters		EROS EROS EROS EROS EROS EROS EROS EROS
Cecal (rodents)	Capybara	Grazer
	Rabbit	Selective herbivore
nt.	Rat	Omnivore
Colonic digester	and the second	Pa t t no
Sacculated	Horse	Grazer
	New World monkey	Selective herbivore
	Pig, human	Omnivore
Unsacculated	Dog	Carnivore
at dig	Cat	Carnivore

consumed.

Two principle gastrointestinal tract types that support herbivory involve enlargements of the stomach or the hindgut to accommodate microbial fermentation (Parra, 1978; Langer, 1988). A third type of gastrointestinal tract seen in a few herbivores (e.g., pandas) involves no enlargements or sacculations (Dierenfeld et al., 1982; Warnell, 1988; Warnell et al, 1988).

Ruminants and other foregut fermenters have a large sacculated foregut that is the primary site of microbial activity. Within the foregut are anaerobic cellulolytic bacteria, and other microbial symbionts, which produce enzymes that degrade plant cell walls and promote access to cellular contents. The foregut fermenter class of gastrointestinal tracts permits microbial fermentation to precede gastric and enzymatic digestion. Moir (1968) listed a number of animals which possess a sacculated foregut-type digestive tract including macropod marsupials, some edentates, hippopotami, camels, and colobine monkeys. These species exhibit evolutionary convergence with ruminants in their adaptations of foregut structure for folivory (Moir, 1968).

The process of rumination (regurgitation and remastication of digesta) is an important mechanism that assists in reducing the protection afforded cell contents by lignin. The additional physical processing by rumination helps reduce

the particle size of lignified plant tissues, creating "openings" that permit access for microbial symbionts. However, not all species that have adaptations for foregut fermentation ruminate. Thus, the type of plant fiber and extent of lignification may have different effects on these species than on ruminants.

In animals with hindgut fermentation, symbiotic microorganisms occupy enlarged areas distal to the gastric. These animals are unable to recover quantitatively as high a proportion of the nutrients produced by fermentation as are foregut fermenters. This decreased return on microbial fermentation may be due, in part, to a faster rate of digesta passage through the lower gut, thus limiting the time available for absorption.

Herbivorous species that lack apparent gastrointestinal adaptations for fermentation present a paradox to nutritionists and physiologists. The contribution of plant material to the diet of these species varies from only a fraction to 100%. However, the extent to which these species benefit from fermentation by microbial symbionts is limited. These species (e.g., red pandas, giant pandas) have developed other strategies to exist on a relatively indigestible diet. These include increased rates of digesta transit and short retention time allowing consumption of relatively large daily amounts of food of low nutrient value (Warnell, 1988; Warnell

et al., 1989).

Anaerobic fermentation of plant material by gastrointestinal microorganisms converts substrate into microbial cells and fermentation end-products. The end-products of fermentation include volatile fatty acids (VFAs), carbon dioxide, and methane. The VFAs produced, acetic, butyric, and propionic, can be important sources of energy for the host. The microbial cells can be important sources of B-complex vitamins, amino acids, and vitamin K.

As indicated above, adaptations to feeding on plant material demonstrated by mammals are not limited strictly to gastrointestinal tract modifications. The chemical composition of plant tissues varies greatly among different plant parts and among stages of plant maturity. This is true with respect to both structural carbohydrates, lignin, and some secondary plant metabolites. For example, mature oak leaves have a higher concentration of tannins than immature oak leaves (Feeny, 1970).

Specific plant parts are selectively consumed by a large number of herbivores. This is a "behavioral" adaptation to the selection of plant parts that may be most appropriate for the anatomy and physiology of the species. Hoffmann (1973) grouped East African ruminants into three major classes based upon gastrointestinal anatomy and feeding habits. Others have extended these categories to additional herbivores.

Herbivores designated as "concentrate selectors" consume plant materials that are relatively low in plant cell walls. These concentrate selectors do not use large amounts of dietary plant fiber effectively and thus selectively feed on those plant parts that are higher in readily fermentable carbohydrates. Herbivores which consume plants that are relatively higher in cell wall components, typically grasses, are classified as "bulk and roughage eaters". The third class consists of "intermediate feeders" that are more adaptable to changing habitats and vegetation than most of the concentrate selectors and bulk and roughage feeders. Opinions differ as to which group is more highly developed. Among ruminants, concentrate selectors are commonly considered the most conservative (Van Soest, 1982). However, the ability of a ruminant species to tolerate and use the end-products of rapid fermentation, as seen in concentrate selectors, might also be considered as more recently evolved.

Primates as Herbivores

The differences in digestive tract anatomy associated with the three classes of herbivores above also can be found within the order Primates. Further, five categories of dietary adaptation can be defined: faunivory, exudivory, granivory, frugivory, and folivory, the latter two representing specializations of herbivory.

Primates which specialize in consuming leaves exhibit several gastrointestinal adaptations. Microbial fermentation occurs in the large sacculated forestomach of Colobidae (e.g., Colobus spp., Presbytis spp., Trachypithecus spp., Pygathrix nemaeus, Nasalis larvatus). Hindgut fermentation, in the sacculated cecum and/or colon, occurs in certain other primates (e.g., Alouatta spp., Indriidae). However, the extent of herbivory demonstrated by a given species is variable. Eisenberg (1978) described five classes or degrees of herbivory:

Class 1. Digestive tract shows little modification; dentition
little modified for grinding; essentially omnivorous, but much
"fruit" is utilized when available; some feeding on buds or
shoots may occasionally be seen.

<u>Class 2</u>. Digestive tract shows little modification for bacterial symbionts. Dentition may show slight modification for grinding or crushing. In addition to some animal food, often feeds on plant parts including seeds, fruit, buds, flowers, leaves, and sap.

Class 3. Digestive tract shows some modification, generally, cecal enlargement and/or lengthening of the intestine.
Dentition may show cusp modification to create crushing surfaces. Diet differs from Class 2 in that buds, blossoms, leaves, and young shoots begin to dominate in the diet (from 30 to 40 percent). If terrestrial, the species may ingest occasional grasses/forbs, as well as leaves.

Class 4. Digestive tract further modified from Class 3. Stomach is often very enlarged and sacculated or cecum is greatly enlarged. Some bacterial digestion of cellulose in the stomach is implied. Leaves and grass comprise a significant percentage of the diet (greater than 40 percent) although blossoms, fruit, and other plant parts contribute to the diet. If terrestrial, the species may be considered either a browser or grazer, depending on the proportion of leaves in its diet.

<u>Class 5.</u> Differs from Class 4 in that the species, if arboreal, is close to being an "obligate" folivore. Gut is specialized for the breakdown of cellulose by bacteria and/or protozoan symbionts. Leaves, buds, and plant stems predominate in the diet. Specimens of arboreal forms taken as advanced juveniles or older for captive rearing are almost impossible to convert to an artificial diet. If terrestrial, the species is generally a specialized browser or grazer.

It has been demonstrated that free-ranging primates do

not eat plant materials at random, but show decided preferences (Milton, 1984). Selective consumption of plant parts, which may be lower in structural carbohydrates, lignin, and/or secondary plant metabolites may be one such adaptation (Oftedal, 1991). Milton (1978a) suggested that the mantled howler monkey (Alouatta palliata), as a "behavioral folivore", utilizes this approach to folivory, consuming young leaves, fruits, and flowers from a diverse number of plant species.

It is difficult to generalize the feeding habits of primates whose diet consists, in part or entirely, of leaf material. Indeed, within the order Primates, a wide range of folivory is demonstrated. Much of this variation is reflected in the gastrointestinal anatomy of these species.

The colobine stomach has four parts: the presaccus and saccus, the long tubus gastricus, and the short pars pylorica (Kuhn, 1964). The large, complex presaccus and saccus are distended and sacculated proximally, with a U-shaped tube distally, and sacculated along the proximal part of the greater curvature (Hill, 1958). The sacculations seen in this region of the gastrointestinal tract are a result of longitudinal muscles, which have been reduced to two or more bands (taenia) (Chivers and Hladik, 1980).

Some variation in gut structure is seen with the subfamily Colobinae (Chivers, 1994). The stomach of the African colobines (e.g., Colobus, Procolobus) is more

elongated, with the tubular portion bent back on the saccular portion. Asian colobines (e.g., Pygathrix, Presbytis, Nasalis) have a saccular portion which is roughly spheroidal. In both groups, the hindgut is comparable, with a long sacculated colon and cecum of moderate size. Drawings of two colobine gastrointestinal tracts, the black and white colobus (Colobus guereza), and northern douc langur (Pygathrix nemaeus nemaeus) are given in Figures 1-1 and 1-2, respectively.

The presence of microbial organisms (Bauchop, 1971) and extensive microbial fermentation (Kuhn, 1964; Bauchop and Martucci, 1968) in the presaccus and saccus demonstrate the ruminant-like digestion possessed by colobine primates. The pH of the colobine foregut has been reported to be between 5.0 and 6.7 in langurs (Bauchop and Martucci, 1968) and 7.0 in colobus (Kuhn, 1964). Interestingly, although significant numbers of cellulolytic (e.g., Bacteroides spp.) and methanogenic (e.g., Methanobacterium ruminantium) bacteria are present in the foregut, protozoa have not been reported. It should be noted that there may also be considerable microbial fermentation in the cecum and colon of colobines (Milton and McBee, 1963; Ullrey, 1986).

The stomach of Alouatta palliata exhibits a great deal of complexity when compared with other cebids. The globular sac is capacious and narrows to the bent tubular pylorus. This portion of the stomach is guarded by strong pillars, and




Figure 1-1. Colobine primate gastrointestinal tract: an adult, black and white colobus (Colobus guereza), body length = 50 cm, re-drawn from Stevens (1988). Scale equals 20 cm.




Figure 1-2. Colobine primate gastrointestinal tract: an adult (3 yr), female northern douc langur (*Pygathrix n. nemaeus*), body length = 52.0 cm, body weight = 6.09 kg, drawn directly from a photograph by Edwards. Scale equals 11 cm.

rugae radiate from the cardia and run longitudinally with the body (Chivers and Hladik, 1980). The cecum of cebids is hookshaped, and the colon is simple and straight. A drawing of the gastrointestinal tract of a red howler (Alouatta seniculus) is given in Figure 1-3.

Chivers and Hladik (1980) conducted a quantitative analysis of gastrointestinal tract morphology in primates. Measurements of body length, gastrointestinal tract surface area and volume for leaf-eating primates are provided in Table 1-2. They used coefficients of gut differentiation (CGD) to make comparative assessments of the degree of specialization of a species for folivory. These values were determined using the following equation:

Those species with higher CGD values have a gastrointestinal tract that is specialized for microbial fermentation of plant material, with an enlarged stomach and/or large intestine.

Brief History of Folivorous Primates in Captivity

Numerous references, as early as 1892, to maintenance of captive folivorous primates in zoological institutions




Figure 1-3. Cebid primate gastrointestinal tract: an adult, red howler (*Alouatta seniculus*) drawn from measurements and descriptions given by Crissey et al. (1989).

	·			
	•			

Table 1-2. Measurements of body length, and surface area and volume of stomach, small intestine, cecum, and colon in howler and colobine monkeys (Chivers and Hladik, 1980)

		T.C.				•	ATPORTIN DIES COLONIA		1	10001
		. да	ns	Surface a	area (cm²)	_		Volume	Volume (cm3)	
Species	Sex	(cm)	Stom.	SI	Cecum	Colon	Stom.	SI	Cecum	Colon
Alouatta	M	51	430	1052	289	1059	839	534	391	959
palliata	M	20	230	719	148	630	328	302	145	471
	ഥ	53	296	815	156	778	479	339	138	642
Alouatta	Σ	61	180	1106	150	578	227	556	163	422
seniculus										
Colobus	Ĺτι	26	1021	556	26	549	2055	107	13	218
polykomos	FI	58	1056	925	15	630	2162	184	9	301
Presbytis	M	63	1585	1673	105	878	3974	633	80	845
entellus	M	65	1439	1167	140	160	3438	330	104	505
Presbytis	[H	48	938	740	139	548	1814	181	77	275
senex	M	59	1416	687	139	548	3357	162	96	336

Table 1-2 (cont'd). Measurements of body length, and surface area and volume of stomach, small intestine, cecum, and colon in howler and colobine monkeys (Chivers and Hladik, 1980).

		BL	Su	rface a	Surface area (cm2)			Volume (cm3)	(cm3)	
Species	Sex	(cm)	Stom.	SI	Cecum	Colon	Stom.	SI	Cecum	Colon
Presbytis	Ħ	20	1694	1929	06	996	4831	604	64	645
cristata	Ŀ	53	1175	1329	30	209	2209	385	12	285
Presbytis	M	20	1363	1953	64	029	3237	669	41	343
obscura	Ĺτι	45	926	1311	52	558	1788	397	27	264
	M	53	1351	1969	88	902	2974	734	51	588
	Ŀ	99	1282	1129	06	869	3139	327	64	546
Presbytis	M	51	1020	1386	09	532	2072	336	34	156
melalophos	M	44	694	1075	23	369	1128	247	80	132
	[24	47	1648	1796	09	552	3521	517	36	224
	M	49	1078	1389	42	695	2158	395	20	362
	E	20	1382	1695	89	848	3327	588	43	469
	न	52	1274	2021	38	512	2296	633	14	292

23

Table 1-2 (cont'd). Measurements of body length, and surface area and volume of stomach, small intestine, cecum, and colon in howler and colobine monkeys (Chivers and Hladik, 1980).

		BL		Surface (cm ²)	(cm ²)	1	37	Volume (cm3)	(cm3)	oli
Species	Sex	(cm) Stom.	Stom.	SI	Cecum	Cecum Colon	Stom.	SI	Cecum	Cecum Colon
Prebytis	M	56	1125	1672	45	637	2259	505	20	329
rubicunda										
Nasalis	Σ	64	1978	3120	100	1234	6523	1127	99	655
larvatus										
Pygathrix	E	53	1243	1512	36	578	2960	444	12	261
nemaeus	Ĺ	09	1431 1601	1601	80	854	3442	531	42	433

exist in the literature. Success in maintaining populations of folivorous primates in captivity has been widely mixed, and to date, is highly related to species (Collins and Roberts, 1978). Mitchell (1911) reported the average life-span of six species of captive langurs (n=56) as 4.3 months (mos), with the maximum being 15 mos. In 1967, the longevity record for the northern douc langur (Pygathrix nemaeus nemaeus) captivity was reported to be 4 mos at the London Zoo (Hick, 1972). Merrit (1982) reported "longevities" of a few mos for both red (Alouatta seniculus) and black howlers (A. caraya) received at the Lincoln Park Zoo (Chicago) in the late 1960s. Although Crandall (1964) described the black and white colobus (Colobus polykomos = guereza) as a very difficult subject to maintain in the zoological garden and Flower (1931) indicated the "African colobus or guereza monkeys appear to be even more difficult than the langurs to keep alive in captivity", this species has proven to be the most adaptable folivore in captive environments. In 1927, a single specimen of black and white colobus was maintained for 3 years, 24 days in the New York Zoological Park (Crandall, 1964). Even though this record seems hardly worthy of mention, at that time this animal held the longevity record. Currently, black and white colobus regularly live beyond 20 years in zoological collections (Jones, 1995). By 1971, births of third generation captive offspring were being reported (Hollihn,

1973).

Species differences in dietary habits in the wild and in the morphology of the gastrointestinal tract suggests that all captive primates should not be fed in the same manner (Ullrey, The importance of a diet that delivers higher fiber concentrations has been alluded to by several authors. adapting wild-caught colobines to captive Success in conditions is reported to be dependent upon the conditions in which the monkeys are kept immediately following capture (Hollihn, 1973; Lindbergh, 1976; Eisenberg, 1978). Urbain et al. (1955) suggested that black and white colobus could not survive in captivity without an abundance of leaves of the vine Vitis vinifera, mulberry (Morus alba), or aucuba (Aucuba These authors point to an "essential element" japonica). provided by these plant species which is not present in cultivated vegetables, fruits, or greens. Additionally, an increased frequency of feeding (i.e., six times daily) in conjunction with appropriate dietary components was suggested as an effective method of dietary husbandry (Ludwig, 1989).

Crandall (1964) indicated that problems surrounding the captive husbandry of howler monkeys could be solved, in part by a diet providing sufficient leafy bulk. Although no attempt has been made to identify the "essential elements" that should be present in the diet of folivores, the contribution of appropriate levels of plant fiber should not

be overlooked.

The development of diets for captive wild animals has historically been one of trial and error. An animal's basic feeding preference (e.g., herbivore, frugivore, omnivore) is a simplistic, but common starting point. Commercially available substitutes for food types selected by wild have been offered to determine acceptance. specimens Unfortunately, this approach placed a great deal of reliance upon "nutritional wisdom", a concept that has been declared invalid for wild animals in captivity (Ullrey, Although bearing similar names (e.g., fruits), the nutrient composition of cultivated plant foods is generally quite different from that of plant foods consumed in the wild (Edwards, 1990). Many animal managers have sought the "secret ingredient" that allows a species to thrive in captivity. However, with few exceptions, controlled studies have demonstrated that animals require specific nutrients, not specific ingredients.

Although the specialized feeding habits of leaf-eating primates have long been recognized by individuals attempting to maintain them in captivity, systems of dietary husbandry are still widely variable. In fact, folivorous primates are often fed diets that are similar to those fed to primate species that are not primarily folivorous and/or do not have pregastric fermentation (Hollihn, 1973; Watkins et al., 1985).

This has resulted in a high incidence of gastrointestinal disorders (Hill, 1964; Hick, 1972; Hollihn, 1973; Benton, 1976; Benirschke and Bogart, 1978; Heldstab, 1988; Taff and Dolhinow, 1989; Janssen, 1994).

The diversity of ingredients and diets offered to captive folivores is remarkable. The following diets were listed by Crandall (1964):

Angola colobus (Colobus angolensis), per animal, daily:

- ·Lettuce, 1 head
- ·White potatoes, raw, 8
- ·Sweet potato, raw, 1
- ·Carrots, raw, 4
- ·Bananas, 2
- ·Peanuts, whole, 25
- ·Bread, rye or raisin, 4 slices
- ·Reconstituted evaporated milk with coddled egg, 1/2 pt
- ·Baby cereal
- ·Cod-liver oil, 1/2 tsp
- ·Celery, leaves of bamboo, and Russian mulberry, when available

Mantled howler (Alouatta palliata), per animal, daily:

- ·Milk gruel (reconstituted evaporated milk, thickened with cooked, rolled oats or PablumTM, with 2 drops of multiple vitamin concentrate added)
- ·Cabbage, raw, 1/2 head
- ·Bread, raisin, slices, 3
- ·Carrot, 1
- ·Celery, stalks, 3
- ·Egg, hard-boiled, 1
- ·Grapes, 1 dozen
- ·Apple, 1
- ·Dog food, canned, 1 oz
- ·Milk gruel, 8 oz (prepared as above)

Interestingly, despite the obviously inappropriate nature of the above diets and some of the ingredients, the author (Crandall, 1964) urged caution concerning the feeding of

certain items, such as oranges or orange juice, which were believed to produce severe diarrhea in these animals. specific concern for excessive use of citrus was reiterated by Merrit (1982). Hollihn (1973) emphasized the importance of leaf material in the diet of colobine primates, and cautioned against the feeding of "easily digestible carbohydrates and fats" such as rice, bread, and nuts. Yet, in the same article, this author promoted the feeding of unpeeled fruits such as bananas, melons, and citrus fruits as a source of "ballast substances" that are "plentiful in the natural diet". Further, Hick (1972) listed the diet of douc langurs housed at the Cologne Zoo as including "a ration of cooked rice, fruits, vegetables, rose leaves, and willow twigs" which, as with Hollihn's (1973) description, included food items that are both relatively high in readily fermentable carbohydrates and in plant cell wall. Additionally, Hick (1972) indicated that the douc langurs were provided with black tea, for drinking in the morning hours, and boiled water in the evening.

These apparent inconsistencies are not limited to the dietary husbandry of colobines. Benton (1976) cited the importance of diet as a primary variable upon which the success of captive groups of howler monkeys (Alouatta spp.) rests. The author noted gastrointestinal adaptations of howler monkeys, including a "semi-sacculated" stomach, which enable members of this genus to "cope with the vast

quantities of high bulk vegetable matter normally consumed in the wild. Despite this background information, the diet fed to these animals included the following, in the morning:

- •Evaporated milk (227 g) mixed with an equal quantity of water, boiled and mixed with:
- ·Soft-boiled eggs
- ·Gevral™ protein powder (protein supplement)
- ·Lactinex[™] tablets (Lacotobacillus acidophilus)
- ·ABDECTM vitamin drops
- ·Clovite capsules
- ·Farbegen™ capsules (iron supplement)

This mixture was then poured over:

•Fresh white bread, to form a semi-solid gruel

The evening feeding, offered in two courses, included:

- ·Boiled chicken
- •Fruit-flavored yogurt
- ·Soft-boiled eggs
- ·Purina Monkey Chow™ (25% crude protein) soaked in tea
- ·Raisins
- · Peanuts
- ·Sunflower seeds
- ·Celery
- · Tomatoes
- · Carrots
- ·Apples
- · Plantains
- ·Canned figs
- · Grapes
- ·Blueberries
- · Strawberries
- ·Fresh corn, on cob
- ·Leafy green matter (lettuce, cabbage, spinach, kale)

As available, additional items were fed, including:

- · Papaya
- · Mango
- ·Blackberries
- ·Canned dog food

The author noted that quantities of individual foods offered were not large enough to allow animals a daily portion of

every ingredient. However, he felt it was more important to prevent "dietary boredom" than to ensure that each individual had the opportunity to consume preferred foods. Although no water was provided to this group of animals, they were given boiled tea, double-strength, sweetened with honey and fortified with an additional protein supplement. This practice was similar to the recommendations of Hick (1972), and appears to be based upon the misconception that these arboreal primates in the wild strictly drink water that is high in tannic acid.

The importance the above authors placed on adherence to these diets and the myths surrounding them focused on the ingredients fed and not the nutrients provided or provision of a diet that approximated the composition of foods consumed by free-ranging animals.

Those institutions that seem to have the greatest success with long-term maintenance of captive folivorous primates are those that have addressed their specialized needs (Edwards, 1990). There are three primary methods of increasing the amount of fiber in these animals' diets. The most widely applied, and easily implemented approach, is the replacement of commercial primate biscuits containing a low concentration of fiber (10-19% NDF, 5-7.5% ADF) with a higher-fiber product (25% NDF, 15% ADF). The lower fiber commercial primate diets were designed originally for omnivorous

primates, such as macaques. The commercial production of higher fiber primate diets is a fairly recently available option, as zoo feed manufacturers respond to the scientific evidence and demands of the enlightened consumer.

Another approach includes the reduction or elimination of commercially available fruits and replacement of those items with leafy greens and higher fiber vegetable material. This method is often used in conjunction with the incorporation of the higher fiber primate biscuit.

The cultivation of plant material for supplemental feeding is also an option (Griner, 1977; Ullrey et al., 1982). This practice is not as widespread, due primarily to the variability of local growing conditions and economic considerations (Koontz et al., 1988). The types of plant material cultivated also are widely variable, from indigenous species to easily grown tropical plants. Surprisingly, no attempts have been made to cultivate widely the browse plants that are consumed by free-ranging representatives of the species held in captivity.

Special considerations should be taken when offering browse to captive primates. Nutrient contributions of the browse material should be determined (Mannchen, 1989), and the impact of seasonal variability and fertilization of browse plant materials should be assessed. Additionally, protective substances such as the secondary compounds previously

described can be present within these plants (Santi and Santi, 1989). The animals to which these plants are offered (most of which have been born in captivity) should be considered naive to such defenses and should be monitored as to the parts of the plants that are consumed. As this approach to feeding becomes more common, reports of deaths directly attributable to the consumption of browse have increased (Hill, 1964; Ensley et al., 1982; Robinson et al., 1982; Janssen, 1994; Edwards, unpublished data). This is an unfortunate demonstration of the evolutionary effectiveness of mechanisms that plants have developed to protect against herbivory, as well as the consequences of relying upon "nutritional wisdom" when offering naive animals novel foods.

An ideal approach to the feeding of captive folivorous primates appears to be a combination of the above three methods:

- 1. Use a higher fiber primate biscuit.
- Reduce or eliminate fruits, and increase green leafy vegetables.
- 3. If browse plants are fed, ensure that they are appropriate.
- 4. Distribute these foods in small quantities over several feedings per day.

Chapter II

Feeding Habits and Nutrient Composition of Natural Diets of Free-Ranging Red Howler Monkeys (Alouatta seniculus)

Background

Howlers (Family, Cebidae; Subfamily, Alouattinae Elliot, 1904; Genus, Alouatta Lacepede, 1799) are large primates, with adults weighing, on average, between 7 and 9 kg (Fooden, 1964). These animals are the largest members of the New World primates (Platyrhinni). Seven species and sixteen subspecies of howler monkeys are currently recognized, including: Alouatta belzebul Linnaeus 1766, A. seniculus Linnaeus 1766, A. caraya Humboldte 1812, A. palliata Gray 1849, A. fusca Ihering 1914, A. pigra Lawrence 1933, and A. coibensis (Hill, 1962; Hershkovitz, 1969; Smith, 1970).

Howlers are the most widely distributed of all the Platyrhinni. These animals range from southern Mexico through Central America and northern South America, extending into Argentina and Brazil. This range covers a wide diversity of habitats including tropical rainforests of Panama and Brazil, cloud-forest regions of Venezuela and

Columbia, the deciduous forests of Guatemala, dry forests in Costa Rica, and llanos regions of Venezuela.

Howlers are highly arboreal primates. However, these animals will descend to ground level under certain conditions (Carpenter, 1934; M.S. Edwards, unpublished data). The tail of the howler is fully prehensile, with a naked patch of skin on the caudal surface of the tip. Howlers are named for their deep roaring call. Beneath the chin is a large swelling which is the site of the specialized hyoid bone. The hyoid bone is larger and more pronounced in the male of the species, but is also modified in the female.

Howler monkeys are considered to be the most folivorous primates within the infra-order Platyrrhini and are strict vegetarians (Carpenter, 1934; Fooden, 1964; Hladik and Hladik, 1969; Richard, 1970; Eisenberg et al., 1972; Smith, 1977; Eisenberg, 1978; Gaulin et al., 1980; Gaulin and Gaulin, 1982; Braza et al., 1983). Most of the published information concerning foraging behavior of howler monkeys has been obtained from the study of one species within the genus, the mantled howler monkey (Alouatta palliatta = Alouatta villosa). This information has been collected from two localities, Panama, specifically Barro Colorado Island (Carpenter, 1934; Hladik and Hladik, 1969; Milton, 1978a; Milton, 1978b; Gaulin et al., 1980) and Costa Rica (Glander, 1975).

Based on data collected by Hladik and Hladik (1969),

Eisenberg described the mantled howler as a species whose digestive tract shows some modification towards herbivory, specifically cecal enlargement and/or elongation of the intestine (Class 3). This is supported by Fooden's (1964) description of the red howler gastrointestinal tract as possessing high ratios of organ length to body length for stomach, cecum, and large intestine. Further, the large diameter of the cecum was argued as hindgut modifications supporting microbial fermentation of a strictly vegetarian and rather folivorous diet (Fooden, 1964; Amerasinghe et al., 1971).

The natural diet is dominated by buds, blossoms, leaves, and young shoots. These plant parts are reported to contain lower quantities of structural carbohydrates, making them more digestible than mature leaves.

When compared to the mantled howler, described as a leaf/fruit-eating primate, the red howler appears to have similar feeding habits. However, many of these descriptions are based on either direct visual observation or gut content analysis. Milton (1978a) suggested that the lack of geographic overlap among the various species of Alouatta is the result of similar dietary niches.

Howler monkeys have been described as "behavioral" folivores rather than "anatomical" folivores (Milton, 1978a). This description is based on the observations and anatomical

adaptations of the mantled howler monkey on Barro Colorado Island, Panama. This description has been accepted as valid across the entire genus, primarily due to the lack of specific information concerning the composition of food items selected by or detailed examination of gastrointestinal anatomy of other species of howlers.

The objectives of this portion of the study were to examine the feeding habits of the red howler monkey (Alouatta seniculus) in the central llanos of Venezuela, including a detailed description of the red howler's natural diet. Calculations were made to determine nutrient intake based upon behavioral observations. Bite counts of plant parts consumed by free-ranging animals and direct measurements of bite weights of the same food plants offered to the same study subjects in captivity were used to complement the behavioral data. Nutrient analysis of specific parts of food plants selected was combined with these two estimates of intake to determine the nutrient profile of the consumed diet.

Limited information has been collected on the feeding habits of the other six species of howler monkeys. Little is known of the dietary habits of either Alouatta belzebul (Braza et al., 1983) or the black howler monkey (Alouatta caraya) (Pope, 1966; Bicca-Marques and Calegaro-Marques, 1993). Carvalho (1975) listed the plants selected by A. guariba. Limited descriptions of the diet of the red howler monkey, the

species observed in this study, are available. These descriptions consist of information collected by direct field observations, gut content analysis, and scat analysis (Fooden, 1964; Izawa, 1975; Hernández-Camacho and Cooper, 1976; Braza et al., 1983). The most detailed dietary description for the red howler was provided by Gaulin and Gaulin (1982) for animals inhabiting the Colombian Andes cloud forest. This single study made estimates of plant material intake via visual observation, in contrast to the more conventional method of reporting the diet as a percentage of total feeding time. Regardless, the information collected on mantled howlers represents the majority of knowledge to date on the feeding strategies within the genus Alouatta.

A single male red howler was examined post-mortem by Racenis (1952). The stomach contents of this animal consisted primarily of Venezuelan ragweed (Ambrosia cumanensis). Racenis (1952) reported that this plant grows in large quantities on the banks of the Orinoco river, where his study was located.

The red howler monkeys found in the basin of the river Caqueta were reported to consume leaves of unidentified trees, the fruit of *Pourouma spp.*, and beans which were similar to the nuts of *Pakia oppositifolia* (Izawa, 1975) These observations were made on six separate occasions between December 1971 and September 1973. The amount of time that the

animals spent feeding and other specific information, such as time spent resting and moving, were collected in this study.

Braza et al. (1983) examined the feeding habits of the red howler monkey in the llanos of Venezuela. Their analyses of the gut contents of howler monkeys identified leaves, of an undetermined phenological stage, in 73.8 % of the stomachs examined. Fruits also were found in 73.8% of the stomach contents, whereas legume pods were found in 21.3% of the stomach contents. The limitations of gut content analysis have been discussed by numerous investigators. Gut contents provide a view only of a "single meal", since they represent only those foods selected shortly before the animal's death. There may be some difficulty in identifying plant species which have undergone mastication and partial, or extensive, peptic digestion in the stomach. Finally, the animals must be killed to conduct such an analysis, thus limiting the potential sample size.

Fecal (scat) analyses also have been conducted to determine the feeding habits of the red howler of the llanos (Braza et al., 1983). The results of these analyses indicate that leaves were found in 95.3 %, fruits in 83.2 %, and woody stalks in 64.2 % of fecal samples examined. Fecal analysis is the most noninvasive method of determining the feeding habits of a given species, but feces represent only the end-products of digestive processes (Moreno-Black, 1978). Food

items which can be identified in the scat sample are typically those which are the most resistant to digestion. This fact may explain the high proportion of woody stalks in the feces, when only 18.0% of the stomach contents contained such material. Additionally, an extensive reference collection of plant materials is required to conduct fecal analyses with accuracy.

An interesting finding of Braza et al. (1983) was the presence of grasses both in stomach contents (4.9 %) and in fecal samples (0.8 %) collected from the howlers of the llanos region. Although, it is known that these animals will descend to the ground for travel and to obtain water (Racenis, 1952; Edwards, unpublished data), direct observations of the consumption of grasses by red howlers have not been reported. Therefore, it seems unlikely that ground vegetation would constitute a dominant portion of the howler's diet (Braza et al., 1983).

Gaulin and Gaulin (1982) reported quantitative estimates of food intake based upon visual observations of red howlers. This method was used in combination with collection of the various foods selected. These estimates were based upon the number of food items consumed per "feeding bout" and then multiplying the number of items by the mean weight of the item. For example, they determined that the mean fresh weight of a mature leaf of Ficus cundinamarcensis is 2.57 g. If an

animal consumed seven of these leaves in a feeding bout, they computed that the animal ate 7×2.57 or 17.99 g of plant material, as fed basis.

Colobine monkeys (e.g., Presbytis spp., Trachypithecus spp., Pygathrix nemaeus, Colobus sp.), which are Old World species, have highly specialized, large sacculated foreguts that permit extensive fermentation of digesta by symbiotic microbes (Hill, 1962). This gastrointestinal arrangement is similar to that of ruminant artiodactylids; however, the colobine primates do not ruminate. The absence of rumination is critical when examining the feeding strategy of these animals. Fermentation by microorganisms gastrointestinal tract of these animals or in ruminants is unable to degrade the lignin in plant cell walls. Ruminants are able to reduce the particle size of this polyphenolic compound through remastication of food during rumination. Without a reduction in physical size, the lignin fibers may become entwined, forming fibrous "ropes" called phytobezoars. These phytobezoars may obstruct the gastrointestinal tract of colobines, and physically block the passage of digesta through the gut (Ensley et al., 1982; Edwards, unpublished data).

Howlers have been described as the New World analog of the Old World folivores (Hill, 1962; Hladik and Hladik, 1969; Richard, 1970; Eisenberg et al., 1972; Jolly, 1972; Eisenberg and Thorington, 1973). Howlers lack the foregut

specialization of colobines and appear to depend on an expanded cecum and colon (Amerasinghe et al., 1971; Crissey et al., 1989) as a site where fermentation processes occur. However, the significance of this adaptation for utilization of fiber has been disputed (Milton, 1980).

The red howler monkey is abundant and widely distributed throughout Venezuela (Bodini and Perez-Hernandez, 1987), and ranges into neighboring Colombia, Guiana, Bolivia, and the Amazon basin and Atlantic coastal regions of Brazil. howlers are extremely adaptable and are found in a wide array of habitats within Venezuela, including montane forests, rain forests, and deciduous forests. Hill (1962) described three subspecies of A. seniculus in Venezuela (Alouatta seniculus seniculus Linnaeus, A. s. arctoidea Cabrera, and A. s. stramineus Humboldt). A fourth, previously unrecognized subspecies which inhabits the Venezuelan llanos, has been described (Bodini and Perez-Hernandez, 1987). This previously undescribed subspecies was identified by its coloration and Based upon genomic rearrangements, Stanyon et al. (1995) placed this subspecies among Alouatta seniculus arctoidea.

Materials and Methods

Study Area

The field work portion of this project was conducted

during a 16-mo period from August 1987 through December 1988 in the central llanos of Venezuela. The llanos comprise a band of savannas extending from the base of the Andes mountains eastward to the Orinoco basin and the Atlantic ocean. The study site was located on a private ranch, Hato Masaguaral, at approximately 8° 34' N, 67° 35' W. Figure 2-1 describes the location of Hato Masaguaral within the political boundaries of Venezuela.


The study site was located approximately 2 km south of the ranch houses. This area was a recently acquired portion of the ranch and had not been previously used for studies of red howlers. Hato Masaguaral covers an area of approximately 5,702 hectares. Figure 2-2 shows the location of the study site in relation to the ranch houses and the sites of previous red howler research at the ranch (Rudran, 1979; Crockett, 1985).

Climate and Photoperiod

The llanos are in a region of dramatic seasonal changes. The 6 mo of rainfall (May through October) constituting the wet season are flanked by 2 transitional mo (April and November) (Table 2-1). The marked dry season, which begins in December, often extends into May. Although this is the typical pattern, both the duration and amount of rainfall are extremely variable (Troth, 1979).



Figure 2-1. Political map of Venezuela with location of field study site (Hato Masaguaral).



Map of Hato Masaguaral with location of study area in box. Scale equals 1 Figure 2-2. km.

Table 2-1. Summary of total annual precipitation for Corozo Pando, a village 8 km south of Hato Masaguaral.

1970 1555 7 (May ? -No.) 1971 938 7 (April ?) 1972 888 7 (April-Oc.) 1973 1424 8 (April-No.) 1974 1272 6 (May-Nove)	
1954 1535 8 (April-No. 1955 1829 8 (April-No. 1955 1829 8 (April-No. 1956 1453 6 (May-Octo. 1957 1152 7 (May-Nove. 1958 1588 6 (May-Octo. 1959 1084 7 (May-Nove. 1960 1495 9 (April-De. 1961 1167 6 (June-Nov. 1962 1600 6 (May-Octo. 1963 1822 8 (April-No. 1964 1346 8 (April-No. 1965 1549 7 (May-Nove. 1966 1510 8 (May-Dece. 1967 1515 6 (May-Octo. 1968 1347 7 (May-Nove. 1969 2284 10 (Februar 1970 1555 7 (May ? -N. 1971 938 7 (April-Oc. 1973 1424 8 (April-No. 1974 1272 6 (May-Nove. 1975 1272 6 (May-Nove. 1	0 mm
1955 1829 8 (April-No 1956 1453 6 (May-Octo) 1957 1152 7 (May-Nove) 1958 1588 6 (May-Octo) 1959 1084 7 (May-Nove) 1960 1495 9 (April-De) 1961 1167 6 (June-Nove) 1962 1600 6 (May-Octo) 1963 1822 8 (April-Nove) 1964 1346 8 (April-Nove) 1965 1549 7 (May-Nove) 1966 1510 8 (May-Dece) 1967 1515 6 (May-Octo) 1968 1347 7 (May-Nove) 1969 2284 10 (Februar) 1970 1555 7 (May ? -Nove) 1971 938 7 (April-Octo) 1973 1424 8 (April-Nove) 1974 1272 6 (May-Nove)	vember)
1956 1453 6 (May-Octo) 1957 1152 7 (May-Nove) 1958 1588 6 (May-Octo) 1959 1084 7 (May-Nove) 1960 1495 9 (April-De) 1961 1167 6 (June-Nov) 1962 1600 6 (May-Octo) 1963 1822 8 (April-No) 1964 1346 8 (April-No) 1965 1549 7 (May-Nove) 1966 1510 8 (May-Octo) 1967 1515 6 (May-Octo) 1968 1347 7 (May-Nove) 1969 2284 10 (Februar) 1970 1555 7 (May ? -N) 1971 938 7 (April-Oc) 1973 1424 8 (April-No) 1974 1272 6 (May-Nove)	vember)
1957 1152 7 (May-Nover 1958 1588 6 (May-Octo) 1959 1084 7 (May-Nover 1960 1495 9 (April-De 1961 1167 6 (June-Nover 1962 1600 6 (May-Octo) 1963 1822 8 (April-Nover 1964 1346 8 (April-Nover 1965 1549 7 (May-Nover 1966 1510 8 (May-Decer 1967 1515 6 (May-Octo) 1968 1347 7 (May-Nover 1969 2284 10 (Februar 1970 1555 7 (May ? -Nover 1971 938 7 (April-Octo) 1973 1424 8 (April-Nover 1973 1424 8 (April-Nover 1974 1272 6 (May-Nover 1975 1975 1975 1272 6 (May-Nover 1975 1975 1975 1975 1975 1975 1975 1975	vember)
1958	ber)
1959 1084 7 (May-Nover 1960 1495 9 (April-De 1961 1167 6 (June-Nover 1962 1600 6 (May-Octo) 1963 1822 8 (April-Nover 1964 1346 8 (April-Nover 1965 1549 7 (May-Nover 1966 1510 8 (May-Decer 1967 1515 6 (May-Octo) 1968 1347 7 (May-Nover 1969 2284 10 (Februar 1970 1555 7 (May ? -Nover 1971 938 7 (April ? 1972 888 7 (April-Octo) 1973 1424 8 (April-Nover 1974 1272 6 (May-Nover 1974 1272 6 (May	mber)
1960 1495 9 (April-De 1961 1167 6 (June-Nove 1962 1600 6 (May-Octo) 1963 1822 8 (April-Nove 1964 1346 8 (April-Nove 1965 1549 7 (May-Nove 1966 1510 8 (May-Dece 1967 1515 6 (May-Octo) 1968 1347 7 (May-Nove 1969 2284 10 (Februar 1970 1555 7 (April ? 1971 938 7 (April ? 1972 888 7 (April-Oc 1973 1424 8 (April-Nove) 1974 1272 6 (May-Nove)	ber)
1961 1167 6 (June-Nove 1962 1600 6 (May-Octo) 1963 1822 8 (April-Nove 1964 1346 8 (April-Nove 1965 1549 7 (May-Nove 1966 1510 8 (May-Deceded) 1967 1515 6 (May-Octo) 1968 1347 7 (May-Nove 1969 2284 10 (Februar 1970 1555 7 (May ? -Nove 1971 938 7 (April ? 1972 888 7 (April-Oc 1973 1424 8 (April-Nove 1974 1272 6 (May-Nove	mber)
1962 1600 6 (May-Octo) 1963 1822 8 (April-Not) 1964 1346 8 (April-Not) 1965 1549 7 (May-Nove) 1966 1510 8 (May-Dece) 1967 1515 6 (May-Octo) 1968 1347 7 (May-Nove) 1969 2284 10 (Februar) 1970 1555 7 (May ? -Nove) 1971 938 7 (April ? Nove) 1972 888 7 (April-Octo) 1973 1424 8 (April-Nove) 1974 1272 6 (May-Nove)	cember)
1963 1822 8 (April-No. 1964 1346 8 (April-No. 1965 1549 7 (May-Nove. 1966 1510 8 (May-Dece. 1967 1515 6 (May-Octo. 1968 1347 7 (May-Nove. 1969 2284 10 (Februar. 1970 1555 7 (May ? -No. 1971 938 7 (April ? 1972 888 7 (April-Oc. 1973 1424 8 (April-No. 1974 1272 6 (May-Nove.	ember)
1964 1346 8 (April-Novement 1965 1549 7 (May-Novement 1966 1510 8 (May-December 1967 1515 6 (May-October 1968 1347 7 (May-Novement 1969 2284 10 (Februar 1970 1555 7 (May ? -Novement 1971 938 7 (April ? 1972 888 7 (April-October 1973 1424 8 (April-Novement 1974 1272 6 (May-Novement 1975 1549 1549 1549 1549 1549 1549 1549 154	ber)
1965 1549 7 (May-Nover 1966 1510 8 (May-Decer 1967 1515 6 (May-Octo) 1968 1347 7 (May-Nover 1969 2284 10 (Februar 1970 1555 7 (May ? -Nover 1971 938 7 (April ? 1972 888 7 (April-Octo) 1973 1424 8 (April-Nover 1974 1272 6 (May-Nover	vember)
1966 1510 8 (May-Decended) 1967 1515 6 (May-Octo) 1968 1347 7 (May-Nove) 1969 2284 10 (Februar) 1970 1555 7 (May? -Nove) 1971 938 7 (April ? 1972 888 7 (April-Octo) 1973 1424 8 (April-Nove) 1974 1272 6 (May-Nove)	vember)
1967 1515 6 (May-October) 1968 1347 7 (May-Nove) 1969 2284 10 (Februar) 1970 1555 7 (May ? -Nove) 1971 938 7 (April ? 1972 888 7 (April-October) 1973 1424 8 (April-Nove) 1974 1272 6 (May-Nove)	mber)
1968 1347 7 (May-Nove 1969 2284 10 (Februar 1970 1555 7 (May ? -Nove 1971 938 7 (April ? 1972 888 7 (April-Oct 1973 1424 8 (April-Nove 1974 1272 6 (May-Nove	mber)
1969 2284 10 (Februar) 1970 1555 7 (May ? -No.) 1971 938 7 (April ?) 1972 888 7 (April - Oc.) 1973 1424 8 (April - No.) 1974 1272 6 (May-Nove)	ber)
1970 1555 7 (May ? -N. 1971 938 7 (April ? 1972 888 7 (April-Oc. 1973 1424 8 (April-No. 1974 1272 6 (May-Nove)	mber)
1971 938 7 (April ? 1972 888 7 (April-Oc 1973 1424 8 (April-No 1974 1272 6 (May-Nove)	y-November)
1972 888 7 (April-Oct 1973 1424 8 (April-Not 1974 1272 6 (May-Nove)	ovember)
1973 1424 8 (April-No. 1974 1272 6 (May-Nover	-October)
1974 1272 6 (May-Nove	tober)
	vember)
1075 1250 7 West North	mber)
1975 1350 7 (May-Nove	mber)
1976 1743 8 (April-No	vember)
1977 ca. 1070 7 (May-Nove	mber)

Troth, 1979.

Little variation in the temperature is observed annually (Figure 2-3). However, the greatest diurnal differences as well as the maximum and minimum annual temperatures occur during the dry season (Troth, 1979).


The photoperiod at this latitude is relatively consistent. A 12-hr daylight period is present year-round. However, there is an approximate 0.5-hr shift in dawn and dusk between seasons.

Vegetation Types

The vegetation found in the llanos is well adapted to the cyclic availability of water. The vegetational types in the savanna of Hato Masaguaral have been described as Medano (sandhill), Banco (non-flooded, low ridge), Bajio (moderately flooded savanna), and Estero (more deeply flooded savanna) (Troth, 1979). Additionally, Troth (1979) categorized gallery forest as the fifth major habitat found on the ranch. The study site included three of these five habitats.

Banco. These areas consist of low ridges which follow the course of old waterways. Troth (1979) stated that "the soils are composed of heavy particles. These areas are never truly flooded, but shallow puddles can be found there during the height of the rainy season. The woody vegetation is very open and is dominated by the common palm (Copernicia tectorum)." Associated with many of these palms are strangler figs (Ficus spp.)

Bajio. This is the most common vegetative zone found on the ranch. The woody vegetation of this moderately flooded savanna is dominated by the common palm and associated



strangler figs. Additionally, several shrub species are found in this zone. Clusters of woody vegetation are found in the bajio forming "islands" within the savanna and are known as matas.

The vegetation of the shrub woodland bajío forms a shrub cover which can be so extensive that the canopy can be 90-100 percent closed. The tree layer of this habitat type is 8-15 m high and the shrub layer is 2-10 m high (Troth, 1979). Several species of herbaceous vines and lianas are found in this region.

Estero. This is the more deeply flooded habitat zone. This region is lacking in woody vegetation, with the exception of an occasional palm (Copernicia tectorum) or shrub (e.g., Annona janhii). The water depth during the peak of the wet season varies from 20-60 cm but may reach a depth of 110 cm (Troth, 1979).

Plant Identification

The vascular plants found within the perimeter of Hato Masaguaral have been previously cataloged (Troth, 1979). However, the study site was located in an area of the ranch which had not been surveyed. Consequently, some species which had not been included in the initial surveys were identified. Numerous sources were utilized, many in combination, in the identification of these plants, including popular texts (Hargreaves and Hargreaves, 1960; Hargreaves and Hargreaves, 1965), keys (Ramia, 1974) and personal communication with colleges on-site and local residents.

Behavioral Observations


Studies of arboreal primates are often inhibited by the dense vegetation in which the animals are found. Keeping

focal animals or other members of the study group continuously in view is difficult, if not impossible, under most field conditions. Investigators studying free-ranging animals have attempted to solve this problem by using a variety of sampling techniques (Chalmers, 1968; Chivers, 1969; Aldrich-Blake, 1970; Richard, 1970; Beck and Tuttle, 1972; Glander, 1975).

The habitat/vegetation types found within the area of this study were integral to the success of the behavioral observations. There was little difficultly in keeping the focal subject in full view due to the relatively low canopy and the absence of a vegetation zone at mid-story, allowing an unobstructed view.

These field observations were conducted from August 1987 through December 1988. The first 2 mo were spent locating a suitable study site, selecting two study groups, identifying plant species within the study site, and finalizing the methodology. The territories of the two study groups were adjacent to each other, and the habitat types found in both areas were essentially identical in composition. Figure 2-4 describes the study site and territories of the two study groups.

In an attempt to decrease variability, the two groups of red howlers selected for the project were of similar size and composition (e.g., sex ratio, age ratio). Further, each group possessed a single adult male, which was an additional



Study site and estimates of study group (MSE, LM1) territories. Figure 2-4.

selection factor necessary for the second phase of this project. The animals were identified by using existing physical characteristics, scars, and missing digits. The composition of each group, MSE and LM1, is given in Table 2-2. On July 6, 1988, a male infant (\$\sigma\$3) was born into group LM1. A subadult/adult male (\$\sigma\$2) left group LM1 in October, 1988.

Table 2-2. Demographic composition of two red howler monkey study groups.

Group ID	Animal ID	Age class ¹	Specific marks
MSE	ď	Adult	
	91	Adult	
	92	Adult	
	ďjuv	Juvenile	
	₽inf	Infant	
	đinf	Infant	
LM1	♂ 1	Adult	Cut on lower R lip
	♂ 2	Sub-Adult	
	91	Adult	Missing upper incisor
	₽2	Adult	Scar on upper R lip; notch in R ear
	9 3	Juvenile	
	₽4	Infant	₽1's infant
	ď3	Infant	92's infant

Age class determined upon initiation of study or upon birth of an individual.

During the first 2 mo of the study, the animals were

gradually habituated to the presence of an observer. Since the ranch was closed to hunting, and contact with humans (ranch workers) was fairly regular, the habituation process was quite rapid. A similar response of red howlers to human presence was reported by Racenis (1952). Both study groups were habituated to approaches within 5 m on foot by the onset of formal observations in October 1987. Toward the end of the study, several individual animals would allow observers to approach within 2 m.

An observation period was established during the first 2 wk of each mo. The animals were observed in a semi-random manner which included watching each animal once daily during a different time block. At the end of the month's observation period, a composite day could be constructed for each animal from the time blocks. The animals were randomized again each mo. Table 2-3 provides an example of a typical month's observation schedule for a group. The groups were located through late afternoon census surveys the day immediately preceding the observation series. Each animal was observed for 12 h each mo, from approximately 0630 to 1830 h. corresponded to the entire daylight period and varied slightly with seasonal changes in dawn/dusk. This schedule helped to decrease the bias created by days of continuous rain, during which the animals were relatively inactive.

The animals selected a specific tree in which to sleep

Table 2-3. Example of a single month's observation schedule

for MSE group.

101 1101	groupt					
Time	MON	TUE	WED	THR	FRI	SAT
06:30- 08:30	¥2	ď	♂inf	¥1	ďjuv	\\ \text{Pinf}
08:30- 10:30	ďjuv	¥1	₽2	<i>ਹ</i> "	₽inf	d inf
10:30- 12:30	ď	₽inf	Ŷ 1	ďjuv	♂inf	\$ 2
12:30- 14:30	9inf	9 2	ďjuv	dinf	Ŷ 1	ਰ
14:30- 16:30	9 1	dinf	₽inf	9 2	<i>ਹ</i> *	♂juv
16:30- 18:30	₫inf	ďjuv	ď	\\ \text{pinf}	\$2	91

each evening. The observer left this site after dusk, and observations were initiated at the same site the following morning. Overnight observations were conducted periodically to determine if the animals left the sleeping tree during the night-time hours.

The MSE group was observed monthly from October 1987 to February 1988. This group was captured for the second phase of the project in mid-March 1988. The LM1 group was observed monthly from October 1987 to December 1988.

Focal animal sampling (Altmann, 1974) was employed during all observations, except for two periods which will be discussed later. A total of five major activity categories were observed. These categories were defined as follows:

Feeding, during which an animal was actively consuming foods; this time may have included some mastication if the animal continued to consume food items.

Resting included both sleeping and resting behavior.

Moving was used to indicate traveling from point to point. No attempt was made to separate traveling from changes in position. A change in position within a space twice the animal's body length was not recorded as a "move".

Other provided a broad category which included drinking, howling, scent marking, urination, defecation, play, nursing, and other extraneous activities.

NIV (not in view) was used when the animal had moved out of the field of vision of the observer or the activity could not be defined.

Feeding behavior was the primary activity of interest in this study. No attempt was made to distinguish between feeding and foraging time (searching for food) during most of the sampling period. During the feeding time, specific information was collected concerning the food item selected. The plant species consumed was recorded. If the plant was not identifiable, the individual plant consumed was tagged and numbered for later identification. The part of the plant selected was recorded, as was the phenologic stage of the plant part. The estimation of stage of growth included the following definitions:

Mature leaves included all of the leaf material which had reached the mature size and coloration characteristic of the plant.

Young leaves were those which had nearly reached

(~80% of mature) the characteristic size of the species but not the color. This was the stage prior to maturation.

New leaves were both smaller and lighter in color than the leaves characteristic of the mature plant. These were typically leaves which had recently erupted from the bud stage.

Leaf buds were those leaves which were still in the bud stage.

Flowers included the entire portion of the flower including the petals and sepals.

Flower buds were those flowers that had not yet opened.

Ripe fruits were those fruits which had reached maturity.

Unripe fruits included those fruits which had not yet reached maturity.

Fruit was a general term used when the maturity of the fruit was unknown or could not be identified.

Stem was used to define the woody portion of a plant, typically near the apex of the branch.

Bark included the cambium layer.

Petiole was the structural portion of the leaf that supports the blade of the foliage.

Pulvinus was that swollen portion of the petiole at which the leaf was attached to the stem or branch of the plant.

In addition to the plant part selected and the time spent feeding on a specific plant part, the numbers of bites taken of the plant part were recorded. In some cases, depending on the plant part (e.g., fruits, flowers), the numbers of individual parts consumed were indicated.

Play activity was defined as wrestling between individuals. Also, some solitary play was observed, typically by infants or juveniles interacting with plants. Animals were considered to be nursing if the female's nipple was in the infant's mouth. An example of the field notes collected during a sampling period is provided in Table 2-4.

Two series of overnight observations were conducted, one in each season, to determine if the howlers were active after dusk. The sampling period was initiated after dusk, at approximately 18:30, and was terminated at dawn, at approximately 06:30 the following day. These observations were integrated into the monthly observations, creating two periods during which the animals were under continuous observation for 36 hr. During the overnight observations, a scan sampling of the entire group was conducted every 10 min.

A digital wrist watch was used to time the duration of specific activities. The length of an activity was rounded to the nearest 0.5 min.

Plant Sample Collection and Analysis

Plant samples were collected monthly following the completion of behavioral observations. The species and plant parts collected were based upon the behavioral results of that mo, with emphasis placed upon obtaining samples of those plant parts comprising the top 90% of feeding time. If a

Table 2-4. Example of field notes used for recording behavioral observations of a focal animal in LM1 group.

ODSELAGET	ons of a focal animal in the	MI grou	ν
Activity	Plant species	Bites	Part
FEED	Marsdenia undulata	6	Leaf
REST			
MOVE			
FEED	Pithecellobium tortum	6	Fruit
MOVE			
FEED	Pithecellobium tortum	9	Fruit
MOVE			
FEED	Pithecellobium tortum	2	Leaf
FEED	Marsdenia undulata	3	Leaf
MOVE			
FEED	Platymiscium diadelphum	45	Leaf
MOVE			
FEED	Pithecellobium tortum	3	Leaf
REST			
MOVE			
FEED	Marsdenia undulata	4	Leaf
REST			
FEED	Marsdenia undulata	22	Leaf
MOVE			
FEED	Marsdenia undulata	5	Leaf
MOVE			
FEED	Pithecellobium tortum	6	Leaf
MOVE			
FEED	Pithecellobium tortum	65	Leaf
MOVE			
	Activity FEED REST MOVE FEED REST MOVE FEED REST FEED MOVE FEED MOVE FEED MOVE FEED MOVE FEED MOVE	FEED Marsdenia undulata REST MOVE FEED Pithecellobium tortum MOVE FEED Pithecellobium tortum MOVE FEED Pithecellobium tortum FEED Marsdenia undulata MOVE FEED Pithecellobium tortum MOVE FEED Pithecellobium tortum REST MOVE FEED Marsdenia undulata REST FEED Marsdenia undulata MOVE FEED Marsdenia undulata REST FEED Marsdenia undulata MOVE FEED Pithecellobium tortum MOVE FEED Pithecellobium tortum MOVE FEED Pithecellobium tortum MOVE FEED Pithecellobium tortum	Activity Plant species Bites FEED Marsdenia undulata 6 REST MOVE FEED Pithecellobium tortum 6 MOVE FEED Pithecellobium tortum 9 MOVE FEED Pithecellobium tortum 2 FEED Marsdenia undulata 3 MOVE FEED Platymiscium diadelphum 45 MOVE FEED Pithecellobium tortum 3 REST MOVE FEED Marsdenia undulata 4 REST FEED Marsdenia undulata 4 REST FEED Marsdenia undulata 5 MOVE FEED Pithecellobium tortum 6 MOVE FEED Pithecellobium tortum 6 MOVE FEED Pithecellobium tortum 6 MOVE

representative sample could not be obtained for that mo, then at least one sample for either season (wet, dry) was collected.

There are several sources of variability inherent in determination of food selection by behavioral observation. is impossible to obtain the specific plant part consumed by the animals. This may introduce uncertainty as to how representative the collected plant parts were of the plant parts consumed. Although a part may remain on a plant, which then can be collected for analysis, that part may have been discriminated against by the animal for reasons not readily apparent to the person collecting the plant part. An additional concern was the impact of sample collection on the availability of foods for the study groups, and the effect this may have on the following month's observations. For this reason, the sample collection occurred after completion of the monthly behavioral observations, with approximately 3 wk to permit regrowth before the next observation period.

Every attempt was made to collect samples from the study site or, if not possible, the area immediately surrounding the two territories. A sample similar to the consumed portion of the plant was collected with a targeted minimum sample size of 200 g. Samples were sealed in plastic bags to prevent moisture loss and delivered to the field laboratory. Fresh samples were weighed and dried in a forced-air drying oven at

60°C until reaching a constant weight (i.e., same weight over three consecutive measurements). The dried samples were then placed in a plastic bag, sealed, and shipped to the Comparative Nutrition Laboratory at Michigan State University (East Lansing, MI) for analysis.

Dried samples were ground in a Wiley mill fitted with a 2-mm screen. Samples were analyzed for dry matter (DM), crude protein ([N]x6.25) (CP), ether extract (EE), ash, gross energy (GE), neutral-detergent fiber (NDF), acid-detergent fiber (ADF), cellulose (C), acid-detergent lignin (ADL), Ca, P, Mg, Na, K, Fe, Cu, Mn, Se, Zn. All nutrients were determined using AOAC official methods (AOAC, 1990). Gross energy was determined by complete combustion of an approximately 1-g sample in a Parr adiabatic oxygen bomb calorimeter (Parr Instrument Co., Moline, IL). Fiber components (NDF, ADF, ADL, and C) were determined by using nonsequential detergent fiber fractionation as described by Goering and Van Soest (1970). An approximately 1-g subsample was used for the procedure. All proximate and fiber analyses were performed in duplicate, unless small sample sample size dictated otherwise.

Mineral concentrations were determined using dried samples digested with nitric-perchloric acid (Fenton and Fenton, 1979). Concentrated nitric acid (20.0 ml) and 70% perchloric acid (5.0 ml) were added to approximately 1 g of ground sample in a Phillip's beaker. The sample was digested

until 1 ml of acid remained in the beaker. Following digestion, the samples were diluted with ddH₂O. All samples were digested in duplicate. All mineral concentrations, except P and Se, were analyzed using an atomic absorption/emission spectrophotometer (Smith-Hieftje 4000, Thermo-Jarrell Ash). Phosphorus was measured by colorometric spectrophotometry. Selenium was determined using the sensitive fluorometric assay of Whetter and Ullrey (1978).

Mean weights were determined for selected parts of plant species consumed by red howlers. These individual weights were determined using part or all of the plant samples collected for nutrient analysis.

Intake Estimation

Six mos (March, 1988) after the initial field observations began, one of the study groups (MSE) was selected for transfer to holding facilities on the ranch for more intensive study. These animals were chemically immobilized with a dart gun using Telazol® (tiletamine/zolazepam) at a concentration of 200 mg/ml. A mean dose of 23.3 mg/kg was used, with a range from 13.0 mg/kg to 37.5 mg/kg (Crissey and Edwards, 1989). Immobilized animals were caught in a large blanket, as they fell from the trees, to prevent injury from impact with the ground. The juvenile male (Juv) was the first individual immobilized. It was felt that this animal

was the most likely to flee when the capture procedure was initiated. The adult females were then immobilized. When the first female (\$2) was immobilized, her male offspring (dinf) also fell, clinging to her. Recognizing this as the ideal method of capturing the infant animals, it was hoped that the second infant (\$\frac{2}{3}\$ inf) would also cling to her mother during immobilization. This, unfortunately, was not the case, as the female infant (\$\frac{2}{3}\$ inf) jumped free of her mother prior to her mother's fall into the blanket. This necessitated the immobilization of the female infant (\$\frac{2}{3}\$ inf), which was conducted without incident. The adult male (\$\frac{2}{3}\$) was the final animal of the group to be immobilized and transferred to the holding facilities.

In addition to the animals in the MSE study group, 7 animals from 2 groups peripheral to the MSE home range were immobilized following the guidelines above. Anatomical data (e.g., body weight, linear measurements) was collected, the animals were fitted with ear tags, and then released.

The holding facilities were located near the principal living area at Hato Masaguaral, which was approximately 2 km from the study site. The holding facilities were designed specifically to house red howlers. The sides of the enclosure, chain link mesh, were all open to the surrounding environment, although a solid roof covered the entire structure. A perimeter fence (~2 m) surrounded the enclosure

to prohibit feral dogs and pigs from entering.

The enclosure was 7.8 m long x 5.75 m high x 4.08 m deep. The floor of the enclosure was 1.0 m above a concrete slab, allowing easy access to the floor beneath for cleaning. The floor was covered by metal pans that could be removed for cleaning or collection of samples without entry into the cage.

The total structure was divided into five sections, four of which were equal in size (1.3 m x 5.75 m x 4.08 m) with the fifth twice as wide as the other four (2.6 m x 5.75 m x 4.08 m). The sections were interconnected with shift doors located 3.05 m above the floor of the enclosure. Divisions between sections were made of the same wire mesh as the outer sides, allowing visual, olfactory, and vocal communication when the animals were physically separated.

The interior of each section was furnished with numerous tree limbs, branches, and ropes to provide climbing structures. At the front of each section was a wooden shelf $(1.3 \text{ m } \times 0.5 \text{ m})$ located 3.05 m above the floor of the enclosure. A graduated container for drinking water was placed upon this shelf. Additionally, there was a hanging platform $(0.75 \text{ m } \times 0.75 \text{ m})$ in the center of each section. The platform in the largest of the five sections had a spring scale attached to it, creating a means by which the animals could be regularly weighed.

The enclosure flooring and hanging platforms were cleaned

daily. The animals were group-fed three times daily. The first feeding consisted of a "higher" fiber primate diet.

Selected nutrient specifications of this extruded diet are provided in Table 2-5. The primate diet was placed in pans on the hanging platforms. This helped entice the animals to use the hanging platforms frequently and thus facilitated the weighing of individuals. The two remaining daily feedings were of fresh browse which was collected from the surrounding area immediately prior to feeding.

Table 2-5. Selected nutrient composition of a "higher" fiber primate diet fed to captive red howlers in Venezuela (dry matter basis).

Nutrient	Concentration (%)
Dry matter	92
Crude protein	25
Ether extract	4.9
Crude fiber	10.9
Acid-detergent fiber	14.7-17.4

^{*}Marion Zoological, Inc., Plymouth, MN.

Water consumption was monitored daily. A container of equal size and volume, placed in an area inaccessible to the howlers, also was monitored to determine daily evaporation rates. This correction factor was needed to accurately determine water intake by the animals. All water containers were replenished daily.


The adult animals in the group (one σ , two \$\$) were used

to determine the mean bite weight of specific food items selected by animals observed in the wild. A pre-weighed portion of each food item was offered to one of the adult animals which had been temporarily isolated from the remainder of the group. The number of bites taken from each portion were recorded, and the remaining portion of the sample was re-weighed. Concurrently, a representative sample of the same food item was weighed, handled in the same manner as the other samples, and re-weighed following the procedure. This sample was used as a control to correct for any desiccation that may have occurred during the procedure.

Daily intake of a nutritionally complete, "higher" fiber extruded primate diet by three (200, 19) of the howlers was measured for 9 d. Weighed quantities of the higher fiber primate diet were offered each morning. Orts were removed in the late afternoon (~18:00), weighed and discarded.

Nutrient Composition of Consumed Diet

By integrating data from the above three areas of study, estimates of the nutrient composition of the diet consumed by free-ranging red howlers were made. For example, a free-ranging, adult of red howler was observed to take 131 bites of Pithecellobium saman leaves. The mean bite weight of leaves of that plant species was 0.4 g as determined with trials involving the captive, adult of red howler. Therefore, the



consumed weight of 131 bites of *Pithecellobium saman* leaves was estimated to be 52.4 g (as fed basis). Selected nutrient analyses of *Pithecellobium saman* leaves indicated that the plant contained 38.3% dry matter and 5.61 kcal GE/g (DMB). Thus, within the 52.4 g of consumed leaf material, the animal had consumed 20.1 g of dry matter (32.3 g of moisture) and 112.8 kcal of GE.

Similar calculations were made from the annual behavioral observations for the three adult animals (15, 299) in LM1 to estimate nutrient composition of the consumed diet.

Results

Behavioral Observations

The behavioral data for the adult animals (10, 299) of group MSE consisted of 180 focal-animal h collected from October, 1987 through February, 1988. This data set was not divided according to season (dry versus wet). Time budgets within the 12-h dawn to dusk observation period are provided in Table 2-6.

The behavioral data for group LM1 consisted of 912 focal animal h. Adult animals (10, 299) of the group comprised 432 focal-animal h of that total time, collected during 12 consecutive mos from January 1988 through December 1988. The 1988 dry season data included observations from January through May (5 mos). Wet season data included

Table	2-6.	Distribution	(%)	of	daily	activities	of	three
adult	(1ď.	299) red howler	s (rou	m MSE)	_		

Activity	Time (%)	Time (hr)
Rest	76.6	9.2
Feed	13.1	1.6
Move	8.2	1.0
Other	1.5	0.2
NIV	0.6	0.1

observations made from June through November (6 mos).

December 1988 was a transition mo between wet and dry season, and was not included in the results. The distribution of daily activities expressed on an annual, dry season, and wet season basis is provided in Table 2-7.

The distribution of time spent feeding (group MSE) on major plant parts (leaf, fruit, flower) is given in Figure 2-5. Consumption data, divided among more specific food plant parts, are shown in Figure 2-6.

Table 2-7. Distribution of daily activities of three adult $(1\sigma, 299)$ red howlers (group LM1) expressed on an annual, dry season, and wet season basis.

Activity	Ann	ual	Dry S	Season	Wet S	Season
	min	8	min	8	min	ક
Rest	543.0	75.4	540.7	75.1	542.9	75.4
Feed	90.7	12.6	88.6	12.3	96.5	13.4
Move	54.0	7.5	62.6	8.7	14.4	6.2
Other	28.8	4.0	24.5	3.4	31.7	4.4
NIV	3.6	0.5	2.9	0.4	4.3	0.6

Figure 2-5. Major plant parts (leaf, fruit, flower) consumed by red howlers (group MSE), listed as percent of observed feeding time.





Figure 2-6. Specific plant parts consumed by red howlers (group MSE), listed as percent of observed feeding time.



The distribution of feeding time (group LM1) on general plant parts (leaf, fruit, flower) is presented in Figure 2-7 (annual), Figure 2-8 (dry season), and Figure 2-9 (wet season). The distribution of time spent feeding on plant parts, at specific stages of maturity (e.g., ripe, unripe), is presented in Figures 2-10, 2-11, 2-12.

Food plant species and parts consumed by group LM1, ranked by total annual feeding time, dry season feeding time, and wet season feeding time, are listed in Tables 2-8, 2-9, and 2-10, respectively.

Throughout the entire study, the howlers were never observed to consume animal-based foods. This does not exclude the possibility of secondary ingestion of insects which may be associated with food plants during some stage of their lifecycle.

The two 12-h dusk (18:00) to dawn (06:00) observation periods revealed that this group of red howlers was not active during that time period, with "resting" comprising 100% of the night time scan samples.

Plant Sample Collection and Analysis

A total of 171 individual plant samples was collected during the study period, representing parts from 36 species.

Dry matter, crude protein, ether extract, ash, and fiber component (NDF, ADF, ADL, C) concentrations for these samples

Figure 2-7. Major plant parts (leaf, fruit, flower) consumed by red howlers (group IM1), listed as percent of total annual feeding time.




Figure 2-8. Major plant parts (leaf, fruit, flower) consumed by red howlers (group LMI), listed as percent of total dry season feeding time.




Figure 2-9. Major plant parts (leaf, fruit, flower) consumed by red howlers (group IM1), listed as percent of total wet season feeding time.





Figure 2-10. Specific plant parts consumed by red howlers (group LMI), listed as percent of total annual feeding time.



Specific plant parts consumed by red howlers (group LMI), listed as percent Figure 2-11. Specific plant parts of total dry season feeding time.





Figure 2-12. Specific plant parts consumed by red howlers (group LMI), listed as percent of total wet season feeding time.



Food plant species and part consumed by red howlers (group LM1), ranked by total 1.99 1.93 5.75 7.38 6.45 5.32 5.30 4.94 4.68 4.54 4.31 3.38 2.69 2.42 2.38 2.05 Feeding time (% of total) 88.0 79.0 188.0 173.5 148.5 78.0 67.0 65.0 63.0 241.5 174.0 161.5 153.0 141.0 110.5 Feeding time 211.0 (mim) Species no.b (81) (21)(87) (34) (74) (64) (46)(41) (62) (84) (48) (65) (26)(22) (4) Parta FLBU NL FL RF RF UF FL E П П Н H Ŀ Н П Н Pithecellobium tortum Pithecellobium saman Pithecellobium saman Albizia guachapele Licania pyrifolia Licania pyrifolia Sterculia apetala Guazuma ulmifolia Guazuma ulmifolia Mangifera indica Unidentified #01 Cassia grandis Zizyphys saeri species Ficus pertusa Ficus pertusa Ficus pertusa feeding time. Plant Table 2-8. annual Rank 10 11 12 13 14 3 2 9 7 ω 0 2 4

Rank 17 18 19 20 21 22 23 24 25 26 27 28 29 30

rable 2-8 by total	Table 2-8 (conf'd). Food plant species and part consumed by red nowlers (group LM1), ranked by total annual feeding time.	n part	consumed by	red howlers (gr	oup LMI), ranked
Rank	Plant species	Parta	Species no.b	Feeding time (min)	Feeding time (% of total)
33	Pithecellobium dualense	FL	(40)	25.5	0.78
34	Unidentified #03	L	(91)	25.5	0.78
35	Sterculia apetala	ST	1	21.5	0.78
36	Perskia guamacho	Ξı	(19)	19.0	0.58
37	Guazuma ulmifolia	NE	1	18.0	0.55
38	Pithecellobium saman	YL	(44)	18.0	0.55
39	Annona jahnii	L	(9)	17.0	0.52
40	Unidentified #05	L	1	17.0	0.52
41	Pithecellobium dualense	L	(39)	16.5	0.50
42	Annona punicifolia	RF	(11)	16.0	0.49
43	Albizia guachapele	YL	(36)	15.5	0.47
44	Pithecellobium saman	ш	(47)	15.5	0.47
45	Cassia grandis	Ĺτι	(27)	15.0	0.46
46	Sapium aucuparium	I	(25)	15.0	0.46
47	Albizia guachapele	NL	(37)	12.5	0.52
48	Pithecellobium saman	LBU	(43)	12.0	0.37
49	Paullinia cururu	ı	(80)	11.5	0,35

Table 2-8 (cont'd). Food plant species and part consumed by red howlers (group LM1), ranked 0.15 0.32 0.32 0.31 0.31 0.29 0.29 0.29 0.24 0.24 0.23 0.21 0.20 0.20 0.18 0.17 0.15 time (% of total) Feeding 10.01 time 9.5 9.5 9.5 8.0 8.0 7.5 7.0 6.5 6.5 0.9 5.5 5.0 5.0 10.5 10.5 10.0 (min) Feeding Species no.b (43) (98) (53) (52)(45) ı ı ı ı ı ı Parta FLBU PET NL XL N NL YL N FL Ē Н Н Н Ē H Н [± Pithecellobium tortum Pithecellobium tortum by total annual feeding time. Randia venezuelensis Randia venezuelensis Pithecellobium saman Pithecellobium saman saman Albizia columbiana Sterculia apetala Sterculia apetala Licania pyrifolia Unidentified #06 Mangifera indica Mangifera indica Ceiba pentandra Ceiba pentandra Pithecellobium species Ficus pertusa Plant Rank 20 51 52 53 54 55 99 57 58 59 09 61 62 63 64 65 99

Table 2-8 (cont'd). Food plant species and part consumed by red howlers (group LM1), ranked by total annual feeding time.

Rank	Plant species	Parta	Species	Feeding time	Feeding time
67	Cassia grandis	YL	1	5.0	0.15
89	Guazuma ulmifolia	ı	-	5.0	0.15
69	Annona jahnii	UF	(7)	4.5	0.14
0.2	Ceiba pentandra	ST	ī	4.5	0.14
7.1	Ceiba pentandra	PET	1	4.0	0.12
72	Zizyphys saeri	YL	1	4.0	0.12
73	Ficus sp.	ı	1	3.5	0.11
74	Melothria trilobata	ı		3.0	0.09
75	Guazuma ulmifolia	YL	-	3.0	0.09
91	Unidentified #07	IJ	1	3.0	0.09
77	Coccoloba sp.	YL	1	2.5	0.08
78	Genipa americana	FT	1	2.0	90.0
61	Unidentified #08	FLBU	1	2.0	90.0
30	Entada sp.	YL	1	1.5	0.05
31	Unidentified #09	I	1	1.5	0.05
32	Unidentified #10	দ্য	1	1.0	0.03
33	Mangifera indica	FLBU	1	0.5	0.02

Table 2-8 (cont'd). Food plant species and part consumed by red howlers (group LM1), ranked by total annual feeding time.

Flant snow;				
Species	Part	Parta Species	Feeding time	Feeding time
84 Marsdonia	1	no.b	(min)	(% of total)
85 material undulata	[E4	(11)	0.5	0.02
*L=mature leaf. NI	ı	,	0.5	0.02
FL=flower; PET=petiole; PUL=pnlvim.s leaf; F=fruit; UF=unripe fruit; RF=ripe fruit;	leaf; F	=fruit; U	F=unripe fruit;	RF=ripe fruit;
A21.	nutrient	analysis	provided in App	endix Tables A1-
Triant parts ranked from 1 through (†) represent 90.2% of total annual feeding time.	epresent	90.2% of	total annual fe	eding time.

Table 2-9. Food plant species consumed by red howlers (group LMI), ranked by total dry season feeding time.

Rank	Dlant				
	tail Species	Parta	Species	Feeding time	Feeding time
1	Bitt		no.b	(min)	(% of total)
1	fithecellobium saman	FL	(46)	174.0	12.03
7	Mangifera indica	RF	(4)	1/8 7	10 01
3	Guazuma ulmifolia	F	(1/0)	136.0	10.27
4	Guazuma ulmifolia	į (r	(60)	135.5	10.0
5	Albizia guachapele	4 -	(00)	0.011	8.03
9	Licania nurifolia	4	(33)	91.0	67.9
1	Pittolia	NL	(21)	65.0	4.49
	Pithecellobium saman	I	(42)	64.5	4.46
8	Licania pyrifolia	I	(22)	62.0	4.29
6	Sterculia apetala	FLBU	(87)	57.0	3.94
10	Mangifera indica	PUL	(5)	52.0	3.59
11	Cassia moschata	I	(29)	42.0	2.90
12	Zizyphys saeri	I	(69)	34.0	2,35
13	Cassia moschata	YL	(31)	32.5	2.25
14	Unidentified #02	YL	(06)	29.0	2.00
15	Licania pyrifolia	YL	1	25.5	1.76
16	Ficus pertusa	T	(62)	21.5	1.49
17	Mangifera indica	UF	(2)	21.0	1.45

Kank					
	fiant species	Part	Species	Feeding time	
18	Pterocarpus acapulcansis		no.	(min)	(% of total)
19	C+011212	NE	(61)	19.5	1.35
	sterculia apetala	ST	(88)	18.0	1.24
07	Cassia moschata	TI	(32)	0 0	
21	Unidentified #04	-	1201	0.71	01.1
22	Pithecellobium saman			6	1.14
23	Unidentified 403	1	(48)	15.5	1.07
		L	(91)	14.5	1.00
67	Albizia guachapele	NE	(37)	12.5	0.86
25	Pithecellobium saman	LBU	-	12.0	0.83
26†	Unidentified #05	17	-	12.0	0 83
27	Cassia grandis	I	1	11.5	0.80
28	Guazuma ulmifolia	NE	-	11.0	0.76
29	Pithecellobium tortum	Ĺτι	(53)	10.5	0.73
30	Pithecellobium saman	PET	1	10.0	69.0
31	Mangifera indica	ı	-	9.5	99.0
32	Ceiba pentandra	ഥ	ī	9.5	0.66
33	Annona jahnii	L	(9)	8.5	0.59
2.4					

SHITTER					
Pithereller.		Parta	Species	Feeding time	Feeding time
ceiloblum tortum	m tortum	MT		(MID)	(% of total)
Pithecellobium	m tort	NF	(52)	7.5	0.52
Ficus pertusa	mn ron ron	I	1	6.5	0.45
Pitherallob	1	ΛΓ	1	6.5	0.45
Pithogoll	m saman	FLBU	(45)	6.5	0 45
Stores 110bium	m saman	NT	(43)	6.0	0.41
Jerculia ape	apetala	FL	(98)	5.5	0.38
Licania pyrifolia	olia	Ĺτί	1	5.0	7 0 35
Ceiba pentandra	Ira	L	1	4.0	0.28
Ceiba pentandra	Ira	PET	1	4.0	0.28
Zizyphys saeri	i.	YL	1	4.0	0.28
Mangifera ind	indica	NL	1	1.0	0.07
Sapium aucuparium	rium	T	1	1.0	0.07
Coccoloba sp.		YL	1	1.0	0.07
Mangifera indica	lica	FLBU	1	0.5	0.03

PReference number corresponds to sample nutrient analysis provided in Appendix Tables Al-+Plant parts ranked from 1 through (†) represent 90.5% of total dry season feeding time.

rable 2-10. Season feedi Rank Plan	feeding time. Plant Species Consumed by red howlers (group LMI), ranked by total wet	med by red		'Stoup LMI), ran	ked by total wet
11	Ficus pertusa	Parta	Species no.b	Feeding time	Feeding time
	Ficus pertusa	UF	(64)	211.0	(% of total)
-	Ficus pertusa	I	(63)	140.0	13.21
	1 0	RF	(65)	110.5	6.92
1	Albizia quachanala	I	(41)	96.5	6.04
	Pithecellohim +	L	(34)	93.5	5,85
	Zizyphys sacti	T.	(49)	92.0	5.76
	Entada en	[E4	(70)	88.0	5.51
	1	I	(54)	59.5	3.73
0	Ziraentiled #01	I I	(74)	54.0	3.38
	Cassia grandis	17	(26)	53.5	3 35
11	Marsdenia undulata	IJ	(16)	52.0	3.26
12	Copernicia tectorum	Ē	(12)	48.0	3.01
13	Pterocarpus acapulcensis	ı	(60)	44.0	2.76
14	Zanthoxylum culantrillo	YL	(78)	38.5	2.41
15	Zanthoxylum culantrillo	I	(16)	37.0	2.32
16	Pithecellobium dualense	FL	(40)	25.5	1.60
1	Guazuma ulmifolia	Ĺ	(81)	21.0	1.31

Table 2-10 (cont'd). Food plant species consumed by red howlers (group LMI),

SOLDER				
Pereskia	Parta	Species	Feeding time	Feeding time
y y damacho	-		(m1n)	(% of total)
Unidentified #04		(19)	19.0	1.19
Guazuma ulmifolia	I	-	19.0	1.19
Pterocarpus accaus	FI	(82)	17.5	1.10
Pitherallow	NL	(61)	17.5	0-1
dualense dualense	I	(30)		21.1
Annona punicifolia	E c	(66)	16.5	1.03
Albizia quachanala	KF	(11)	16.0	1.00
Cassis are all pere	AL	(36)	15.5	76.0
accera grandis	[zi	(27)	15.0	0
Sapium aucuparium	7	(36)		70.0
Mangifors india		(62)	14.0	0.88
2	UF	(3)	12.5	0.78
29T Paullinia cururu	ı	(80)	11 2	0 72
Unidentified #03	,		0	7/.0
		(37)	11.0	0.69
Kandia venezuelensis	I	1	10.5	0.66
Albizia columbiana	ц	(33)	10.0	0.63
Randia venezuelensis	YL	(71)	9.5	0.59
Annona jahnii	T	(9)	8.5	0.53
4000000	1	(28)	0 8	0 50

rable cotal Rank	Table 2-10 (cont'd). Food plant species Rank Plant species	Consum	led by red	howlers (group	Consumed by red howlers (group LM1), ranked by
36	Unidentie:	Parta	Species	Feeding time	Feeding time
37	Guazima Guazima	I	1011	(min)	(% of total)
38		111		8.0	0.50
2 8	Cassia moschata	NL	-	7.0	0.44
5	Mangifera indica	UF	(32)	6.5	0.41
30	Cassia grandis	NL	1	0.9	0.38
41	Guazuma ulmiforia	YL	,	5.0	0.31
42	Annona interior	П	1	5.0	0.31
43	Ceiha nort	UF	(7)	4.5	0.28
44	Inidontic	ST	1	4.5	0.28
45	E:	YL	1	4.5	0.28
100	ricus sp.	T	1	3.5	0.22
0 5	Sterculia apetala	ST	(88)	3.5	0.22
15	Melothria trilobata	Ţ	(23)	3.0	0.19
48	Guazuma ulmifolia	YL	1	3.0	0.19
49	Unidentified #07	Ţ	,	3.0	0.19
50	Genipa americana	Fi	1	2.0	0.13
51	Unidentified #08	FLBU	1	2.0	0.13
52	Entada sp.	YL	1	1.5	60.0

Table 2-10 (cont'd). Food plant species consumed by red howlers (group IM1), ranked by total wet season feeding time.

Rank	Rank Plant species	Parta	Parta Species	Feeding time (min)	Feeding time (% of total)
53	Unidentified #09	I	1	1.5	0.09
54	Ceiba pentandra	L	1	1.0	90.0
55	Coccoloba sp.	AL	-	1.0	90.0
56	Unidentified #10	H	1	1.0	90.06
57	Tetracera volubris	L	1	0.5	0.03
L=matu fruit; Refere	"I-mature leaf; YI-young leaf; NL-new leaf; LBU=leaf bud; F=fruit; RF=ripe fruit; UF=unripe fruit; T=f=lower; RLBU=flower bud; PET=petlob; PUL-pullyinus; STEM=stem. Reference number corresponds to sample nutrient analysis provided in Appendix Tables Al-	LBU=leg tiole; utrient	af bud; F PUL=pulvi analysis	=fruit; RF=ripe nus; STEM=stem. provided in App	fruit; UF=unripe endix Tables A1-
A21.	421. Plant parts ranked from 1 through (†) respresent 90.2% of total wet season feeding time.	spresent	90.28 0	f total wet seas	on feeding time.

are provided in Appendix Tables A1-A7. Macro-mineral (Ca, P, Na, K, Mg) concentrations are given in Appendix Tables A8-A14. Trace element (Fe, Cu, Mn, Se, Zn) concentrations are given in Appendix Tables A15-A21.

Average weights of specific plant parts, which were used to supplement the bite weight data (see below), are provided in Table 2-11.

Intake Estimation

The average bite weights of an adult male (σ) and two adult females (\$1, \$2) were determined for parts of some of the plants. The plants were those upon which 90% of the feeding time of free-ranging animals was spent, determined by the field observations. Table 2-12 provides the average bite weight for the specific plant parts tested.

The mean body weights of free-ranging red howlers, collected during immobilization procedures, are provided in Table 2-13. Levels of daily consumption of the higher fiber primate diet by three of the captive animals during a 9-d trial are given in Table 2-14.

Nutrient Composition of Consumed Diet

Integrating information from the results of the three areas of study, daily food intake by free-ranging howlers was calculated to be 544.0 g, on an as-fed basis, or 151.0 g of

Table 2-11. Average fresh weight of plant parts consumed by red howlers (Alouatta seniculus) in the central lianos region of Venezuels.

Family/ Species	Species	Part	и	Weight (g)	(g)
	no.			Mean	SD
Anacardiaceae					
Mangifera indica	(2,3)	UF	28	135.24	19.446
Mangifera indica	(4)	RF	54	139.13	29.619
Mangifera indica	(5)	PUL	280	0.13	0.036
Annonaceae					
Annona jahnii	(9)	ī	100	1.11	0.306
Annona jahnii	(7)	UF	7	10.90	1
Annona jahnii	(8)	RF	11	69.50	1
Annona punicifolia	(10)	UF	14	24.90	-
Annona punicifolia	(11)	RF	41	49.64	6.336
Arecaceae					
Copernicia tectorum	(15)	RF	33	6.22	0.790
Asclepidaceae					
Marsdenia undulata	(16)	I	63	0.88	0.455
Marsdenia undulata	(17)	UF	20	127.87	18.020
Cactaceae					
Pereskia quamacho	(19)	R	18	3.94	0.918

Table 2-11 (cont'd). Average fresh weight of plant parts consumed by red howlers (Alouatta seniculus) in the central llanos region of Venezuela.

Family/ Species	Species	Parta	и	Weight	t (g)
	no.			Mean	SD
Dilleniaceae					
Tetracera volubris	(24)	FL	15	0.04	0.074
Euphorbiaceae					
Sapium aucuparium	(25)	Ţ	76	0.51	0.218
Fabaceae (Caesalpinoideae)					
Cassia grandis	(36)	Llft	108	0.62	0.050
Fabaceae (Mimosoideae)					
Pithecellobium saman	(43)	NL	100	0.87	0.550
Pithecellobium saman	(44)	YL	57	3.53	1.229
Pithecellobium saman	(48)	UF	7	19.34	10.057
Pithecellobium tortum	(49)	Lflt	54	0.44	0.197
Pithecellobium tortum	(53)	UF	20	1.79	0.872
Fabaceae (Papilionoideae)					
Platymiscium diadelphum	(56)	Lflt	100	0.42	0.186
Rubiaceae					
Randia venezuelensis	(73)	UF	64	1.10	0.224
Rutaceae					
Zanthoxylum culantrillo	(26)	Lflt	54	0.47	-

Table 2-11 (cont'd). Average fresh weight of plant parts consumed by red howlers (Alouatta seniculus) in the central llanos region of Venezuela.

Family/ Species	Species	Part	и	Weight (g)
	no.			
				Mean SD
Sterculiaceae				

Sterculia apetala 0.26 0.074
* L=mature leaf; Lflt=leaflet; NL=new leaf; YL=young leaf; F=fruit; UF=unripe fruit; RF=ripe fruit; FL=flower;
PET=petiole; PUL=pulvinus

b All fruit weights include seeds unless otherwise indicated.

Table 2-12. Wean bite weight (g) of plants consumed by red howlers (Alouatta seniculus) in the central llanos region of Venezuela.

Family/ species	Species no.	Part	*0	п	41	п	42	п
Anacardiaceae								
Mangifera indica	(1)	YL	0.47	7	0.78	4	1	1
Mangifera indica	(2), (3)	UF	2.19	20	2.03	10	1.81	11
Mangifera indica	4	RF	2.39	11	1.82	22	1.96	21
Annonaceae								
Annona punicifolia	(11)	RF	6.41	11	4.09	10	7.37	11
Arecaceae								
Copernicia tectorum	(15)	RE	0.87	6	1.07	10	1.09	10
Asclepidaceae								
Marsdenia undulata	(16)	L	0.89	10	0.55	11	0.82	10
Marsdenia undulata	(11)	UF	1.20	2	1.49	4	1.33	9
Cactaceae								
Pereskia guamacho	(19)	RF	'	1	4.18	11	3.40	2
Chrysobalanaceae								
Licania pyrifolia	(21)	NL	0.64	11	0.46	10	0.32	10
Licania pyrifolia	(22)	L	0.34	10	0.35	17	0.29	10
Dilleniaceae								
Tetracera volubris	(24)	FL	0.29	σ	0.32	11	0.23	0

Table 2-12 (cont'd). Mean bite weight (g) of plants consumed by red howlers (Alouatta seniculus) in the central llanos region of Venezuela.

Family/ species	Species no.	Part	ъ	п	9.1	п	42	п
Fabaceae (Caesalpinoideae)								
Cassia grandis	(26)	T	0.39	4	0.33	10	0.33	8
Cassia moschata	(28)	T	0.39	2	0.33	13	0.22	2
Cassia moschata	(32)	UF	1	1	1.32	3	1.64	1
Fabaceae (Mimosoideae)								
Albizia guachapele	(34,35)	Г	0.55	12	0.49	12	0.36	10
Albizia guachapele	(36)	YL	0.57	6	0.37	6	0.41	10
Pithecellobium dualense	(39)	ı	0.39	1	1	1	1	,
Pithecellobium dualense	(40)	FL/BU	0.35	9	0.15	80	1	,
Pithecellobium saman	(41)	IJ	0.40	10	0.32	11	0.27	10
Pithecellobium saman	(45)	FLBU	0.93	14	0.81	12	0.67	12
Pithecellobium saman	(46)	FL	0.83	14	0.81	12	0.67	12
Pithecellobium tortum	(49)	ı	0.39	11	0.28	12	0.17	13
Pithecellobium tortum	(52)	BU, NL	-	1	0.18	4		'
Fabaceae (Papilionoideae)								
Entada polystachya	(54)	T	0.46	7	0.40	12	0.27	4
Pterocarpus acapulcensis	(09)	ı	0.58	2	0.51	2	,	1
Pterocarpus acapulcensis	(61)	NE	0.65	15	0.56	10	09.0	15

Table 2-12 (cont'd). Mean bite weight (g) of plants consumed by red howlers (Alouatta seniculus) in the central llanos region of Venezuela.

Family/ species	Species no.	Part*	ъ	п	41	п	4.2	d .
Moraceae								
Ficus pertusa	(62)	I	0.45	10	0.37	10	0.43	10
Ficus pertusa	(64)	UF	0.24	11	0.26	10	0.24	10
Ficus pertusa	(65)	RE	0.27	10	0.30	10	0.34	5
Rubiaceae								
Unidentified #01	(75)	YL	0.31	12	0.16	11	0.36	10
Rutaceae							7.97	
Zanthoxylum culantrillo	(78)	YL	0.40	10	0.34	13	0.30	12
Sapindaceae								
Paullinia cururu	(18)	YL	0.35	10	0.22	10	0.24	10
Sterculiaceae								
Guazuma ulmifolia	(81)	Ĺι	1.03	10	0.67	11	1.02	10
Guazuma ulmifolia	(84)	FL	0.44	14	0.19	12	0.11	9
Sterculia apetala	(87)	FLBU	0.40	10	0.27	10	0.23	10
UNIDENTIFIED PLANTS								

Unidentified #02 7 0.29 14 1- | Unidentified #02 7 0.29 14 | Limature leaf; NL-new leaf; NL-young leaf; F-fruit; UF-unripe fruit; FF-ripe fruit; FL-flower; PET-petiole; PUL=pulvinus

Table 2-13. Mean body weight of free-ranging red howlers (Alouatta seniculus) in the central llanos region of Venezuela.

Age Class/Sex	n	Mean BW (kg) ± SD
Adult dd	4	6.28 ± 0.90
Adult 99	4	4.10 ± 0.37

Table 2-14. Body weight and daily feed intake of nutritionally complete, "higher" fiber primate diet by captive red howlers (Alouatta seniculus).

Animal	BW (kg)	Intake (g/d) (AFB)	Intake (g/d) (DMB)	Intake (% BW) (DMB)
Adult o	6.58	146.1	134.4	2.04
Juvenile &	4.31	101.5	93.4	2.17
Juvenile ?	NDb	100.7	92.6	-

^{*}Marion Zoological, Inc., Plymouth, MN.

dry matter. Dry matter intake, as a percentage of mean body weight, was calculated as 3.13%. Mean gross energy intake was estimated to be 711.2 kcal/d. The calculated concentrations of proximate and fiber components in the diet of free-ranging animals are provided in Table 2-15. The calculated macro and trace mineral concentrations are given in Table 2-16. Estimates of dietary nutrient composition based alone on distribution of time feeding on specific plant parts are presented in Tables 2-17 and 2-18 for proximate and mineral concentrations, respectively.

The distribution of feeding among general plant parts (leaf, fruit, flower) based upon calculated mass intake is

bNot determined.

Table 2-15. Calculated proximate and fiber composition of the mean annual diet selected by three adult $(1\sigma, 299)$ red howlers (group LM1).

Nutrient	Concentration (DMB)
Dry matter (%)	27.8
Crude protein (%)	13.4
Crude fat (ether extract) (%)	3.6
Ash (%)	6.3
Gross energy (kcal/g)	4.71
Neutral-detergent fiber (%)	51.5
Acid-detergent fiber (%)	37.8
Acid-detergent lignin (%)	18.3
Cellulose (%)	19.0

Table 2-16. Calculated macro- and trace-mineral composition of the mean annual diet selected by three adult $(1\sigma, 299)$ red howlers (group LM1).

Nutrient	Concentration (DMB)
Calcium (%)	0.78
Phosphorus (%)	0.19
Sodium (ppm)	97.3
Potassium (%)	1.6
Magnesium (%)	0.35
Iron (ppm)	126.8
Copper (ppm)	11.6
Manganese (ppm)	100.6
Selenium (ppm)	1.04
Zinc (ppm)	19.5

Proximate nutrient composition (dry basis) of total diet as consumed by red Table 2-17.

howlers (group IM1) in the central llanos region of Venezuela calculated based on time spent <u>f</u> eeding.	anos region of Ven	ezuela calculated ba	ased on time spent
Nutrient	Annual	Wet Season	Dry Season
Dry matter (%)	27.1	29.4	27.3
Crude protein (%)	11.7	12.7	11.3
Ether extract (%)	2.7	3.2	2.2
Ash (%)	5.5	5.7	4.6
Gross energy (kcal/g)	4.18	4.30	3.91
Neutral-detergent fiber (%)	47.6	49.1	44.6
Acid-detergent fiber (%)	33.5	17.4	29.7
Acid-detergent lignin (%)	16.1	17.6	13.4 86

15.6

18.0

16.7

Cellulose (%)

Macro and trace mineral composition (dry basis) of total diet as consumed by red howlers (group LM1) in the central llanos region of Venezuela calculated based on time 09.0 0.19 3,16 0.25 0.18 18.0 87.2 1.7 105.2 10.1 59.4 Dry Season 96.0 0.14 0.49 4.39 98.9 52.6 6.6 115.7 2.1 122.5 17.2 Season Wet 0.15 2.39 0.89 5.93 0.38 93.3 1.4 9.7 91.6 17.1 105.1 Annual Calcium: Phosphorus Manganese (ppm) Phosphorus (%) (mdd) spent feeding. Potassium (%) Magnesium (%) Sodium (ppm) Copper (ppm) Table 2-18. Iron (ppm) Zinc (ppm) Selenium Nutrient Calcium

presented in Figure 2-13. The distribution of calculated mass intake of plant parts, at specific stages of maturity (e.g., ripe, unripe), is presented in Figure 2-14.

Discussion

Behavioral Observations

The distribution of daily activities for adult howlers in both groups MSE and LM1 is comparable to that previously reported for both red howlers and other members of the genus Alouatta. Gaulin and Gaulin (1982) indicated that red howlers (Alouatta seniculus) in Andean cloud forest spent 78.5% of their time resting, 12.7% feeding, and 5.6% moving. Mantled howlers (A. palliata) in Costa Rica spent 72.5% of the observation time resting, 18.1% feeding, and 7.8% moving (Glander and Teaford, 1995). Milton (1978a) reported that mantled howlers on Barro Colorado rested 66.2%, and fed 15.6% of sample days.

The distribution of time spent feeding on general and specific plant parts was markedly different between the seasons. For the 12-mo study period, group LM1 spent over 50% of their feeding time consuming leaf material. Results from the 5-mo observations of group MSE were very similar to those of the main study group (LM1). Similarities are most likely due to the inclusion of animals in the same age/sex classes. Perhaps even more important are the similarities

Figure 2-13. Major plant parts (leaf, fruit, flower) consumed by red howlers (group LM1), listed as percent of calculated mass intake.





Figure 2-14. Specific plant parts consumed by red howlers (group LM1), listed as percent of calculated mass intake.



in habitat and vegetation between the two home ranges.

Although seasonal fluctuations were seen in the foods selected during the observations, these variations in plant part consumption appear to be more a function of phenology than preferential selection. Similar variation was reported in red howler consumption of fruits by Braza et al. (1983).

Regardless of how the feeding time was analyzed (seasonally, or general versus specific plant parts), mature leaf material was consistently the largest component of the diet (27.8%-53.5%). This is supported by observations of the brown howler (A. fusca clamitans) that spent 80% of feeding time consuming leaf material, 10% flowers, and 10% fruits (Galetti, et al., 1987). These foods were selected from more than 30 different species of plants. In contrast, red howlers in French Guiana selected from 195 plant species as food sources. The major food categories included young leaves (54%), mature fruits (21.5%) and flowers (12.6%). The remainder consisted of immature fruits (4%), mature leaves (3%), termitaria (1.5%), and bark (0.4%), respectively.

Major food categories selected by mantled howlers on Barro Colorado Island, Panama included leaves (43.5%), fruit (46.5%), and flowers (10%). Although these findings seem to coincide with the results presented here for red howlers, the conclusions provided by Milton (1978a) describe howlers as "behavioral folivores". This suggests that the animals seek

"less fibrous" and more digestible plant materials, due to their apparent lack of gastrointestinal modifications for fermentive digestion of plant cell wall. Although the time spent feeding on young leaves, fruits and flowers may increase, as they become seasonally available, it is apparent from the nutrient analyses presented below, as well as the quantified digestive capabilities of howlers (see Chapter III), many species within the genus Alouatta are quite capable "anatomical folivores".

A similar selective feeding pattern was also observed in both red colobus (Colobus badius tephrosceles) and black-and-white colobus (Colobus guereza occidentalis) (Baranga, 1982), both of which have been previously defined as "anatomical folivores". When available, these colobine primates demonstrated preferences for young plant growth of two species, which were both lower in fiber components and higher in crude protein, than the mature plants of the same plants.

When feeding time devoted to the plant part of an individual species was examined, the prominence of seasonally available food items became more apparent. During the dry season, over 40% of the time spent feeding focused on the flowers of Pithecellobium saman and Guazuma ulmifolia and the fruits of Mangifera indica and Guazuma ulmifolia. More than 20% of the wet season feeding time was involved in the consumption of ripe and unripe fruits of Ficus pertusa.

When expressed as a percent of the calculated mass intake, the quantities of consumed fruits (45.4%) are greater than either leaves (34.3%) or flowers (20.2%). This "shift" in distribution of selected foods is mostly likely due the amount of processing time involved when the animals are consuming leaves versus fruits. Additionally, the quantity of food harvested per individual bite, as measure with the captive individuals, was greater when comparing fruits, across species and animal sex (1.86 g/bite) to leaves (0.40 g/bite). Therefore, per unit of time, a red howler could ingest more fruit than leaves, by weight. This is the major limitation of making assessments of distribution of food plants among specific parts or nutrient content of the diet, based on time feeding alone.

The reliance on leaves provides the howler with a food source which presents essentially the same relative composition throughout the year (Rosenberger, 1992). The inclusion of fluctuating food resources reported in several species of howlers, within a single species, and even within a single group over a 12-mo period, indicates the adaptability of the genus. These variable feeding habits, which conform to seasonal fluctuations of plant phenology, are suggestive of an "intermediate feeder", a description proposed by Hoffman (1973), for ungulate herbivores with similar dietary habits.

Plant Sample Collection and Analysis

Nutrient analyses of plant materials provided a new level of insight as to the composition of natural foods selected by a free-ranging primate.

Suggestions that wild fruits, flowers, or immature leaves are consistently lower in plant cell wall and higher in readily fermentable carbohydrates constitute a very broad and over-simplified generalization. Much of this misconception is based on an image of "fruits" as the commercially propagated items purchased in a grocery store for human consumption. These cultivated foods are substantially different in nutrient composition from those wild foods analyzed. The hazards of the above generalization become even more apparent when one examines the dietary ingredients offered to many captive folivores and observes the unfortunate results of using inappropriate foods (see Chapter III).

Intake Estimation

Calculated estimates, using bite count and bite weight data, of dry matter intake by an adult, free-ranging red howler (151.0 g/d) were similar to measured intakes determined with the temporarily captive adult red howlers (134.4 g/d), as well as long-term (6 mo+) dry matter intakes determined with captive adult red howlers (134.0 g/d) (see Chapter III).

Gaulin and Gaulin (1982) estimated the dry matter intake

of red howlers living in the Andean cloud forest. Their method relied upon the average weight of individual plant parts and the number of items consumed per feeding bout. Although their reports of time spent feeding (12.7%) are nearly identical to those reported here (12.6%), their estimates of dry matter intake were nearly double (266 g/d) these calculated estimates. Dry matter intake of mantled howlers was estimated by Nagy and Milton (1979). Their estimates were based upon an indirect method combining energy expenditure assessments with measurements of caloric content of food items, feces, and urine. These results combined with the assumption that the mantled howlers' fresh food consisted of two-thirds water, were the basis for estimates of dry matter intakes of 348 g/d.

Estimates of fresh food intake by mantled howlers, using similar methods combining focal-animal sampling with collections of representative food items, were made by Hladik and Hladik (1969). They calculated that a typical mantled howler consumed 1.145 and 1.220 kg/d of fresh food in the mid and late wet seasons, respectively. Nagy and Milton (1979) estimated that a mantled howler required approximately 1 kg of fresh food/d. Gaulin and Gaulin's (1983) estimated a red howler's fresh food intake was 1.23 kg/d. Each study reported fresh food intakes greater than those reported here (544.0 g/d).

Dry matter intake is a more reliable estimate when making comparisons between studies, as it removes the variable effects of water. Unfortunately, dry matter intake was not determined by Hladik and Hladik (1969). Differences in plants selected and the nutrient composition of those plants, environmental and climatic stresses, and animal body mass all play a role in the regulation of food intake. These variables would certainly explain the differences seen between the estimate made in this study and the results of the captive feeding trials. However, it is difficult to conceive of conditions which could produce a nearly two-fold increase in dry matter intake within individuals of the same species.

These consistently inflated estimates of food intake may partially explain why previous researchers have suggested that howlers have relatively poor digestive capabilities. Herbivores lacking gastrointestinal adaptations that promote microbial fermentation of plant material typically exhibit a feeding strategy involving consumption of large quantities of food, rapid rates of passage, and short periods of retention. These adaptations allow the animal to process quantities of material per unit of time. None of these traits, characteristic of herbivores like the red panda (Ailurus fulgens) and ruffed lemur (Varecia variegata), are seen in howlers (see Chapter III).

Nutrient Composition of Consumed Diet

Estimates of dry matter composition and nutrient intake, based only on time spent feeding, were consistently lower than estimated composition of the diet based upon bite count and weight data. Further discussion concerning nutrient intake is based upon the bite count/weight estimates.

Moisture content of the red howlers' diet (72.2%) was slightly lower than that predicted by Gaulin and Gaulin (78%) (1982). A higher moisture content might be predicted in vegetation found in the Andean cloud forest, which is subjected to less dramatic changes in total annual rainfall, as compared to the Venezuelan llanos. The 5.8% difference in moisture concentrations is not sufficient, on its own, to explain the difference in results observed between these two studies.

Estimated crude protein (CP) concentration (13.4%) of the consumed diet was lower than NRC (1978) recommendations for captive primate diets, including both Old (15% CP) and New World (<25% CP) Primates. The higher estimates of dietary protein need by New World Primates may apply more appropriately to the smaller sized tamarins and marmosets (Oftedal, 1991).

Previous estimates of protein concentrations in diets consumed by wild red howlers, one made by Gaulin and Gaulin (1982), used a method combining bite counts per plant part and

average plant part weight previously described. These authors predicted a protein content of 5%. Hladik (1978) estimated the average crude protein content in the annual diet as 9.6%.

The dietary nutrient concentrations which were the most remarkable included the components of the plant cell wall, cellulose, hemicellulose, and lignin. These fiber fractions, measured as neutral-detergent fiber (NDF), acid-detergent fiber (ADF), and acid-detergent lignin were greater than anticipated, especially in view of earlier suggestions that howlers were unable to utilize fibrous foods. The calculated concentrations (51.5% NDF, 37.8% ADF, 18.3% ADL) in the diet selected by the red howlers were comparable to concentrations found in orchardgrass hay (55.1% NDF, 30.7% ADF) or timothy hay (64.2% NDF, 37.5% ADF) (NRC, 1989).

The earlier estimates of low concentrations of these complex carbohydrates and polymers in the wild diet may have provided the basis for the previous inferences that howlers are limited folivores compared with the Old World colobines. Obviously, the presence of high concentrations of plant fiber in the diet selected by red howlers in the llanos, suggests otherwise. Evidence supporting the red howlers' adaptations to folivory also is seen in modifications of the gastrointestinal tract (Table 2-19). Elongation of the large intestine and cecum create regions of reduced digesta flow, allowing populations of symbiotic microorganism to flourish.

The mechanism by which these organisms degrade plant cell wall and the benefits to the host have been described in detail elsewhere in this text.

The presence of lignin, a nearly indigestible compound that functions to protect plant materials from consumption by herbivores, was higher than would be predicted based on

Table 2-19. Length of different portions of the gastrointestinal tract of red howlers (mm) (Braza et al., 1983).

Component	ರ್ರ್ (n = 27)	♀♀ (n =22)
Stomach (cardia-pylorus)	152.2 ± 29.3	145.5 ± 24.7
Stomach (pylorus-fundus)	188.6 ± 43.6	185.1 ± 37.6
Small intestine	2426.9 ± 166.7	2285.5 ± 205.8
Large intestine	1287.0 ± 247.4	1198.0 ± 181.1
Cecum	107.1 ± 7.8	101.8 ± 7.1

previous descriptions of howlers as "behavioral folivores". The relatively high ADL content (18.3%) of the selected diet was in excess of quantities recommended in captive animal diets, but still lower than those which promoted the formation of obstructive phytobezoars in folivorous primates fed highly lignified (23.6%) plant materials (Ensley et al., 1982). It should be noted that the formation of these phytobezoars has been reported primarily in colobine primates, although primates with a relatively less complex digestive system have also been affected (McClure et al., 1978).

Another variable which could explanation of the measured ADL concentrations include the effects of tannins. Tannins will often confound the crude lignin analysis (Oftedal, 1991). Sequential analysis of the detergent fiber fractions will help to resolve contamination of the ADL by tannins.

The estimated macro mineral (calcium, phosphorus, potassium, magnesium, sodium) concentrations in the selected diet were typical for many range plant materials. All macro minerals, except sodium (see below) were consumed in quantities which met or exceeded the estimated nutrient requirements for primates (NRC, 1978).

Phosphorus concentrations (0.18%) were in the lower range of the NRC (1978) guidelines (0.2%). Marginal to deficient phosphorus concentrations in vegetation under range conditions have been commonly noted (Preston, 1977). The first reports of phosphorus deficiency in ungulates involved pica in animals in South Africa. These herbivores subsequently died from botulism as a results of consuming bones from rotted carcasses (Van Soest, 1982).

The concerns over marginal phosphorus concentrations are compounded when one considers the ratio of phosphorus which is available to the animal. A less available form of phosphorus bound by phytate, a hexaphosphoric acid ester of inositol, is commonly found in plant materials. Phytate also binds calcium and zinc producing less available forms. No measurements were

made to determine the extent of phytate binding of minerals in this study. However, phytate phosphorus may be partially available to the red howlers. As seen in other hindgut fermenters (Hintz et al., 1973), red howlers may possess some phytase activity produced by the microbial populations of the cecum and large intestine. Approximately 50% of phytate phosphorus is absorbed in animals and man that lack extensive microbial fermentation (Parfitt et al., 1964; NRC, 1989).

Estimates of dietary sodium concentrations (97.3 ppm DMB) were surprisingly low, when compared to both recommended dietary allowances for humans (1000 ppm) (RDA, 1989) and nonhuman primates (2200-6400 ppm) (NRC, 1978). One would presume that the howler's dietary requirement for sodium under the climatic conditions of the Venezuelan llanos would be slightly greater due to the high temperatures. In horses, prolonged exercise and elevated environmental temperatures increase the need for sodium because significant amounts are lost in sweat (NRC, 1989). The sodium concentration in natural feedstuffs for horses is often less than 0.1 percent (1000 ppm) (NRC, 1989). Although howlers may not sweat appreciably, they must either be very efficient in conserving limited dietary sodium or they acquire sodium from sources which were not included among the wild foods collected.

Consumption of soils (Izawa, 1993), or soils processed by termites (termitaria), by red howlers has been reported in

both French Guiana (Julliot and Sabatier, 1993) and Colombia (Izawa and Lozano, 1990). Although not observed in these studies, numerous reports exist in the literature concerning the consumption of soils (geophagia) by primates, including red leaf monkeys (Davies and Baillie, 1988), colobus (Mackey, 1978; Harrison and Hladik, 1986), and gibbons, gorilla, red colobus, and indris (Hladik and Guegen, 1974).

Two hypotheses are typically presented explaining this behavior. One hypothesis is the assumption that geophagia is associated with high intakes of tannins in folivorous primate diets, so that the chemical and physical properties of ingested soils may assist in adsorption and excretion of these secondary plant metabolites. Another hypothesis presumes that soils are consumed as a supplemental dietary source of minerals.

There is no NRC estimated requirement for selenium concentrations in the diet of primates. This trace element, which functions as an essential component of the enzyme glutathione peroxidase (Rotruck et al., 1973), is required by swine at 0.1 and 0.3 ppm (NRC, 1988) in the diet. The selenium requirement of the horse was estimated at 0.1 ppm of diet (Stowe, 1967). The concentration of selenium in most feedstuffs, which is a function of soil selenium content and pH, ranges from 0.05 to 0.3 ppm (NRC, 1989). Regions of Venezuela have been previously recognized as seleniferous

zones (Jaffe, 1976).

The average concentration of selenium (1.04 ppm) in the diet selected by red howlers was higher than anticipated. Although some foods were high in this element, this average concentration was well within the limits of safety for domestic animals (NRC, 1989).

CONCLUSION

The red howler found in the central llanos of Venezuela were strictly herbivorous. The dramatic seasonal changes in amounts of rainfall, and the impact of those changes on vegetation abundance and diversity is apparent. The diet of the red howler fluctuated with these seasonal changes, yet mature leaves were consistently the primary food item selected during all times of the year. The remainder of the diet, composed of fruit, flowers, and leaves in various stages of maturity, was consumed opportunistically as seasonally available.

Chemical analyses of the selected plant parts, including those parts previously stated to be "more readily digestible" indicate that the red howler of this region consumes a highly fibrous diet. The structure and capacity of the hindgut permit the retention of digesta required for extensive microbial fermentation of plant cell wall. That howlers consistently spend over 75% of their day resting suggests a

conservation of energy expenditure. This may be a result of the lower digestible energy content of the selected foods, devotion of metabolic energy to utilization of fermentation by-products, and/or detoxification of secondary plant metabolites.

The selected diet also provides moderate amounts of protein and minerals. Although the red howler would appear to be a perfect candidate to benefit from coprophagy, this behavior was not observed. The variability in mineral concentrations seen in range plants consumed by free-ranging ungulates was also observed in the plants selected by the red howlers.

The red howler, although not a strict folivore, is a highly folivorous and adaptable feeder. The adaptations of this species to a fibrous diet, as with most herbivores, are seen at all levels, behavioral, anatomical, and physiological. Each of these adaptations complements the others, permitting the species to maximize the food potential of the vegetation with which it has coevolved.









LIBRARY Michigan State University PLACE IN RETURN BOX to remove this checkout from your record. TO AVOID FINES return on or before date due.

DATE DUE	DATE DUE	DATE DUE

MSU is An Affirmative Action/Equal Opportunity Institution

COMPARATIVE ADAPTATIONS TO FOLIVORY IN PRIMATES

Volume II

Ву

Mark Stephen Edwards

A DISSERTATION

Submitted to
Michigan State University
in partial fulfillment of the requirements
for the degree of

DOCTOR OF PHILOSOPHY

Department of Animal Science

1995

Chapter III

Digestive Capabilities and Digesta Transit in Leaf-eating Primates

Feeding Habits of Folivorous Primates

It is difficult to generalize the feeding habits of primates whose diet consists, in part or entirely, of leaf material. Indeed, within the order Primates, a wide range of folivory is exhibited (Oftedal, 1991). Much of this variation is a reflection of the differences in gastrointestinal anatomy of these species (Oftedal, 1991; Chivers, 1994).

The interaction of plants and animals has been studied extensively. Green plants, which are autotrophic and can manufacture essential organic substances for life, supply the nutriment for herbivorous animals and other organisms that feed on those plants (Van Soest, 1982). Plants also have developed mechanisms of defense which protect the plant, or parts of the plant, from consumption by herbivores. One of these defenses is plant fiber.

Dietary fiber influences a number of factors including passage rates of food through the gastrointestinal tract and

digestibility of nutrients (Cummings, 1978). There are no established minimum requirements for plant fiber concentrations in the diet of domestic animals or humans (RDA, 1989). In fact, dietary fiber has often been considered a "negative feed factor" that diminishes the value of a feed for support of production. However, the presence of fiber in the diet of herbivores plays a very important role.

The reduction or removal of fiber from the diet can compromise an animal's health (Cummings, 1978). Alternatively, inclusion of dietary fiber at appropriate concentrations may reduce the occurrence of intestinal disorders (Morin et al., 1978; Ensley et al., 1982; Ullrey, 1986; Helstab, 1988; Janssen, 1994).

The relative contributions of plant materials and the plant parts selected by primates vary widely. Numerous field studies have examined the types of foods selected by free-ranging primates, and the manner in which these studies were conducted has been found to influence the results (See Chapter II).

Based on field observations, hanuman langurs (Semnopithecus entellus) were found to spend only 25.7% of the daytime feeding (Newton, 1992). Feeding time was divided among the following food items: mature leaves (34.9%), fruits (24.4%), leaf buds (10.6%), flowers and flower buds (9.5%), young leaves (3.6%), insects (3.0%), and plant exudates/gums

(1%). The ability of hanuman langurs to utilize plant materials, the consumption of which comprises 96% of the time spent feeding, is a reflection of the specialization of the animal's gastrointestinal tract.

Black and white colobus (Colobus guereza) have been studied extensively in their natural habitat. Guatier-Hion (1983) concluded, based upon analysis of the stomach contents of five animals, that their diet included 50.5±34% leaf material, 49±34% fruit and seeds combined, and 0.5% other materials. Leaves of all phenological stages were consumed. Based upon visual observations of feeding behavior, Clutton-Brock (1975) found that black and white colobus fed mostly on shoots, flowers, and fruit during a 3-mo period. Leaf selection appeared to be a function of seasonal availability, with highest levels of consumption occurring during the dry season.

Analyses were conducted of the nutrient composition of leaf parts from two tree species, Celtis durandii and Markhamia platycalyx, that were preferred as food by both red colobus and black and white colobus (Branga, 1982). It was found that there were marked differences in the nutrient content of young and mature leaves, with young leaves being preferred over mature leaves by both species. The young leaf blades of Markhamia platcalyx had higher crude protein (33.1% vs. 22.0%), lower ADF (47.9% vs. 55.6%), and lower ADL (15.8%

vs. 23.7%) concentrations than mature leaf blades. A similar pattern was seen in the young leaf blades of *Celtis durandii*, with higher crude protein (39.0% vs. 22.1%), lower ADF (27.5% vs. 37.5%) and lower ADL (9.7% vs. 11.5%) concentrations than in mature leaf blades. Branga (1982) concluded that the most likely factors affecting food preference of free-ranging colobus were nutrient composition, inhibitory compounds (i.e., tannins, lignins), and the seasonal availability of alternate food items.

The proboscis monkey (Nasalis larvatus) has been the subject of several studies focusing on natural diet (Kern, 1964; Kawabe and Mano, 1972; Salter et al., 1985; Yeager, 1989). Yeager (1989) indicated that proboscis monkeys utilize at least 55 different plant species as food, with a marked preference for Eugenia spp. Although diverse species were selected, 11 plants species contributed 75.1% of the diet based on time spent feeding (Yeager, 1989). Similar to findings with other folivorous primates, seasonal shifts in foods selected by proboscis monkeys, were largely a function of availability.

Very little information exists on the feeding habits of or the plant species selected by the douc langur (*Pygathrix nemaeus*) (Lippold, 1977). Recently, a list of plants that were observed to be selected by this primate has been published (Nhat et al., 1993). However, no indication was

given of the proportion that individual plants or plant species contributed to the total diet.

It is clear that the pregastric fermentation seen in leaf-eating primates of the subfamily Colobinae and the food choices of these species are similar to those of selective ruminants (Kuhn, 1964; Bauchop and Martucci, 1968; Bauchop, 1978; Chivers and Hladik, 1980; Chivers, 1994). The importance of plant fiber for the health and maintenance of normal gastrointestinal function in leaf-eating primates should not be overlooked.

Captive Feeding Studies

The numbers of controlled feeding studies that have actually examined the digestibility of feeds and/or the rates of digesta passage in leaf-eating primates are few. This may be due to the limited number of animals available for digestibility trials in captivity, the lack of physical facilities required for proper collection of excreta, and concerns that the dietary manipulation required for such studies might be unduly stressful. As a result, there are few studies in which herbivorous primates have been individually housed and fed a single food at maintenance.

The studies found in the literature were either restricted by small sample size or by an apparent lack of control over multiple variables. It should be noted that

these constraints are not unusual in nutrition studies conducted with captive wild animals. The limited number of specimens held in captivity and the complex social structure of primate groups makes removal of animals from established groups for short-term isolation during feeding trials very difficult. Also, the physical features of zoo exhibits often are not appropriate for the collection of digestibility data.

One of the first studies examining the digestive capabilities of leaf-eating primates involved the silvered langur (Trachypithecus cristatus) (Bauchop and Martucci, 1968). The focus of this study was measurement of the production of volatile fatty acids as a result of microbial fermentation of alfalfa. The authors reported that the animals maintained a pH (5.0- 6.7) in the foregut that would allow anaerobic bacteria to ferment ingested plant material. Although cellulolytic bacteria, such as a gram-positive coccus and a gram-negative Bacteriodes spp., were present in the forestomach of the langurs, no protozoa (characteristically present in ruminants) were found. Celluloytic microorganisms produce end-products of fermentation that include ammonia, carbon dioxide, methane, and volatile short-chain fatty acids. These volatile fatty acids (VFAs), including acetic, butyric, and propionic acids, are primary source of energy for ruminant animals.

Fresh alfalfa was fed to the silvered langur (0 h), and

stomach contents were withdrawn via stomach tube at regular intervals (1, 1.5, 2, 2.5, 3.5, 4.5, 5.5, and 6.5 h). The VFA concentration peaked within 2 h after feeding, and held relatively constant over 6.5 h. Ammonia concentrations were also measured and were maintained at a high level, which the authors suggested was indicative of active bacterial proteolysis.

Another assessment of VFA concentrations in the gastrointestinal tract of captive leaf-eating monkeys involved the colobus monkey (Colobus polykomos) (Kay et al., 1976). Although balance trials were not conducted in this study, the progressive disappearance of food constituents along the entire length of the digestive tract was determined in two animals. The diet included only one item. This was a locally available exotic succulent, Commelinum benghalense. The two study subjects were sacrificed to determine the nutrient content of digesta at four points in the digestive tract.

Oxnard (1969) evaluated serum levels of vitamin B_{12} in two species of langurs (*Trachypithecus obscurus*, *Semnopithecus entellus*). The concentrations (1000 to 3000 pg/ml) were high when compared with anthropoid primates (150 to 500 pg/ml) (Oxnard, 1966). He proposed that vitamin B_{12} is synthesized by microbial organisms in the foregut of these langurs as seen in ruminant herbivores.

Milton et al. (1980) conducted two feeding trials with

wild-caught, mantled howler monkeys (Alouatta palliata) examining the digestibility of natural feedstuffs. Diet 1 consisted of 81% of four species of ripe fruits and 19% of Cecropia insignis leaves. Diet 2 consisted primarily (87%) of young leaves of different species and phenological stages and 13% of mixed species of fruits. The three animals used in the study, one adult male, one adult female, and one juvenile were housed individually; however, the variable nature of the diet was a primary limitation of the study.

Oftedal et al. (1982) examined dry matter intake and digestibility of a mixed-ingredient diet of constant composition that was fed to 2 black and white colobus monkeys (Colobus guereza). These animals were housed together, although both animals were similar in body weight (8.9 kg male, 8.8 kg female).

The acceptance of commercial primate biscuits, browse plants, and a produce mixture by a number of langur species including Francois' langurs (Trachypithecus francoisi), Douc langurs (Pygathrix nemaeus), hanuman langurs (Semnopithecus entellus), purple-faced langurs (Trachypithecus vetulus) and silvered langurs (Trachypithecus cristatus) was examined by Ullrey et al. (1982). The animals in this study were housed in groups of two to four individuals. The foods were offered in a manner that would permit differential selection by individual langurs, although the foods were weighed prior to

feeding. This study represents one of the most ambitious projects to date, yet the results were limited by the restrictions previously described.

Watkins et al. (1985) also examined the digestive capabilities of the black and white colobus (Colobus guereza). This study was one of the first to examine the acceptance and suitability of higher fiber primate biscuits, formulated to meet the specific needs of folivorous primates. Although this study also included a diet of mixed ingredients (Table 3-1), attempts were made to limit daily variability in intake by feeding fixed quantities of each food item at 95% of the animal's predetermined intake. Additionally, the two study subjects were individually housed, permitting collection of feces and orts without questions of their origin.

Table 3-1. Diet fed to two black and white colobus during pretrial and collection periods to determine digestibility (Watkins et al., 1985).

Feed group	Feed item
Commercial product	High fiber biscuit
Root vegetables	Sweet potato
Fruit	Apple
	Grapes
	Pear
	Orange
	Tomato
Green produce	Lettuce

Fiber digestion and digesta retention also were measured in silvered leaf monkeys (Semnopithecus cristatus) in a comparative study with Japanese (Macaca fuscata) and Rhesus macaques (Macaca mulatta) (Sakaguchi et al., 1991). Five adult langurs, each individually housed, were included in these trials. A single food item, a higher fiber "cubed" diet was offered ad libitum to the langurs in this study. The nutrient composition of the higher fiber "cubed" diet is provided in Table 3-2.

Measurements of food intake, digestion, and passage rate were made in captive proboscis monkeys at the New York Zoological Society (Dierenfeld et al., 1992). Although this is the only study in the literature in which attempts were made to quantify these measures in this species, this study has limitations that must be considered when interpreting the result.

Table 3-2. Concentrations of selected nutrients in a higher fiber "cubed" diet fed to silvered leaf monkeys (Sakaguchi et al., 1991).

Nutrient	Concentration (%)
Dry matter	91.9
Organic matter	81.4
Crude protein	18.3
Neutral detergent fiber	37.5
Acid detergent fiber	15.1

Unfortunately, all measurements were taken on a group of animals, one male and five females, that were not individuallyhoused specimens. Therefore, variability between individuals or sexes cannot be isolated nor can individual intakes be quantified. Additionally, the animals were fed a mixed diet, the contents of which are outlined in Table 3-3. These compounding variables preclude the authors from drawing conclusions concerning food intake or digestibility.

A summary of digestibility trials conducted with leafeating primates is presented in Table 3-4. A summary of studies examining rates of digesta passage and retention is given in Table 3-5. Due to the diversity of foods used and the variable consumption of those foods, the variation in collection procedures and analytical techniques used, the variations in animal body weight and stages of growth, and the feeding of mixed diets, no attempt has been made to compare these individual experiments.

Digestive Markers

A variety of markers has been used to determine the rate of passage and/or retention of consumed food items by animals. Additionally, markers may be included in a diet to serve as an indirect indicator of digestibility. Markers can be classified by their function and characteristics (Table 3-6).

Table 3-3. Diet fed to a group of six (one male, five females) proboscis monkeys at the New York Zoological Park (Dierenfeld et al., 1992).

Food group	Food item
Browse plants	Alder
	Grape
	Mulberry
	Willow
	Mangrove
Green produce	Snap beans
	Broccoli
	Celery
	Kale
	Spinach
	Bok choy
	Sunflower sprouts
Fruit	Banana
	Plantain
Commercial products	Canned primate diet (Zu*Preem TM)
	Dry biscuits (unspecified type)
Root vegetables	Carrots
	Yams
Miscellaneous	Eggs, hard-cooked
	Rice
	Peanut butter sandwich

<pre>Table 3-4. Apparent digestibility energy in various diets fed to lea</pre>	ibility to lea	44		for ates	dietary dr and humans		dietary dry matter, fiber, and gross and humans in previous studies.
Species	Diet	ΜO	NDF	ADF	ADL	GE	Author
	typeª						
Mantled howler	7	ı	0.23	ı	_	1	Milton et al., 1980
Mantled howler	8	1	0.41	-	_	-	Milton et al., 1980
Purple-faced langur	2	0.87	1	0.78	_	98.0	Ullrey et al., 1982
Purple-faced langur	4	0.84	1	0.72	_	0.81	Ullrey et al., 1982
Purple-faced langur	5	0.88	1	0.81	_	98.0	Ullrey et al., 1982
Purple-faced langur	9	0.88	1	0.74	_	18.0	Ullrey et al., 1982
Douc langur	2	0.81	1	08.0	1	0.78	Ullrey et al., 1982
Douc langur	4	0.81	1	0.71	1	0.78	Ullrey et al., 1982
Douc langur	5	0.83	1	0.78	_	0.82	Ullrey et al., 1982
Douc langur	9	0.85	•	0.73	_	0.82	Ullrey et al., 1982
Silvered langur	2	0.83	1	0.82	_	68.0	Ullrey et al., 1982
Silvered langur	4	0.85	_	0.70	ę	0.81	Ullrey et al., 1982
Silvered langur	5	0.89	_	0.79	1	0.84	Ullrey et al., 1982
Silvered langur	9	06.0	-	0.73	-	18.0	Ullrey et al., 1982
Silvered langur	1	0.76	0.69	0.62	ı	1	Sakaguchi et al., 1991

energy in various diets fed to leaf-eating primates and humans in previous studies. 3-4 (cont'd). Apparent digestibility coefficients for dietary dry matter, fiber, and gross (Table

Species	Diet	DM	NDF	DM NDF ADF ADL	ADL	GE	Author
	rype						
Black and white colobus	2	0.83	0.68	0.83 0.68 0.68 0.31	0.31	1	Oftedal et al., 1982
Black and white colobus	1	0.87	0.81	0.87 0.81 0.69 0.21	0.21	T	Watkins et al., 1985
Proboscis monkey (summer)	3	0.88	0.86	0.88 0.86 0.85 0.82	0.82	1	- Dierenfeld et al., 1992
Proboscis monkey (winter)	3	0.89	0.86	0.89 0.86 0.86 0.82	0.82	1	- Dierenfeld et al., 1992
Humans	IC	1	- 0.70	1	1	1	Slavin et al., 1981
Humans	HC	_	- 0.23	1	1	-	Slavin et al., 1981

1=nutritionally complete, higher fiber biscuit or cube.

including a standard primate biscuit, a canned primate diet, assorted green produce, root and fruit. 2=mixed diet, 3-mixed diet,

including a standard primate biscuit and canned primate diet (11%), assorted browse and including a standard primate biscuit, Eugenia browse, assorted green produce, root produce (55%), root vegetables (9%), fruit (11%), and miscellaneous items and fruit. 4=mixed diet, vegetables, green

including a standard primate biscuit, Hibiscus browse, assorted green produce, root 5=mixed diet,

including a standard primate biscuit, romaine lettuce, assorted green produce, root and fruit. and fruit. vegetables, 6=mixed diet, vegetables,

8=young leaves (87%), including Ficus indipida, Cecropia insignis, and immature fruits (13%), including (81%), including Ficus yoponensis, F. insipida, and young Cecropia insignis leaves (19%). Cecropia insignis, C. obtusifolia.

LC=low cellulose, including bread, fruits, vegetables, and refined grains. HC=high cellulose, including bread with added Solka Floc (BW=200, Brown Co., Berlin, NH), fruits, vegetables, and refined grains.

and mean retention time (R_{GIT}) of various recovery, markers fed to leaf-eating primates. (TT,), Transit time Table 3-5.

Species	type ^a	type	(h)	(h)	Rgit (h)	Author
Spider monkey	3	plastic	4.4	1	1	- Milton, 1981
Mantled howler	2,3	2,3 plastic	20.4	1	-	- Milton, 1981
Red howler	1	seed	21	54	134	134 Crissey et al., 1989
Proboscis monkey	4	bead	14	1	49	49 Dierenfeld et al., 19
Silvered leaf monkey	1	Cr	13.6	1	46.8	46.8 Sakaguichi et al., 19
Silvered leaf monkey	1	CO	13.6	1	44.8	44.8 Sakaguichi et al., 19
"1=nutritionally complete, higher fiber biscuit or cube	gher fiber	: biscuit or	cube.			

992

3-mixed diet, including bread, cottage cheese, green produce, and fruit.
4-mixed diet, including a standard primate biscuit and canned primate bit (11%), assorted browse and green produce (35%), root 'vegetables (3%), fruit (11%), and miscellancous items (16%). 2=ripe fruits (81%), including Ficus yoponensis, F. insipida, and young Cecropia insignis leaves (19%).

Table 3-6. Classification of markers used to evaluate rates of digesta passage (Kotb and Luckey, 1972).

Elements

Inert metals (heavy & rare earth metals)

Natural isotopes (40K)

Artificial isotopes (144Ce)

Compounds

Inorganic Metal oxides (Cr_2O_3)

Mineral salts (BaSO₄)

Organic Natural dyes (carmine, chromogen)

Synthetic dyes (methylene blue)

Other (lignin, cellulose, plant sterols)

Particulates

Polymers Polyethylene glycol, plastics, glass

Cells Yeast, bacteria

Charcoal

Metal particles (e.g., sized aluminum particles)

Other particulates Seeds, cotton string

The use of these markers in nutrition and physiological studies has been outlined by Kotb and Luckey (1972), and Udén (1978). Soluble markers must meet the following criteria, as modified by Udén (1978):

- 1. Be strictly nonabsorbable.
- Not be subject to degradation or precipitation by the stomach or intestine.
- 3. Not be trapped by mucus or other gastrointestinal

contents during its flow through the intestine.

- 4. Not influence gastrointestinal motility.
- 5. Be easy to measure and give accurate results.
- 6. Have the same solubility characteristics as the test substance with which it might be used.
- 7. Not influence digestion.
- 8. Not interfere with the measurement of other substances.
- 9. Be measurable in opaque or strained samples.
- 10. Be estimated in very small concentrations thereby causing only a minimal osmotic disturbance.
- 11. Have the same "diffusion space" as the test substance.
- 12. Have the same physical characteristics as the fraction of interest during its residence in the gastrointestinal tract.
- 13. If used as a stain or mordant, should be firmly attached, and unaffected by digestion, pH, or other factors.
- 14. Be reasonable in price.

These specific requirements are more critical for markers used for rate of passage experiments than for markers used in balance trials (Udén et al., 1980).

MATERIALS AND METHODS

General

Four separate experiments were conducted at two zoological institutions, the Cleveland Metroparks Zoo (CMZ), Cleveland, OH, and the San Diego Zoo (SDZ), San Diego, CA. All trials were conducted between August 1992 and August 1993. The experimental protocol was coordinated within the established routines of daily animal care and exhibit or enclosure maintenance.

In vivo apparent digestibilities of dry matter, fiber fractions, and energy, rate of passage, and time of retention were measured for two diets of differing fiber concentrations. The experiments are listed in Table 3-7.

Table 3-7. Experimental feeding trials and data collected.

Experiment	Data collected
In vivo apparent digestibility	Total food intake, total fecal collection
Acetate bead marker	Rate of marker passage and retention
Co-EDTA marker	Rate of marker passage and retention
Cr-mordanted fiber marker	Rate of marker passage and retention

The staffs of both facilities were extremely cooperative

in providing individual housing of study subjects for the duration of the study. Each animal's response to the experimental procedure was monitored very closely. As a result, three individuals who failed to adapt to one of the two test diets were removed from that portion of the trial. A fourth animal was never integrated into the study because of inadequate consumption of the test diet. The welfare of these animals, all of which were endangered species, was considered paramount.

In each experiment, a cross-over design was used so that each individual might consume each of the test diets. Although the intention was to allow all individuals to remain in the study until completion, some individuals were removed from the study prior to the final collection, for reasons unrelated to the experimental treatments.

Test Diet Formulation

Two test diets formulated to offer a medium and high level of fiber were designated 15ADF and 30ADF, respectively, in reference to their targeted acid-detergent fiber (ADF) concentrations. Both diets were formulated to be nutritionally complete, meeting the estimated nutrient requirements of both New and Old World Primates for maintenance (NRC, 1978). The diets were manufactured as

extruded morsels approximately 27 x 15 x 5 mm (Marion Zoological, Inc., Plymouth, MN 55441). No additional food items were offered.

The two diets were made of the same ingredients, using variations in proportions of soy hulls and corn grain to produce the desired fiber concentrations. The composition of the two test diets is given in Table 3-8. The fiber concentration of the two test diets was the principal variable. The estimated concentrations of selected nutrients in the diets are presented in Table 3-9.

A sample of each test diet was ground in a Wiley mill fitted with a 2-mm screen. The samples were analyzed for dry matter (DM), ash, gross energy (GE), neutral-detergent fiber (NDF), ADF, cellulose (C), hemicellulose (HC), acid-detergent lignin (ADL), Ca, P, Mg, Na, K, Fe, Mn, Zn, and Se.

Animals, Housing, Feeding, and Collection of Excreta During Digestion Trials and Handling of Samples

Ten separate species/subspecies were used. Their scientific and common names are listed in Table 3-10. Based on records of previous physical examinations and current medical status, there were no apparent health problems with these animals.

Table 3-8. Composition (percentage by weight, air-dry basis) of the lower and higher fiber test diets.

Ingredient	Die	et
	15ADF	30ADF
Soybean meal (48% crude protein)	13.00	5.00
Corn gluten meal	11.00	15.00
Corn grain	24.00	
Sugarbeet pulp	5.00	5.00
Soybean hulls	19.762	52.762
Corn hominy feed	10.00	5.00
Sucrose	10.00	10.00
Soybean oil	3.00	3.00
Dicalcium phosphate	1.85	2.20
Calcium carbonate	0.80	0.45
L-Lysine (78%)	0.30	0.30
Sodium chloride	0.50	0.50
Vitamin-trace mineral premix ^a	0.50	0.50
Mold inhibitor ^b	0.10	0.10
Red cabbage extract (color)	0.10	0.10
Vitamin C (ethocel coated)	0.05	0.05
Feed flavor	0.038	0.038

^{*}Formulated to provide per kg of diet: 8000 IU vitamin A, 1750 IU vitamin D₃, 250 IU vitamin E, 5 mg menadione, 3 mg thiamin, 4 mg riboflavin, 22.5 mg niacin, 1 mg pyridoxine, 0.1 mg biotin, 15 mg D-Ca-pantothenate, 0.3 mg folic acid, 25 mcg vitamin B₁₂, 200 mg choline, 25 mg Fe, 10 mg Cu, 90 mg Zn, 45 mg Mn, 1 mg I, 0.23 mg Se.

b MonoProp™ (50% propionic acid on verxite), Anitox Corp., Buford, GA.

Table 3-9. Estimated nutrient concentrations (dry matter basis) of the test diets.

Nutrient	D:	iet
	15ADF	30ADF
Dry matter (%)	91.2	91.7
Gross energy (kcal/g)	4.46	4.39
Crude protein (%)	20.5	20.0
Lysine (%)	1.1	1.0
Ether extract (%)	5.9	5.2
Ash (%)	6.1	6.7
Neutral-detergent fiber (%)	24.0	42.0
Acid-detergent fiber (%)	15.0	29.6
Acid-detergent lignin (%)	1.2	1.5
Crude fiber (%)	10.8	22.5
Calcium (%)	1.0	1.1
Phosphorus (%)	0.7	0.7

All animals were housed individually for the duration of the study. Every attempt was made to house conspecifics in adjacent enclosures to permit continued visual, olfactory, and vocal communication. However, due to the nature of the study, isolation for measurement of individual feed consumption and fecal output was necessary.

CMZ animals were housed indoors. SDZ animals were housed outdoors, with the exception of *Pygathrix nemaeus nemaeus*, Colobus guereza kikuyuensis, Cercocebus galeritus chrysogaster, and Mandrillus leucophaeus leucophaeus which had

free access to indoor/outdoor enclosures, except during cleaning and collection procedures. The enclosures varied in size and offered a variety of climbing structures. Concrete flooring was present in all enclosures.

Table 3-10. Scientific and common names of species included in the study.

Family/ subfamily	Scientific name	Common name
Lemuridae	Varecia variegata rubra	Red ruffed lemur
	Varecia variegata variegata	Black & white ruffed lemur
Alouattinae	Alouatta caraya	Black howler
	Alouatta palliata	Mantled howler
	Alouatta seniculus sara	Red howler
Colobinae	Colobus guereza kikuyensis	Black & white colobus
	Trachypithecus francoisi francoisi	Francois' langur
	Pygathrix nemaeus nemaeus	Douc langur
Cercopithecinae	Cercocebus galeritus chrysogaster	Golden-bellied mangabey
	Mandrillus leucophaeus leucophaeus	Drill

The animals had free access to water. Feed intake was monitored daily. Over a 4-wk period, animals were gradually converted from the mixed diets (which included commercially

available foods, produce, and browse) fed prior to the study to the test diets. The test diets were incrementally added to the daily diet, as the previously fed items were removed.

During preliminary trials and the conversion period, the quantities of test diets offered were increased until each animal was leaving 30 to 50 g daily to ensure ad libitum intake. Weighed quantities of the test diet were offered daily at approximately 0900 h. After approximately 24 h, remaining quantities of food (orts) were collected and weighed. The test diets were fed exclusively for a minimum of 1 wk (adaptation period) prior to the 1st d of fecal collection.

Following the adaptation period, feces were collected for 14 d. All feces were collected daily during the morning cleaning procedures using a plastic putty knife. Daily fecal samples were kept separate. Collected feces were sealed in plastic bags to prevent moisture loss and stored at 0°C prior to analysis. Urine was not collected and no attempt was made to prevent urinary contamination of the fecal material. Due to the arboreal habits of these primates, urinary contamination was not a primary concern.

Chemical Analysis of Feed and Feces

The weight of the frozen fecal material was recorded and

all samples were freeze-dried in preparation for analysis. Following freeze-drying, samples were reweighed, crushed into a fine powder, and thoroughly mixed. Test diets and feces were analyzed for DM, GE, NDF, ADF, and ADL. moisture/dry matter concentration was determined using the freeze-dried DM and a final DM. The final DM was determined on approximately a 1.0-g freeze-dried subsample in an aluminum weighing dish by overnight drying in a vacuum oven at 60°C. Gross energy was determined by complete combustion of an approximately 1-g sample in a Parr adiabatic oxygen bomb calorimeter (Parr Instrument Co., Moline, IL). Following combustion, the remaining residue was titrated with 0.0725 N sodium carbonate to correct for the heat of formation of nitric and sulfuric acids. Sequential fiber analysis as described by Goering and Van Soest (1970) was used to determine NDF, ADF, and ADL. Seguential analysis produces lower AD-nitrogen values (Udén, 1978). Sulfite was not used in the ND treatment because it solubilized part of the lignin. When sulfite was not used, somewhat higher NDF-nitrogen values resulted. Approximately a 0.5-g subsample was used for the procedure. A 2% α-amylase (A-3051, Sigma Chemical Co., St. Louis, MO) solution was used in the NDF analysis (Robertson and Van Soest, 1977). The NDF residue was treated with acid detergent (AD) to determine ADF. ADL was determined on the

ADF residue using 72% sulfuric acid (Goering and Van Soest, 1970). ADL residue was ignited at 500°C in a muffle furnace to determine the AD-insoluble ash residue. The fiber residues were calculated as follows (Udén, 1978):

ND residue - AD residue = hemicellulose

AD residue - AD lignin = cellulose

AD lignin - AD-insoluble ash residue = lignin

All analyses were performed in duplicate.

Apparent dry matter digestibility (ADMD) and digestibility of dietary components, fiber, and energy were estimated according to the simple relationship:

When possible, the animals were weighed at the beginning of the study, during the crossover transition period, and at the end of the study. Individual animals were moved into plastic airline kennels and weighed on an electronic balance. Recently recorded weights obtained from medical records were used for those individuals who would be unacceptabily stressed by the weighing procedure.

RATE OF DIGESTA PASSAGE MEASUREMENTS

Rate of passage experiments using acetate beads were conducted in conjunction with digestibility trials. Co-EDTA and Cr-mordanted fiber trials were conducted separately.

Marker Administration and Sampling Frequency

Inducing domestic animals to consume markers at the desired time can lead to a variety of problems. Encouraging primates to consume these markers produced an entirely new set of obstacles. All markers were mixed with ground test diet (\$ 20 g) and fruit nectar (typically mango, \$ 25 g) to enhance palatability. The fruit nectar also provided adequate moisture to hold the marker/test diet mixture together. The dough-like mixture was rolled into balls, placed into individually labeled bags, and hand-fed to each subject.

CO-EDTA (14.35% Co) was dosed orally at 0.50 g per individual, providing 71.75 mg Co. Cr-mordanted fiber was fed in quantities to deliver 40 mg Cr to the test subject.

These methods were successful, but a few individuals continued to refuse the marker after several attempts. Thus, the marker was placed in a small portion (\leq 10 g) of either white grape or banana. The marker was consumed by

most of the remaining animals using these vehicles.

Every attempt was made to feed the markers to each individual within the hour from 0900 to 1000 h. Only when an individual refused the initial attempt was a follow-up attempt made the following day, during the same time period (0900-1000 h).

All feces were collected. Following the introduction of the markers, feces were collected at 3-h intervals. The collections from the study subjects were made in the same order that the markers were fed. A post-dusk sample was collected at 2100 h and a pre-dawn sample was collected at 0300 h. Fecal samples were not collected or observed at 0000 h, so as not to disrupt the animals during the night. This collection schedule was continued for 2 wk following introduction of the marker.

Samples were immediately transferred to plastic storage bags upon collection and stored at 0°C until analyzed.

Marker Synthesis and Mordanting of Plant Fiber Co-EDTA

Udén (1978) found that a Li Co-EDTA salt produce diarrhea in some sheep and goats, but the Na Co-EDTA did not yield similiar conditions. For that reason, a Na Co-EDTA salt was selected as a liquid marker for all liquid phase digesta

transit experiments.

The Na Co-EDTA was synthesized using an adaptation of the procedure described by Udén (1978) as follows:

- Weigh 25 g of Co(II) acetate•4H₂O (cobaltous acetate),
 29.2 g EDTA (ethylenediaminetetraacetic acid), 4.0 g sodium hydroxide, and 200 ml of distilled water into a 2-L beaker.
- Record initial volume. Dissolve, stirring continuously with moderate heat (if necessary) for 4 to 5 h.
- 3. Discontinue heat and restore to the initial volume with ddH_2O (double distilled water). Stir overnight.
- 3. Place beaker in ice bath on stir plate to permit continued stirring. Add SLOWLY 20 ml of hydrogen peroxide (30% solution). ** CAUTION: WILL FOAM OVER IF TOO MUCH HYDROGEN PEROXIDE IS ADDED AT ONCE. **
- 4. Remove from ice bath. Continue stirring for 5 to 6 h.
- 5. Divide equally into two 2-L beakers.
- 6. Add 150 ml ethanol (95% v/v) to each beaker. Leave at room temperature for 1 to 2 h. Refrigerate overnight (or longer). NOTE: Longer period of refrigeration, such as a weekend, will increase final

yield.

- 7. Filter on a large Buechner funnel under vacuum and wash with ethanol (80% v/v).
- 8. Equilibrate with moisture in air for 24 h.
- 9. Place in a storage container. Take a sample (5 g) for Co assay.

Cr-mordanted fiber

Chromium was used as the particulate marker in all solid-phase digesta transit trials. Chromium has been described as one of the easiest particulate markers with which to work, inexpensive, easy to analyze, easy to mordant, and accepted as a safe marker for human studies (Udén, 1978). Cr-mordanted fiber has been shown to be indigestible (Martz et al., 1974).

The Cr was mordanted to a macroamount of NDF from each of the test diets. The procedure for mordanting Cr to the plant fiber was as follows (modification of Uden et al., 1980):

- Prepare a macroamount of plant cell wall by boiling in neutral detergent (ND) solution for 1 h, approximately 50 g of diet per 1 L of ND solution.
- 2. Rinse thoroughly with hot ddH_2O through a brass sieve followed by a rinse with acetone.
- 3. Place in a 65°C oven until dry.

- 4. Weigh cell wall fraction.
- 5. Add a water solution providing 375 g sodium dichromate $(Na_2Cr_2O_7)$ for every 1000 g of plant fiber (an amount of chromium equivalent to 10% of the amount of fiber used) in amounts sufficient to cover the fiber.
- 6. Boil cell wall fraction at 100°C for 3 h.
- 7. Wash the fiber thoroughly with tap water.
- 8. Suspend the fiber in tap water and add ascorbic acid equivalent to 50% of the weight of plant cell wall (the liquid should taste slightly sour). Let soak for at least 1 h.
- 9. If the fiber has turned green, rinse until free of soluble material and dry at 65°C in a forced-draft oven.

Analysis of Markers

Frozen samples were weighed, freeze-dried, and re-weighed. The freeze-dried samples were crushed to a fine powder and thoroughly mixed. Ten samples were randomly selected from the 14-d collection period to determine DM using approximately a 1.0-g subsample in an aluminum weighing dish by overnight drying in a vacuum oven at 60°C. Freeze-dried feces were digested with nitric-perchloric acid (Fenton and

Fenton, 1979). Concentrated nitric acid (20.0 ml) and 70% perchloric acid (5.0 ml) were added to 1 g of ground sample in a Phillip's beaker. The sample was digested until 1 ml of acid was left in the beaker. Following digestion, the samples were diluted with ddH_2O . All samples were digested in duplicate, unless a small sample size demanded otherwise.

All samples were analyzed using an atomic absorption/emission spectrophotometer (Smith-Hieftje 4000, Thermo-Jarrell Ash). Chromium samples were analyzed in a high-temperature nitrous oxide/acetylene flame with the emission wavelength set at 425.4 nm. Cobalt samples were analyzed in an air/acetylene flame with the absorption wavelength at 240.7 nm. All analyses were performed in duplicate.

Mathematical Calculations of Retention Time, Transit Time, and Rate Constants.

The mean retention time, including transit time (R_{GIT} + TT), was calculated according to Blaxter et al. (1956).

$$R_{GIT} + TT = \frac{\Sigma (Y_i \cdot t_i)}{\Sigma Y_i}$$

Transit time (TT) was determined as the time between dosing and first appearance of the marker in the feces. R_{GIT} was determined by subtracting TT from R_{GIT} + TT.

Statistical analyses were completed using QuattroPro (Novell Applications Group, Orem, UT). Comparisons among

gastrointestinal tract type within diet were made using a oneway analysis of variance. Comparisons between diets within the gastrointestinal tract group were made by paired t-test.

RESULTS

Body weights of all individuals before, during, and following the trials are shown in Appendix Table B1. The results from the digestibility trials for each individual are shown in Appendix Tables B2-B12. The results from the digesta marker trials for each individual are shown in Appendix Tables B13-B19. The digestibility and marker results for the single golden-bellied mangabey and drill are not included in the statistical analyses but can be found in Appendix Tables B17, B18, and B19, respectively.

Body Weight Changes

The body weights of *Varecia variegata* were significantly lower when maintained on diet 30ADF than when consuming diet 15ADF (P<0.01). There were no statistically significant differences (P>0.05) among the body weights of the two other groups, *Alouatta* spp. and colobine primates, fed diets 15ADF or 30ADF (Table 3-11).

The mean body weights of species and/or subspecies included in statistical analyses are shown in Table 3-12.

2.86

28.6

Table 3-11. Mean body weight (BW) and mean dry matter intake (DMI) of primate study subjects fed diets 15ADF and 30ADF grouped by gastrointestinal tract (GII) type.	and mean by gast	n dry roint	ht (BW) and mean dry matter intake (DMI) grouped by gastrointestinal tract (GIT)	e (DMI) t (GIT)	of primate study type.	subjects
GIT Type/Species	Diet	п	BW (kg)	DMI (g/d)	DMI (g/kg BW)	DMI (% BW)
SIMPLE GIT						
Varecia variegata spp.	15ADF	5	4.68ª	136.3°	° 29.0°	2.90ª
	30ADF 5	2	4.04 ^b	222.6 ^d	d 54.9 ^b	5.49b
HINDGUT FERMENTER						
Alouatta spp.	15ADF	7	7.52	132.4	18.9	1.89
	30ADF 7	7	7.61	161.9	3 22.4	2.24
FOREGUT FERMENTER						
Colobus, Pygathrix,	15ADF	7	7.54	213.7	7 29.7	2.97

30ADF 7 7.46 204.0 column with different superscripts differ (P<0.01). c,d Values in same column with different superscripts differ a,b Values in same Trachypithecus

Table 3-12. Mean body weight (BW) fed diets 15ADF and 30ADF grouped	and mean by spec	dry mies/su	and mean dry matter intake (DMI) by species/subspecies.		of primate study subjects	subjects
Species	Diet	п	BW (kg)	DMI (g/g)	DMI (g/kg BW)	DMI (% BW)
Varecia variegata rubra	15ADF	m	4.58	139.0	30.0	3.00
	30ADF	٣	4.00	183.8	45.6	4.56
Varecia variegata variegata	15ADF	2	4.83	132.4	27.5	2.75
	30ADF	7	4.10	280.8	8.8	6.88
Alouatta caraya	15ADF	8	8.11	128.0	17.6	1.76
	30ADF	m	8.11	185.1	25.5	2.55
Alouatta villosa palliata	15ADF	2	5.85	141.0	23.9	2.39
	30ADF	7	6.45	131.5	20.3	2.03
Alouatta seniculus sara	15ADF	7	8.32	130.6	16.1	1.61
	30ADF	2	8.03	157.7	19.7	1.97
Colobus guereza kikuyuensis	15ADF	1	10.90	154.8	14.2	1.42
	30ADF	7	10.70	153.8	14.4	1.44
Pygathrix nemaeus nemaeus	15ADF		12.10	428.9	35.5	3.55
	30ADF	н	11.75	391.1	33.3	3.33
Trachypithecus f. francoisi	15ADF	5	5.96	182.4	31.6	3.16
	30ADF	5	5.94	176.6	31.8	3.18

Dry Matter Intake

Dry matter intakes were expressed as grams of dry matter consumed per d (24 h), and as a function of the animal's body weight, g/kg BW or %BW. Statistically significant differences between diets 15ADF and 30ADF were observed only in Varecia variegata spp. Dry matter intakes of diet 30ADF by Varecia variegata spp. were statistically higher than of diet 15ADF when expressed as g/d (P<0.05) and as a function of body weight (P<0.01). However, there were no statistical differences (P>0.05) between the dry matter intakes of either diet by Alouatta spp. or the colobine species (Table 3-11).

The mean dry matter intakes of species and/or subspecies included in the statistical analyses are shown in Table 3-12.

Apparent Dry Matter Digestibility

The apparent dry matter digestibility (ADMD) of diet 30ADF by both Varecia variegata and colobine species was statistically (P<0.05) lower than that of diet 15ADF. The ADMD of diet 30ADF by Alouatta spp. was also significantly (P<0.01) lower than that of diet 15ADF (Table 3-13). Statistically significant differences were observed in ADMD between gastrointestinal tract type (simple, hindgut, foregut), within either diet 15ADF or 30ADF (P<0.0005).

The mean ADMD of diets 15ADF and 30ADF by species and/or

(DMD), digestible energy (DE) of diets for primate study subjects grouped by and digestible energy intake (DEI) Mean apparent dry matter digestibility tract 15ADF and 30ADF, gastrointestinal Table 3-13.

GIT Type/Species	Diet	n	DMD	DE	ΞQ	DEI
			(8)	(%)	(kcal/g)	(kcal/kg ^{0.75})
SIMPLE GIT						
Varecia variegata ssp.	15ADF	5	51.0ª,e	47.0	2.20	92.80
	30ADF	₂	41.7 ^{b.e}	39.7	1.74	134.49
HINDGUT FERMENTER						
Alouatta spp.	15ADF 7	7	69, 3°, f	68.5°	3.05	92.74
	30ADF	7	61.7 ^{d, f}	61.5 ^d	2.70	98.16
FOREGUT FERMENTER						
Colobus, Pygathrix,	15ADF 7	7	81.200	°6.08	3.61	173.79
$\it Trachypithecus$	30ADF 7	7	76.6 ^{d,9}	75.6 ^d	3.32	152.31
a,b Values in same column with different superscripts differ (P<0.05).	ifferent s	Silber	scripts differ	P<0.05		

e,f,g Values in same column with different superscripts differ (P<0.0001). " Values in same column with different superscripts differ (P<0.01).

subspecies shown in the statistical analyses are included in Table 3-14.

Energy Digestibility and Energy Intake

The digestible energy (DE) of diet 30ADF was significantly (P<0.01) lower than that of diet 15ADF for both the colobine species and Alouatta spp. No statistically significant differences were demonstrated between diets 15ADF and 30ADF fed to Varecia variegata (Table 3-13).

The mean DE (% or kcal/g) of diets 15ADF and 30ADF, and DE intake (kcal/kg $^{0.75}$) by species and/or subspecies included in the statistical analyses are shown in Table 3-14.

Fiber Component Digestibility

No statistical differences were seen in the digestibility of the individual cell wall components (NDF, ADF, C, HC, ADL) between diets 15ADF and 30ADF within each gut type (P<0.01), except for the digestibility of HC by Alouatta spp., which was significantly (P<0.01) lower in diet 15ADF than in diet 30ADF (Table 3-15).

The mean percent digestibility of various fiber components (NDF, ADF, C, HC) in diets 15ADF and 30ADF by species and/or subspecies included in the statistical analyses are shown in Table 3-16.

(DMD), digestible energy (DE) of diets for primate study subjects grouped by 94.35 DEI (kcal/kg^{.75}) 108.44 90.45 89.01 195.95 156.25 83.72 111.33 114.12 87.80 84.89 88.78 90.28 173.57 240.92 177.07 2.25 1.79 2.96 2.70 3.19 3.18 2.68 3.50 3.42 3.64 3.62 3.32 3.07 1.71 2.71 (kcal/g) 48.8 44.4 40.8 66.4 61.6 68.89 61.8 71.5 78.5 78.0 81.6 72.5 81.2 75.7 39.1 61.1 DE (%) 42.5 62.0 70.0 73.9 53.3 41.2 71.0 61.8 80.3 78.7 76.2 76.7 47.7 67.7 82.4 62.1 rable 3-14. Mean apparent dry matter digestibility 15ADF and 30ADF, and digestible energy intake (DEI) DWD DWD (%) C \sim \sim \sim $^{\circ}$ \sim 2 \sim 2 2 S Ŋ 15ADF 30ADF 15ADF 30ADF 30ADF 30ADF 15ADF 30ADF 15ADF 15ADF 30ADF 15ADF 15ADF 30ADF 15ADF 30ADF Diet Varecia variegata variegata Colobus guereza kikuyuensis Trachypithecus f. francoisi Alouatta villosa palliata Pygathrix nemaeus nemaeus variegata rubra sara seniculus species/subspecies Alouatta caraya Alouatta Varecia Species 15ADF

Table 3-15. Mean digestibility (%) of various fiber components (NDF, ADF, C, HC) in diets	sof va	rious s gro	fiber compc uped by gas	nents (NDF, trointestina	ADF, C, HC)	in diets
15ADF and 30ADF 101 Prime	Diet	n	NDF	ADF	Ŋ	HC
GIT Type/ special						
SIMPLE GII	15ADF	2	20.4	9.4	11.5	38.8
Varecia varios	30ADF	5	20.7	12.6	8.7	38.9
FERMENTER						
HINDOOT TITLE	15ADF	7	44.8	42.2	39.2	50.3ª
Alouatta orr	30ADF	7	47.4	40.6	39.1	62.4 ^b
FOREGUT FERMENTER						
Colobus, Pygathrix,	15ADF	7	77.1	7.67	80.4	71.9
S. (100)	30ADF 7	7	74.7	71.3	75.2	77.0
ne column	erent s	npersc	with different superscripts differ ($P < 0.01$)	r (P<0.01).		

15ADF and 30ADF for primate study	subjects '		grouped by spe	species/subspecies.	ies.	
Species	Diet	n	NDF	ADF	ວ	HC
Varecia variegata rubra	15ADF	٣	22.9	12.1	15.9	40.3
	30ADF	ю	19.0	10.6	9.9	38.0
Varecia variegata variegata	15ADF	7	16.6	5.3	5.9	36.5
	30ADF	7	23.3	15.8	11.9	40.3
Alouatta caraya	15ADF	Ж	46.5	40.5	38.0	53.6
	30ADF	ю	45.8	37.7	36.8	63.6
Alouatta villosa palliata	15ADF	7	43.7	43.8	39.5	51.8
	30ADF	7	52.5	46.2	44.5	8.99
Alouatta seniculus sara	15ADF	7	43.4	43.1	40.8	43.8
	30ADF	7	44.8	39.5	37.3	56.4
Colobus guereza kikuyuensis	15ADF	1	77.0	80.1	81.4	6.79
	30ADF	Ч	74.3	56.2	76.1	78.0
Pygathrix nemaeus nemaeus	15ADF	-	66.5	9.99	59.7	66.4
	30ADF	Н	8.69	9.79	66.1	74.8
Trachypithecus f. francoisi	15ADF	Ŋ	79.3	82.3	84.3	73.7
	30ADF	5	75.7	75.0	76.9	77.3

Markers

There was no statistical difference (P>0.01) seen in either TT_1 (Table 3-17) or R_{GIT} (Table 3-19) between diets 15ADF and 30ADF for acetate bead markers, Co-EDTA, and Cr-mordanted fiber fed to Varecia variegata, Alouatta spp. or the colobine species. Unfortunately, due to previously mentioned circumstances, eight animals had to be removed from the second Cr-fiber trial, thus eliminating them from the statistical analyses. The results of the first Cr-fiber trial for these animals are provided in the Appendix (B15-B16).

The mean TT_1 and R_{GIT} of species and/or subspecies fed diets 15ADF and 30ADF and included in the statistical analyses are presented in Tables 3-18 and 3-20, respectively.

Markers (acetate bead, Co-EDTA, and Cr-mordanted fiber) were compared to each other in order to determine the level of correlation for parameters measured (TT_1 , R_{git}). TT_1 and R_{git} as estimated by different markers within diet 15ADF or 30ADF were not significantly correlated.

Discussion

General

Fermentation of the plant cell wall is a relatively slow process, requiring the retention of digesta in specific segments of the gastrointestinal tract (Parra, 1978). The rate at which digesta transit the tract is influenced by

Table 3-17. Mean transit time (TT_1 , h) of various markers, acetate bead, Co-EDTA, and Cr-mordanted fiber, in diets 15ADF and 30ADF for primate study subjects grouped by gastrointestinal tract (GIT) type.

GIT Type/Species	Diet	Acetate bead	CO-EDTA	Cr-fiber
SIMPLE GIT				
Varecia variegata spp.	15ADF	3.0 (6)	2.7 (5)	2.1 (5)
	30ADF	2.5 (6)	2.7 (5)	2.7 (5)
HINDGUT FERMENTER				
Alouatta spp.	15ADF	32.3 (4)	15.6 (7)	23.3 (4)
	30ADF	24.8 (4)	16.1 (7)	15.8 (4)
FOREGUT FERMENTER				
Colobus, Pygathrix,	15ADF	31.5 (2)	16.5 (6)	46.5 (1)
${\it Trachypithecus}$	30ADF	37.5 (2)	21.0 (6)	43.5 (1)
an in parentheses				

Table 3-18. Mean transit time (TT_1 , h) of various markers, acetate beads, Co-EDTA, and Cr-mordanted fiber, in diets 15ADF and 30ADF for primate study subjects grouped by species/subspecies.

Species	Diet	Acetate bead	CO-EDTA	Cr-fiber
Varecia variegata rubra	15ADF	2.5 (3)ª	3.5 (3)	2.5 (3)
	30ADF	3.5 (3)	2.5 (3)	3.5 (3)
Varecia variegata variegata	15ADF	3.0 (2)	1.5 (2)	1.5 (2)
	30ADF	1.5 (2)	3.0 (2)	1.5 (2)
Alouatta caraya	15ADF	!	16.5 (3)	I
	30ADF	I	9.5 (3)	ı
Alouatta villosa palliata	15ADF	18.0 (2)	7.5 (2)	12.0 (2)
	30ADF	13.1 (2)	7.5 (2)	4.5 (2)
Alouatta seniculus sara	15ADF	46.5 (2)	22.5 (2)	34.5 (2)
	30ADF	36.0 (2)	34.5 (2)	27.0 (2)
Colobus guereza kikuyuensis	15ADF	31.5 (1)	43.5 (1)	46.5 (1)
	30ADF	31.5 (1)	43.5 (1)	43.5 (1)
Pygathrix nemaeus nemaeus	15ADF	31.5 (1)	1.5 (1)	1.5 (1)
	30ADF	43.5 (1)	19.5 (1)	I
Trachypithecus f. francoisi	15ADF	1	13.5 (4)	I
	30ADF	-	15.8 (4)	1
an in narentheses				

'n in parentheses.

Table 3-19. Mean retention time $(R_{git}, \ h)$ of various markers, acetate bead, Co-EDTA, and Crmordanted fiber, in diets 15ADF and 30ADF for primate study subjects grouped by gastrointestinal tract (GIT) type.

GIT Type/Species	Diet	Acetate bead	CO-EDIA	Cr-fiber
SIMPLE GIT				
Varecia variegata spp.	15ADF	5.2 (6) a	4.1 (5)	3.8 (5)
	30ADF	4.9 (6)	2.4 (5)	2.3 (5)
HINDGUT FERMENTER				
Alouatta spp.	15ADF	30.1 (4)	34.1 (7)	38.8 (4)
	30ADF	24.2 (4)	40.4 (7)	38.8 (4)
FOREGUT FERMENTER				
Colobus, Pygathrix,	15ADF	30.7 (1)	41.1 (6)	37.6 (1)
${\it Trachypithecus}$	30ADF	37.6 (1)	38.4 (6)	68.6 (1)
2 2 20 20 20 20 20 20 20 20 20 20 20 20				

an in parentheses.

Species	Diet	Acetate bead	CO-EDIA	Cr-fiber
Varecia variegata rubra	15ADF	3.8 (3)	4.0 (3)	3.1 (3)
	30ADF	6.3 (3)	3.3 (3)	5.0 (3)
Varecia variegata variegata	15ADF	2.2 (2)	4.3 (2)	1.6 (2)
	30ADF	2.3 (2)	1.1 (2)	3.4 (2)
Alouatta caraya	15ADF	1	18.4 (3)	
	30ADF	1	38.4 (3)	1
Alouatta villosa palliata	15ADF	25.7 (2)	33.6 (2)	37.8 (2)
	30ADF	25.4 (2)	30.6 (2)	36.8 (2)
Alouatta seniculus sara	15ADF	34.5 (2)	58.2 (2)	39.9 (2)
	30ADF	22.9 (2)	53.3 (2)	40.8 (2)
Colobus guereza kikuyuensis	15ADF	20.7 (1)	44.0 (1)	37.6 (1)
	30ADF	37.6 (1)	53.4 (1)	68.6 (1)
Pygathrix nemaeus nemaeus	15ADF	1	35.5 (1)	33.8 (1)
	30ADF	1	32.5 (1)	•
Trachypithecus f. francoisi	15ADF	1	41.1 (4)	•
	30405	1	36 1 ///	

factors such as diet composition, rate of digestion, and level of intake (Allen and Mertens, 1988; Church, 1988). The effect of intake on dietary digestibility is much more pronounced when an animal is fed a mixed-ingredient diet than one composed of a single food, as in this study. In general, as intake increases, dry matter digestibility is depressed as a consequence of an increased rate of digesta passage through the gastrointestinal tract. The depression of digestibility related to increased intake is not limited to fiber but also includes more readily fermentable components of the diet such as starch.

Digesta transit time in herbivores, when compared across dietary and gastrointestinal types, tends to decrease from ruminants to nonruminants with hindgut fermentation to more generalist feeders (e.g., omnivores). The rate of passage in herbivores generally increases with decreasing body size (Parra, 1978). It is generally believed that nonruminant herbivores exhibit a faster rate of passage and a shorter mean retention time as compared to foregut fermenters (ruminants) (Moir, 1968; Parra, 1978).

It is apparent that these factors are closely interrelated and dependent upon one another. The level of intake
influences transit time, and thus influences digestibility.
In turn, the digestibility of the diet influences the level
of intake. These are the complex relationships that make

comparisons between this study and other studies so challenging. Although several reviews have been published (Hintz, 1969; Olsson, 1969; Parra, 1978) drawing comparisons between the digestive capabilities of foregut and hindgut fermenters, they were limited by the previously described variables. However, the reviewers suggested that the fiber digesting capabilities of nonruminant herbivores are commonly less than those of ruminants.

Generally speaking, animals with a larger body mass consume a smaller percentage of their body weight on a daily basis because their mass-specific energy requirements (kcal/kg) decrease with increasing body size (Van Soest, 1982). The level of consumption also is a function of the caloric density of the diet.

Varecia variegata spp.

Surprisingly, very little information could be located in the literature concerning the natural feeding habits or gastrointestinal anatomy of either black and white or red ruffed lemurs (Tattersall, 1982). Additionally, no previous measurements or estimates of dry matter intake by Varecia variegata spp. were available. Petter et al. (1977) described Varecia as frugivorous, a feeding strategy that Tattersall (1982) suggested is in agreement with inferences from their dental morphology.

Recent studies of Varecia sp. in their native habitat indicate that these animals utilize fruits and nectar as 70.8% and 15.3% of their total food resources, respectively (White, 1991). Leaves (11.1%), undesignated as to maturity, and flowers (2.8%) make up the remaining portions of the diet. The author noted that the flowers were licked but not consumed, suggesting their use as a source of pollen and/or nectar. Although the majority (86.1%) of the diet appeared to consist of what would be considered "more digestible" food items, there was no chemical analysis to support this assumption.

Food items selected by a captive group of black and white ruffed lemurs were described in a study of species behavior by Pereira et al. (1988). As with the previously mentioned study, no attempt was made to quantify food items selected. The author did indicate that the animals consumed leaves, grasses, buds, flowers, and acorns (Quercus spp.). However, there was no suggestion that this diet was comparable to that selected by animals in their native habitat.

The digestive capabilities of *Varecia variegata* ssp. have been a subject of curiosity among many who have cared for this species in captivity. When fed a mixed diet similar to those described in previous studies, food items such as yams, carrots, and apples appear in the feces in a physical form similar to that which was offered (e.g., cubed, sliced). From

these observations, an assumption has been made that these animals utilize very little, if any of these foods and raises questions concerning the ability of this species to digest other foods as well. This assumption seems to be supported by the relatively low dry matter digestibilities seen in this study (51.0% of 15ADF and 41.7% of 30ADF). The reduced dry matter digestibility of 30ADF is a further representation that this species lacks specific gastrointestinal adaptations for utilization of a fibrous diet.

The significant reduction in body weight of Varecia variegata ssp. when consuming diet 30ADF also was indicative of the limited ability of this species to use a diet with elevated levels of plant cell wall. Although dry matter intakes increased significantly, there was a concurrent reduction in digestibility of dry matter. Since food was always provided ad libitum, it appears that the 30ADF diet alone was not suitable to support the caloric requirements of Varecia variegata spp.

Ruffed lemurs have demonstrated a coprophagic behavior in captivity (Edwards, personal observation), although the animals were not permitted access to their feces during these trials. This type of nutrient recycling may permit the ruffed lemur to recover some of the undigested nutrients lost in the fecal matter.

Another common observation concerning Varecia ssp. in

captivity is the apparent rapid rates of digesta transit. As one keeper noted, "you can return 20 min after feeding the animals, and find the food you just offered in the feces". In fact, some have questioned whether the material was fecal in origin or regurgitated food, due to the similar physical characteristics and rapid appearance. Although the first fecal collection was conducted at 3 h post ingestion of the dietary marker, visual observations made during the collection period suggest that transit time (TT₁) is more rapid (<1 h) than the 3.0 h reported here.

Although there were no statistically significant differences between the estimates of DE for 15ADF and 30ADF for Varecia, the animals fed 30ADF consumed more dry matter (5.49% BW) than those fed 15ADF (2.90% BW). This presumably was a metabolically driven attempt to maintain caloric intake. However, this strategy appears not to have been effective since there was a significant reduction in mean body mass (4.68 kg vs. 4.04 kg) while consuming 30ADF. These animals were simply unable to digest and absorb adequate calories to maintain their body mass.

The above observations all suggest that lemurs lack gastrointestinal adaptations that permit the retention of digesta for a time sufficient to allow microbial fermentation of plant cell wall. Even so, this species selects a natural diet entirely of plant origin. Similar digestive patterns

and dietary habits have been observed in another herbivore of comparable body size, the red panda (Ailurus fulgens) (Table 3-21).

Table 3-21. Digestible energy (DE) and dry matter intake (DMI) of two nonspecialized herbivores of similar body weight.

Species	Diet	DE (kcal/g DM)	DMI (g/kg BW)
Varecia variegata	15ADF	2.20	29.0
Varecia variegata	30ADF	2.74	54.9
Ailurus fulgens	G/alf ^b	3.14	28.6
Ailurus fulgens	G/bam ^c	3.69	27.0

aWarnell (1988).

The high rate of digesta passage suggests that limited digestion of insoluble plant material requires an increased turnover of ingesta (Table 3-22). This is reflected in the

Table 3-22. Transit time (TT_1, h) and total mean retention time (R_{git}, h) of particulate markers for two nonspecialized herbivores of similar body weight.

Species	Diet	\mathtt{TT}_1	$R_{\tt git}$
Varecia variegata	15ADF	3.0	5.2
Varecia variegata	30ADF	2.5	4.9
Ailurus fulgens	Bamboo	5.0	5.7
Ailurus fulgens	G/alf ^b	5.2	9.6
Ailurus fulgens	G/bam ^c	6.1	13.3

aWarnell (1988).

bGruel mixture with 10% alfalfa meal.

^cGruel mixture with 10% bamboo (*Psuedosasa japonica*) meal.

bGruel mixture with 10% alfalfa meal.

^{&#}x27;Gruel mixture with 10% bamboo (Psuedosasa japonica) meal.

slower rates of transit and longer retention in the red pandas fed bamboo versus a forage/concentrate mixed diet. With lower gross energy digestibilities (DE), Varecia ssp. have adapted by attempting to fulfill their energy needs by exposing a large mass of food to digestive processes per unit of time and by rapidly passing the undigested portion.

Rapid rates of transit and reduced capacities for fiber digestion indicate that a substantial portion of the diet selected by free-ranging ruffed lemurs would be passed in the fecal material. This feeding/digestive strategy would be an ideal method of seed dispersal for many of the fruits consumed by this species, presuming the seed coat can withstand the limited digestive assault demonstrated in this study.

Strict adherence to descriptions of the five classes or degrees of herbivory in primates (Eisenberg, 1978) (see Chapter I) would place Varecia variegata spp. in Class 1, based upon their "apparent lack of digestive tract modification". Yet the dietary habits of primates in this class (omnivorous feeding strategies) do not correspond with current information on the natural dietary habits of ruffed lemurs. Therefore, if Varecia spp. are classified according to a specific degree of herbivory, this species appears to fit the description of Class 2 more appropriately. Future descriptions of the gastrointestinal anatomy of Varecia spp. will help to clarify this issue.

Alouatta spp.

Members of the genus Alouatta have been described as "behavioral folivores" suggesting that howlers, lacking the extensive gastrointestinal adaptations of the Old World folivores, select foods perceived as more readily digestible (e.g., young leaves and fruits). Although this description may apply to a particular species as it behaves in a specific region (i.e., Alouatta palliata on Barro Colorado Island; Milton, 1978), the description may be too broad to encompass the entire genus.

The digestibilities of NDF by the three species of Alouatta examined in this study were higher than those reported by Milton et al. (1980) (Table 3-23). The differences between studies may be explained by the dramatic differences in diet types and fiber sources tested, as well as by differences in methodology. One would predict that a diet composed of foliage and fruits, both containing unprocessed or "long-stem" fiber would be lower in digestibility than the extruded test diets, which consist of "processed" fiber particles.

The presence of a voluminous, complex large intestine is presumed to have evolved before the adaptations for foregut fermentation exhibited by the Old World colobines. The results of this study demonstrate that howler monkeys possess the capability of digesting and utilizing fibrous feeds, but

Table 3-23. Apparent digestibility (%) of plant cell wall (NDF), hemicellulose (HC), and cellulose (C) by howler monkeys

(Alouatta spp.).

Species	Diet	NDF	НC	С
A. caraya	15ADF	46.5	53.6	38.0
	30ADF	45.8	63.6	36.8
A. seniculus	15ADF	43.4	43.8	40.8
	30ADF	44.8	56.4	37.3
A. palliata	15ADF	43.7	51.8	39.5
	30ADF	44.8	56.4	37.3
A. palliata ^a	Fruitb	23	18	29
	Leaf	41	63	67

aMilton et al. (1980).

the ADMD of diets 15ADF and 30ADF represent only 85.3% and 80.5%, respectively, of the same values measured in colobines.

Comparable differences have been reported (Udén, 1978) for other hindgut and foregut fermenters fed similar diets. Horses digested 85.8% as much DM as goats, and ponies digested 81.0% as much DM as heifers.

The significant reduction (11%) in ADMD seen in howlers fed diet 30ADF versus 15ADF supports the relatively lower ability of a hindgut fermenter to utilize a higher fiber food. The colobine species exhibited only a 5.7% reduction in ADMD when comparing diets 30ADF and 15ADF.

bFruit diet consisted of 81% ripe fig fruit from Ficus yoponensis, and F. insipidia, and 19% flush leaves from Cecropia insignis.

^cLeaf diet consisted of 87% young leaves from Ficus insipida and, Cecropia insignis, and 13% immature fruits from Cecropia insignis and C. octusifolia.

Howler monkeys digested relatively more of the hemicellulose fraction of both diets 15ADF and 30ADF than the cellulose fraction as compared to digestion of those components by the foregut fermenters. This has been observed in hemicellulose digestion by domestic hindgut fermenters (equids) versus foregut fermenters (cattle, sheep, goats) as Although there is no single well (Van Soest, 1994). explanation for this, there has been speculation that xylan cannot be processed by microorganisms until the arabinosic side chains are removed. Another view is that the digestion of xylan is dependent upon the removal of cellulose that surrounds it. It is possible that some linkages which are sensitive to gastric digestion, must be removed from the xylan to allow further digestion, which in this case would occur in the hindgut. Finally, it is thought that some hemicellulosic carbohydrates may exist as glycoproteins that are released by the action of acid pepsin. Apparent digestibilities of cellulose and hemicellulose for three species of howlers are compared to values determined in other hindgut fermenters in Table 3-24.

In general, it is believed that nonruminant herbivores have shorter retention times than ruminants. These trends were not consistently exhibited in this study. Estimates of digesta transit reported here are comparable with two previous studies examining these parameters in *Alouatta* sp. (Milton,

Digestibility (%) of cellulose (C) and hemicellulose (HC) by various nonruminant Table 3-24. animals.

diringro.					
Species	Diet	υ	ЭН	Author	
Alouatta caraya	15ADF	38.0	53.6	Current study	ybu
	30ADF	36.8	63.6	Current study	ndy
Alouatta villosa palliata	15ADF	39.5	51.8	Current st	study
	30ADF	44.5	8.99	Current st	study
Alouatta seniculus sara	15ADF	40.8	43.8	Current st	study
	30ADF	37.3	56.4	Current st	study
Elephant, Asian	Alfalfa	99	49-68	Foose, 1	1982
	Timothy	45-57	52-54	Foose, 1	1982
Elephant, African	Alfalfa	44-62	69-75	Foose, 1	1982
	Timothy	42-46	38-49	Foose, 1	1982
Horse, pony	Timothy	37	38-45	Udén, 1	1978
	Timothy	37-42	38-45	Foose, 1	1982
Zebra, mountain	Alfalfa	54-64	36-48	Foose, 1	1982
	Timothy	38-43	47	Foose, 1	1982
Tapir, Asian	Alfalfa	33-35	10-27	Foose, 1	1982
	Timothy	29-47	36-57	Foose,]	1982

1981; Crissey et al., 1989). Although not statistically significant, TT_1 was more rapid for Co-EDTA and slower for Cr-mordanted fiber when the howlers were fed 15ADF than when fed 30ADF. With acetate beads, TT_1 with values determined on diet 30ADF were more rapid than those for 15ADF and $R_{\rm git}$ values were shorter. The only other $R_{\rm git}$ determined with howlers and reported in the literature (134 h) (Crissey et al., 1989) was significantly longer than those reported in this study (24.2-38.8 h). Although the diets used in these two studies were comparable in composition, the types of markers used were not similar and this may account, in part, for the differences between the studies.

It appears that the three species of Alouatta studied exhibited all of the criteria proposed by Eisenberg (1978) for classifying howlers as Class 4 herbivores, having both the gastrointestinal tract modifications and natural diets containing at least 40% leaves. This classification also is supported by the results presented in Chapter II. If one were to apply Hoffman's (1973) classification of herbivores, the howler might be considered an intermediate feeder, exhibiting the ability to shift from leaf material to plant parts that are relatively more digestible as a function of seasonal availability.

Colobines (Colobus, Pygathrix, Trachypithecus)

One of the important characteristics of herbivores is an increased capacity for temporary retention of food in the gastrointestinal tract, when compared to omnivorous or carnivorous species (Parra, 1978). The fermentation contents (digesta in the rumen or saccus + cecum + colon) expressed as a percentage of BW, of langurs and other herbivores with foregut fermentation are compared in Table 3-25.

Table 3-25. Fermentation contents (FC) (rumen or saccus + cecum + colon) of species with foregut fermentation, expressed as a percentage of BW (Parra, 1978).

Species	FC (%BW)	Author
Langur	13.0	Kuhn (1964)
Langur	17.4	Bauchop and Martucci (1968)
Sheep	10.1	Elsden et al. (1946)
Sheep	15.6	Boyne et al. (1956)
Cattle	10.0	Nevens (1928)
Cattle	17.8	Elsden et al. (1946)
Eland	11.9	Hungate et al. (1959)
Camel	12.3	Hungate et al. (1959)

Dry matter intakes of both diets 15ADF and 30ADF, 1.42% and 1.44% of body weight, respectively, seen in *Colobus guereza* in this study were lower than those reported by both Oftedal et al. (1982), 1.8%, and Watkins et al. (1985), 1.99%. Watkins et al. (1985) suggested that the elevated levels of

consumption of the test diet in their study were a function of the higher (25% versus 16%) cell wall (NDF) content of the test diet. However, one would predict that intake levels of 15ADF (24% NDF) in this study would be similar to those reported by Watkins et al. (1985). The absolute dry matter intakes, 154.8g/d of 15ADF and 153.8g/d of 30ADF, which were not statistically different between the two treatments, are similar to those reported in other studies with colobus (159 g/d; Oftedal et al., 1982; and 158 g/d; Watkins et al., 1985). Variations seen between the three studies may be a result of the differences in composition of the test diets, differences in their caloric densities, and differences in the body weights and body compositions of study subjects.

Dierenfeld et al. (1992) found that mean dry matter consumption of an individual proboscis monkey in a group of male and female animals fed a mixed ingredient diet was 3% of estimated body mass. The group of foregut fermenters involved in this study (Colobus, Trachypithecus, Pygathrix) consumed comparable levels of dry matter as a percentage of body weight (2.97% of 15ADF, 2.86% of 30ADF) with no significant differences between dietary treatments.

The single douc langur, which was the heaviest animal (\bar{x} = 11.9 kg) included in the study, had the highest level of dry matter intake as a function of body weight (3.55% 15ADF, 3.33%

30ADF). This species is also the closest in body size to the proboscis monkey. However, Chivers (1994) indicated that the gastrointestinal tracts of these two species may be different. The gastrointestinal tract of the douc langur was described as the heaviest and most voluminous, relative to the other colobines examined by Chivers (1994). The proboscis monkey had only average volume. Although the proboscis had a very high functional absorptive area (FAA), the potential fermenting volume was low (Table 3-26).

Plant cell wall is also important as a primary restrictive determinant of intake (Van Soest, 1982). This is due in part to its effect on gut fill. However, the intake of forages and other fibrous feeds can be increased through the reduction of particle size by grinding and/or pelleting.

Dry matter digestibilities of 88 and 89% were calculated by Dierenfeld et al. (1992) for proboscis monkeys fed a mixed-ingredient diet during summer and winter trials, respectively. Although these digestibility estimates appear unusually high, Ullrey et al. (1982) also reported dry matter digestibilities as high as 90% in silvered leaf monkeys fed a mixed diet without browse plant supplementation. This higher level of dry matter digestibility appears to be related to the presence of readily fermentable and highly digestible foods (e.g., Zu-Preem Primate biscuits, fruits, lettuce) in the diets offered during the collection periods.

	Gut len	ength	Gut weight	ight	Abso	Absorptive a	areab	Fermenting volume ^c	ng vc	olume ^c		CGD⁴	
Species	GL	GL/BL	M.S	8 BW	Ą	% of total	AA/BL	FV &	% of total	FV/BW	Area	Wgt	Vol
Colobus guereza	483	7.8	363	5.8	2175	89	41	3332	92	392	2.1	2.7	12
S. entellus	445	7.0			2650	89	41	4473	06	447	1.7		9.3
T. vetulus	334	6.2			1650	64	31	2977	95		5.6		17
T. cristatus	540	10.4	372	6.1	2770	71	53	4023	68	099	1.4	4.9	8.1
T. obscurus	472	6.3	278	4.8	2460	70	48	3192	87	550	1.5	3.9	7
Presbytis melalophos	547	11.2	256	3.9	2477	73	51	2715	98	418	1.2	2.5	9
Presbytis rubicunda	548	10.5	229	4.4	2343	75	45	2598	98	499	н	3.6	9
Pygathrix nemaeus	526	9.5	276	6.7	2613	71	46	3575	88	872	1.4	4	7.3
Nasalis larvatus	988	13.8	598	3.8	4776	74	75	7244	87	456	1.1	2.9	6.4

*GL=gut length, cm; BL=body length, cm; GW=gut weight, g; BW=body weight; AA=absorptive area, FV=fermenting volume, cm³

bArea of small intestine+1/2(stomach+cecum+colon), cm³.

cvolume of stomach+cecum+colon, cm³.

d(stomach+cecum+colon)/small intestine.

Sakaguchi et al. (1991) found the dry matter digestibility of a manufactured ("cubed") diet (16.4% ADF, DMB) fed to silvered leaf monkeys (Semnopithecus crisatus) was 75.7%. This value is lower, but comparable to the results of this project. François' langurs, a closely related species, digested 82.4% of the dry matter in diet 15ADF. The douc included in this study exhibited dry langur digestibilities of 76.2% and 73.9%, for diets 15ADF and 30ADF, respectively. These values were consistently lower than for other colobines (Colobus, Trachypithecus) in this study maintained on the same diets. Additionally, these values were lowest apparent dry matter digestibility estimates reported for a colobine primate. Dry matter digestibility can be affected by the level of intake during the collection period (Church, 1988).

The significantly lower ADMD of diet 30ADF by all foregut fermenters (Colobus, Pygathrix, Trachypithecus) studied suggests that the elevated fiber concentrations in this diet provided a digestive challenge to these species. ADMD of mixed diets, including those containing browse plants, were not dramatically lower than those seen in this study.

Fiber component digestibility exhibited by the colobines demonstrates that these species are suited to consume a diet containing higher concentrations of plant cell wall. The lack of any significant differences in fiber digestibility between

the two diets further supports these findings. There were slight numerical differences between fiber digestibilities of the two diets, with digestibilities of NDF, ADF, and C in diet 30ADF of 96.9%, 89.5%, and 93.5%, respectively, of those digestibilities in diet 15ADF. Interestingly, as seen in the hindgut fermenting howlers, hemicellulose digestibility was numerically higher in diet 30ADF (77.0%) than in diet 15ADF (71.9%).

Similar, but numerically lower digestibilities of fiber by proboscis monkeys fed a mixed ingredient diet were as follows: 86.2-86.4% NDF, 87.1-90.4% HC, 85.0-86.0% ADF, 87.8-89.1% C, and 81.7-82.3% ADL (Dierenfeld et al., 1992.). results support Chivers (1994) description of the proboscis monkey, the largest colobine included in his study, possessing smallest stomach and largest large intestine. He indicated that this anatomical arrangement results in a low fermenting ability and high absorbing capability. This is further supported by natural dietary information indicating that this species eats few mature leaves (<50%), but selects fruit and young leaf parts (Yeager, 1989). Additionally, seed consumption approaches 20% in free-ranging proboscis. extent of microbial fermentation which results contributions of the hindgut in the proboscis was not examined.

Herbivores with foregut fermentation are more efficient

in their ability to digest plant fiber due to their adaptations to: consume large quantities of plant material, retain it for longer periods of time, and provide the environment required to support microbial digestive processes (Stevens, 1982). Table 3-27 includes dry matter and NDF digestibility estimates for three species of colobines included in this study compared to values determined for other species with foregut fermentation. The digestible energy estimates reported by Ullrey et al. (1982) are comparable to those seen in this study, when the differences in test diet are considered. Digestible energy was not determined by any other authors.

Bauchop and Martucci (1968) estimated the daily energy produced from microbial fermentation of ingesta in the foregut of an Asian colobine using the following assumptions. stomach contents average 12% of body weight, and contents are, on average, 15% dry matter. The dry matter in the stomach contents of a 4.5-kg langur would be equivalent to 81 g. The quantified VFA molar composition o f acetic:propionic:butyric:valeric acids was 56:24:13:7, and the average measured rate of fermentation per h was 460 µmole/g of dry matter. They estimated that this would result in a daily energy equivalent of 283 kcal based upon the heat of combustion of each fatty acid. Using Klieber's (1961) interspecific value (BMR = $70BW_{kg}^{0.75}$), the basal energy

Digestibility (%) of dry matter and neutral-detergent fiber (NDF) by species 1982 1982 1973 1973 1982 Current study Current study Current study Current study Current study Current study Foose, 1982 1982 Foose, 1982 Foose, Foose, Foose, Hintz et al., Foose, al., Author et Hintz 66.5 79.3 75.7 6.09 54.9 49.8 64.5 39.8 69.4 54.2 .5 8.69 58, NDF 78.7 73.9 76.7 52.9 71.5 70.3 54.6 63.8 80.3 76.2 82.4 56.4 60.2 72.7 M Alfalfa Timothy Alfalfa Timothy Alfalfa Alfalfa Timothy Alfalfa 30ADF 15ADF 15ADF 30ADF 30ADF Diet 15ADF Colobus guereza kikuyuensis Trachypithecus f. francoisi Pygathrix nemaeus nemaeus fermentation Llama-guanaco foregut Table 3-27. Barasingha Species Cattle Sheep Eland with

requirement for a 4.5-kg langur would be 216 kcal per day. This latter value is 76% of Bauchop and Martucci's (1968) estimate of energy yield from microbial fermentation.

When the same assumptions used to calculate the above values are applied to the colobine primates involved in this study, 61 to 79% of the energy yield of microbial fermentation would be required for basal metabolic needs. The remainder presumably would be available toward support of the total maintenance requirement. These values are shown in Table 3-28.

The rate of digesta passage in small ruminants is expected to be more rapid than that of larger ruminants to support the animal's elevated energy demands, as a function of metabolic body weight $(BW^{0.75})$. One would predict a similar trend when comparing different species of colobines. Rates of passage (TT_1) determined with colobines in this study were slightly longer than those reported in two other studies (Sakaguchi et al., 1991; Dierenfeld et al., 1992). Based upon acetate bead marker recoveries of 96, 88, and 52% from proboscis monkeys fed a mixed ingredient diet, Dierenfeld et al. (1992) found a TT_1 of 14 h and a R_{GIT} of 49 h. This suggests that the proboscis monkey, when fed a mixed-ingredient diet has a TT_1 nearly twice as rapid as other colobines fed a higher fiber diet.

Results comparable to those determined with Francois'

Table 3-28. Body weight (BW), estimated daily energy production from microbial fermentation (MFE) of digesta based upon calculations by Bauchop and Martucci (1968), estimated basal metabolic rate (BMR) and percent of MFE used for BMR by colobines.

Species	ID	BW (kg)	MFE (kcal/d)	BMR (kcal/d)	BMR/MFE (8)
Trachypithecus cristatus	NA⁴	4.50	283	216	92
Colobus guereza kikuyuensis	Flower	11.34	713	433	61
	Gomez	10.90	685	420	61
Pygathrix nemaeus nemaeus	Rhett	12.10	761	454	09
	1001	17 7	787	324	67
rachypichecus i. mancoisi	400	•	7	F 30	ò
	Janet	5.67	357	257	72
	Chrissie	6.67	419	291	69
	Stanley	3,99	251	198	79
	Fungar	5.75	362	260	72
	GooNgoot	5.85	368	263	71
abata from Banchon and Martucci	.1 (1968)				

*Data from Bauchop and Martucci (1968).

langurs are reported with silvered leaf monkeys fed a higher fiber "cubed" diet using like markers (Table 3-29). The rates of passage (TT_1) determined with Cr-mordanted fiber were slower and retention times (R_{git}) shorter in this species when compared to the silvered leaf monkey.

Table 3-29. Rates of digesta passage (TT_1 , h) and mean retention time (R_{git} , h) determined in two langur species fed similar higher fiber (15% ADF) diets.

Species	Marker type	TT ₁	R_{git}
Silvered leaf monkey	Cr	13.6	46.8
Silvered leaf monkeya	Со	13.6	44.8
Francois' langur	Cr	27.0	30.5
Francois' langur	Со	13.5	41.1

^aSakaguchi et al. (1991).

The anatomical and physiologic adaptations that permit colobines to support microbial fermentation of plant fiber and to utilize the end-products of that fermentation limit their ability to adapt to more digestible diets. The VFAs produced as a principle end-product of the fermentation process have a pK_a of 4.8. The symbiotic microorganisms responsible for VFA production cannot survive at a pH below 5.5 (Stevens, 1982). The pH of the colobine saccus contents is thought to be maintained in a manner similar to that of the rumen contents, i.e., by rapid absorption of VFAs and secretion of HCO₃ as a buffer by the salivary glands, pancreas, and other digestive

tract cells. The extent to which these mechanisms are present in the colobine saccus has yet to be determined (Kay and Davies, 1994).

If a colobine primate consumes excessive amounts of simple or readily fermentable carbohydrates (starch, sugar), the result would include rapid formation of VFAs in the foregut. This rapid production of VFAs in ruminants produces a dramatic drop in the pH of the digesta, increased absorption of VFAs, and replacement of the normal microbial organisms with lactobacilli. Lactobacilli produce lactic acid that may result in formation of hypertonic digesta, pulling water into the rumen and producing systemic dehydration. D-lactic acid produced in the forestomach is absorbed into the bloodstream, where it is only slowly metabolized, and leads to a systemic acidosis. Some of these changes have been observed in colobines receiving a concentrate-type diet (Bauchop and Martucci, 1968) or selecting a diet high in seeds (Ohwaki et al., 1974).

This rapid production of VFAs can produce necrosis and ulceration of the rumen epithelium, rumen immotility, and systemic acidosis, all of which are signs commonly observed in captive colobines (Janssen, 1994). Additionally, acute acidosis causes proliferation of coliforms and clostridial species that may produce enterotoxins and contribute to diarrhea (Church, 1988).

Eisenberg's (1978) categorizations of primate herbivory (Chapter I) would place both African and Asian colobines in Class 5, species that are close to being "obligate" folivores. Certainly, within such a broad grouping, there will be variability. As with Alouatta, there is strong evidence from field data that some colobines may utilize plant parts that are relatively more digestible when they are seasonally available.

Markers

Markers may be used to measure the movement of digesta through the gastrointestinal tract as well as indirect indicators of food digestibility. Those markers used for the purpose of estimating digestibility are typically fed or administered at a constant level in the diet (Van Soest et al., 1983). Acid-detergent lignin has been used as an internal indicator of food digestibility (Van Dyne, 1968; Ullrey et al., 1987). Application of this method assumes that the lignin fraction of the diet is totally indigestible by the host and the gastrointestinal microflora. Van Dyne (1968) stated that the apparent digestibility (AD) of nutrients may be determined, using the following lignin-ratio technique:

AD = 100- (% Nutrient in feces) (% Lignin in ingesta)
(% Nutrient in ingesta) (% Lignin in feces)

The primary limitations of this method lie in the assumption of lignin indigestibility and the analytical error involved in lignin determination. These limitations of lignin analysis are well documented (Van Soest, 1994). To use lignin as a marker for estimates of food or nutrient digestibility, it has been stated that the dietary lignin concentration should be at least 6.0% (Van Soest, 1995).

Relatively high apparent digestibility of acid detergent lignin was observed in this study (39.7-68.9%). Similar results were reported in feeding trials with black and white colobus (Watkins et al., 1985; Oftedal et al., 1982). These values appear to originate from a combination of low lignin concentrations (< 2%) in the test diets used in these trials and the analytical errors involved in lignin determination previously described.

Particulate Markers

Concerns surrounding the application of particulate markers, such as acetate beads, seeds, and glass balls, have been expressed by several authors. The primary argument against these markers is their potential to traverse the gastrointestinal tract in a manner different from the digesta. Some of this deviation from natural movement is a result of differences in specific gravity. Suitable markers should behave as part of the digesta the investigator wishes to mark,

without producing other effects. However, the suitablility of a marker is related, in part, to how it is used. For example, the acetate bead may be a suitable marker for solid phase digesta within species lacking a complex gastrointestinal tract. In animals with gastrointestinal sacculations and folds, acetate beads may become entrapped and separate from the remainder of the digesta. The type of food offered during the trial and the desired measurement (e.g., transit time) will dictate which marker may be successfully applied. Finally, the facilities and equipment available to analyze the excreted marker will influence the marker choice.

Inorganic Compounds

It has been demonstrated that liquid and particulate matter flow differently through the gastrointestinal tract. Large ungulates, ruminants, horses, and pigs tend to retain fiber (solid) relative to liquid while some rodents, and rabbits selectively retain liquid relative to fiber (solid). The two inorganic markers most commonly used to measure these fractions, Cr-fiber (solid) and Co-EDTA (liquid), have been applied to numerous species. Comparative passage rates of liquid and solids in some animals, including the results of this study, are presented in Table 3-30. The lack of statistical differences between the rates determined for the two markers in this study appear to be a function of the

Comparative passage rates of solid and liquid Table 3-30. digesta in species with simple GIT, hindgut, and foregut fermentation (Udén, 1978; Van Soest et al., 1983).

Species	Diet	BW	Whole tract re	etention (R _{git})
	typeª	(kg)	Solid (h)	Liquid (h)
SIMPLE GIT				
Ruffed lemur	1	4.7	3.8	4.1
Ruffed lemur	2	4.0	2.3	2.4
Man	4	70	41	39
HINDGUT				
howler	1	7.5	38.8	34.1
howler	2	7.6	38.8	40.4
horse	3	388	21	17
pony	3	132	25	16
rabbit	3	3	5	193 ^b
FOREGUT				
colobine	1	7.5	37.6	41.1
colobine	2	7.5	68.6	38.4
cattle	3	243	48	20
sheep	3	30	57	26
goat	3	29	41	28

al=15ADF.

sample size.

In general, there was an increase in passage rate (decreased retention) of both phases of digesta seen in ruffed lemurs when fed 30ADF as compared to 15ADF. This would

²⁼³⁰ADF.

³⁼Timothy hay.

⁴⁼Standard diet including 20 g of dietary fiber from wheat bran.
bProlonged retention results form selective retention of liquid as well as selective coprophagy of the less fibrous night feces.

further support the previously described digestive strategies of these two subspecies.

The approximately equal retention times for solid and liquid digesta exhibited by howlers are similar to the relative rates $R_{\rm git}$ reported for other hindgut, noncoprophagic species. Species with foregut fermentation exhibit a more rapid passage of liquid than solid digesta. This is consistent with relative liquid and solid retention times $(R_{\rm git})$ seen in colobines fed 30ADF; however, when fed 15ADF, the solid digesta $R_{\rm git}$ appears to be appreciably shorter.

CONCLUSION

The gastrointestinal and dietary adaptations that are well known in domestic herbivores also are seen in leaf-eating primates. Rapid rates of passage and reduced digesta retention are particularly evident in herbivorous species such as red ruffed lemurs and black and white ruffed lemurs that lack specific gastrointestinal tract modifications for microbial fermentation of plant fiber. This results in a reduced ability to digest ingested plant material and requires increased rates of intake. One would predict the time spent feeding by these species, as a percent of daily activity would be greater than that of other leaf-eating primates.

Increased retention times and reduced rates of passage are observed in leaf-eating primates with adaptations for

either hindgut or foregut fermentation. Digestive efficiencies of hindgut fermenters (mantled howler, red howler, and black howler) are significantly less than those of reduced digestive efficiencies colobines. These comparable to those seen when contrasting horse and cattle and appear to be related to the lack of extensive gastrointestinal modifications anterior to the sites of gastric and enzymatic digestion. Hindgut fermenters also have a reduced absorptive surface area posterior to the site of microbial fermentation.

Colobine primates (black and white colobus, Francois' langur, douc langur) exhibit the most extensive gastrointestinal modifications for microbial fermentation of plant cell wall. Although there is a diversity of feeding habits among these species, foregut fermentation is present throughout the subfamily. As a result, regardless of the type of plant material consumed, that food is subjected to fermentative processes.

Primates with either hindgut or foregut fermentation subjected to thousands of years of evolutionary pressure, have developed mechanisms to accommodate readily available plant food and effectively degrade plant fiber, yielding a significant amounts of energy through symbiotic relationships with gut microorganisms. The unfortunate consequences of presenting rapidly fermentable foods to these primates are illustrated in much of the literature concerned with their

captive husbandry. The impression that plant fiber is a negative component in diets, and that a diet low in this fraction is "preferred" by these captive primates, has produced many of the health problems commonly seen.

It is apparent, from these and other studies that plant fiber should be considered with equal importance to "true" nutrients when formulating diets for herbivorous primates.

LITERATURE CITED

Literature Cited

Aldrich-Blake, F.P.G. 1970. Problems of social structure in forest monkeys. In: Social behaviour in birds and mammals (J.H. Crook, ed.). Pp. 79-99. Academic Press, New York, NY.

Allen, M.S., and D.B. Mertens. 1988. Evaluating constraints on fiber digestion by rumen microbes. Journal of Nutrition 118:261-270.

Altmann, J. 1974. Observational study of behavior: sampling methods. Behavior 49:227-267.

Amerasinghe, F.P., B.W.B. Van Cuylenberg, and C.M. Hladik. 1971. Comparative histology of the alimentary tract of Ceylon primates in correlation with diet. Ceylon Journal of Science and Biological Science 9:75-87.

Applebaum, S.W., and Y. Birk. 1979. Saponins. In: Herbivores: Their Interactions with Secondary Plant Metabolites (G.A. Rosenthal and D.H. Janzen, eds.). Pp. 539-566. Academic Press, New York, NY.

Association of Official Analytical Chemists. 1990. Official Methods of Analysis. (15th ed.) AOAC, Washington, D.C.

Baranga, D. 1982. Nutrient composition and food preferences of colobus monkeys in Kibale Forest, Uganda. African Journal of Ecology 20:113-121.

Bauchop, T. 1971. Stomach microbiology of primates. Annual Review of Microbiology 25:429-436.

Bauchop, T. 1978. Digestion of leaves in vertebrate arboreal folivores. In: The Ecology of Arboreal Folivores (G.G. Montgomery, ed.). Pp. 193-204. Smithsonian Institution Press, Washington, D.C.

Bauchop, T., and R.W. Martucci. 1968. Ruminant-like digestion of the langur monkey. Science 161:698-700.

Beck, B.B., and R. Tuttle. 1972. The behavior of the grey langurs at a Ceylonese waterhole. In: The Functional and

Evolutionary Biology of Primates (R. Tuttle, ed.). Aldine-Atherton. Chicago, IL.

Benirschke, K., and M. Bogart. 1978. Pathology of douc langurs (*Pygathrix nemaeus*) at San Diego Zoo. Erkrankungen der Zootiere 257-261.

Benton, L., Jr. 1976. The establishment and husbandry of a black howler, *Alouatta caraya*, colony at Columbia Zoo. International Zoo Yearbook 16:149-152.

Bicca-Marques, J.C., and C. Calegaro-Marques. 1993. Feeding postures in the black howler monkey, *Alouatta caraya*. Folia Primatologica 60:169-172.

Blaxter, K.L., N. McGraham, and F.W. Wainman. 1956. Some observations on the digestibility of food by sheep and on related problems. British Journal of Nutrition 10:69.

Bodini, R. and R. Perez-Hernandez. 1987. Distribution of the species and subspecies of Cebids in Venezuela. Fieldiana: Zoology 39:231-244.

Boyne, A.W.R., R.M. Campbell, J. Davidson, and D.P. Cuthbertson. 1956. Changes in composition of the digesta along the alimentary tract of sheep. British Journal of Nutrition 10:325-333.

Braza, F., F. Alvarez, and T. Azcarate. 1983. Feeding habits of the red howler monkeys (*Alouatta seniculus*) in the llanos of Venezuela. Mammalia 47:205-214.

Brown, A.J., and P.E. Kolattukudy. 1978. Evidence that pancreatic lipase is responsible for the hydrolysis of cutin, a biopolyester present in mammalian diet, and the role of bile salt and colipase in this hydrolysis. Archives of Biochemistry and Biophysics 190:17-26.

Carpenter, C.R. 1934. A field study of the behavior and social relationship of howling monkeys (*Alouatta palliata*). Comparative Psychology Monographs 19:1-168.

Carvalho, C.T. 1975. Acerca da alimentacao dos bugios (Mammalia, Cebidae). Silviculture em Sao Paulo 9:53-56.

Chalmers, N.R. 1968. Group composition, ecology, and daily activities of free living mangabeys in Uganda. Folia Primatologica 8:247-262.

Cheeke, P.R. 1976. Nutritional and physiological properties of

saponins. Nutrition Reports International 13:315-324.

Chivers, D.J. 1969. On the daily behavior and spacing in howling monkey groups. Folia Primatologica 10:48-102.

Chivers, D.J. 1994. Functional anatomy of the gastrointestinal tract. In: Colobine monkeys: their ecology, behaviour, and evolution (A.G. Davies and J.F. Oates, eds.). Pp 205-228. Cambridge University Press, Cambridge.

Chivers, D.J., and C.M. Hladik. 1980. Morphology of the gastrointestinal tract in primates: Comparisons with other mammals in relation to diet. Journal of Morphology 166:337-386.

Church, D.C. 1988. The Ruminant Animal. (D.C. Church, ed.). Prentice Hall. Englewood Cliffs, NJ.

Clutton-Brock, T.H. 1975. Feeding behaviour of red colobus and black and white colobus in East Africa. Folia Primatologica 23:165-207.

Collins, L., and M. Roberts. 1978. Arboreal folivores in captivity: Maintenance of a delicate minority. In: The Ecology of Arboreal Folivores (G.G. Montgomery, ed.). Pp. 193-204. Smithsonian Institution Press, Washington, DC.

Crandall, L.S. 1964. Management of Wild Mammals in Captivity. The University of Chicago Press, Chicago, IL.

Crissey, S.D., and M.S. Edwards. 1989. Telazol immobilization of red howlers (*Alouatta seniculus*) under free-ranging conditions. AAZV Annual Conference Proceedings. Greenville, NC.

Crissey, S.D., M.S. Edwards, O.T. Oftedal, J. Curier and R. Rudran. 1989. The role of fiber in natural and manufactured diets fed to red howler monkeys (*Alouatta seniculus*). In: Proceedings of the Dr. Scholl Conference on Zoo Nutrition. Pp. 135-147. Chicago, IL. 8-9 December, 1989.

Crockett, C.M. 1985. Population studies of red howler monkeys (Alouatta seniculus). National Geographic Research 1:264-273.

Cummings, J.H. 1978. Nutritional implications of dietary fiber. American Journal of Clinical Nutrition 31:S21-S29.

Davies, A.G., and I.C. Baillie. 1988. Soil-eating by red leaf monkeys (*Presbytis rubicunda*) in Sabah, northern Borneo. Biotropica 20:252-258.

- Deinum, B. 1971. Prediction of forage digestibility from some laboratory procedures. 2. Comparison of *in vivo* digestibility at two institutions. Netherlands Journal of Agricultural Science 19:106-113.
- Dierenfeld, E.S., H.F. Hintz, J.B. Robertson, P.J. Van Soest, and O.T. Oftedal. 1982. Utilization of bamboo by the giant panda. Journal of Nutrition 112:636-641.
- Dierenfeld, E.S., F.W. Koontz, and R.S. Goldstein. 1992. Feed intake, digestion and passage of the proboscis monkey (*Nasalis larvatus*) in captivity. Primates 33:399-405.
- Edwards, M.S. 1990. Using natural diet composition when formulating captive animal diets: fiber concentrations. In: Proceedings of the American Association of Zoological Parks and Aquariums. Pp. 696-700. Wheeling, WV.
- Eisenberg, J.F. 1978. The evolution of arboreal herbivores in the class Mammalia. In: The Ecology of Arboreal Folivores (G.G. Montgomery, ed.). Pp. 135-152. Smithsonian Institution Press, Washington, DC.
- Eisenberg, J.F., N.A. Muckenhirn, and R. Rudran. 1972. The relation between ecology and social structure in the primates. Science 176:863-874.
- Eisenberg, J.F., and R.W. Thorington. 1973. A preliminary analysis of Neotropical mammal fauna. Biotropica 5:150-161.
- Elseden, S.R., M.W.S. Hitchcock, R.A. Marshall, and A.T. Phillipson. 1946. Volatile acid in the digesta of ruminants and other animals. Journal of Experimental Biology 22:191-202.
- Ensley, P.E., T.L. Rost, M. Anderson, K. Benirschke, D. Brockman, and D.E. Ullrey. 1982. Intestinal obstruction and perforation caused by undigested *Acacia* sp. leaves in langur monkeys. Journal of the American Veterinary Medical Association 181:1351-1354.
- Feeny, P. 1970. Seasonal changes in oak leaf tannins and nutrients as a cause of spring feeding by winter moth caterpillars. Ecology 51:565-580.
- Fenton, T.W., and M. Fenton. 1979. An improved procedure for the determination of chromic oxide in feed and feces. Canadian Journal of Animal Science 59:631-634.
- Flower, S.S. 1931. Contributions to our knowledge of the duration of life in vertebrate animals. 5. Mammals. Pp.145-

- 234. Proceedings of the Zoological Society of London.
- Fooden, J. 1964. Stomach contents and gastrointestinal proportions in wild shot Guianan monkeys. American Journal of Physical Anthropology 22:227-232.
- Foose, T.J. 1982. Trophic Strategies of Ruminant versus Nonruminant Ungulates. Ph.D. Dissertation. Department of Biology, University of Chicago, Chicago, IL.
- Galetti, M., R. Laps, and F. Pedroni. 1987. Feeding behavior of the brown howler monkey (*Alouatta fusca clamitans*) in a forest fragment in state of Sao Paulo, Brazil. International Journal of Primatology 8:542.
- Gaulin, S.J.C., and C.K. Gaulin. 1982. Behavioral ecology of Alouatta seniculus in Andean cloud forest. International Journal of Primatology 3:1-32.
- Gaulin, S.J.C., D.H. Knight, and C.K. Gaulin. 1980. Local variance in *Alouatta* group size and food availability on Barro Colorado Island. Biotropica 12:137-143.
- Gautier-Hion, A. 1983. Leaf consumption by monkeys in western and eastern Africa: a comparison. African Journal of Ecology 21:107-113.
- Glander, K.E. 1975. Habitat and resource utilization: An ecological view of social organization in mantled howling monkeys. Ph.D. Dissertation. Department of Anthropolgy, University of Chicago, Chicago, IL.
- Glander, K.E., and M.F. Teaford. 1995. Seasonal, sexual, and habitat effects on feeding rates of Alouatta palliata. American Journal of Primatology 36:124.
- Goering, H.K., and P.J. Van Soest. 1970. Forage Fiber Analyses (Apparatus, Reagents, Procedures, and Some Applications). Agricultural Handbook 379. U.S. Department of Agriculture, Washington, DC.
- Griner, L.A. 1977. Nutritional, veterinary and breeding requirements of the East Bornean Proboscis monkey within a controlled environment. In: AAZV Annual Conference Proceedings. Pp. 48-59. San Diego, CA.
- Harbourne, J.B. 1977. Introduction to Ecological Biochemistry. Academic Press. New York, NY.
- Harbourne, J.B. 1979. Flavonoid Pigments. In: Herbivores:

Their Interactions with Secondary Plant Metabolites (G.A. Rosenthal and D.H. Janzen, eds.). Pp. 619-655. Academic Press. New York, NY.

Hargreaves, D., and B. Hargreaves. 1960. Tropical Blossoms of the Caribbean. Hargreaves Company, Inc., Kailua, HI.

Hargreaves, D., and B. Hargreaves. 1965. Tropical Trees found in the Caribbean, South America, Central America, and Mexico. Hargreaves Company, Inc., Kailua, HI.

Harrison, M.J.S., and C.M. Hladik. 1986. Un primate granivore: Le Colobe noir dans la foret du Gabon; Potentialite d'evolution du comportement. Terre Vie 41: 281-298.

Heldstab, A. 1988. Management and disease problems in douc langurs at the Basle Zoo. AAZV Annual Conference Proceedings. Pp. 184-187. Toronto, Ontario.

Hernandez-Camacho, J., and R.W. Cooper. 1976. The non-human primates of Colombia. In: Neotropical Primates: field studies and conservation (R.W. Thorington and P.G. Heltne, eds.). Pp. 35-69. National Academy of Sciences, Washington, DC.

Hershkovitz, P. 1969. The evolution of mammals on southern continents, VI: The recent mammals of the Neotropical region, a zoogeographic and ecological review. Quarterly Review of Biology 44:1-70.

Hick, U. 1972. Breeding and maintenance of douc langurs at Cologne Zoo. International Zoo Yearbook 12:98-103.

Hill, W.C.O. 1958. Pharynx, oesophagus, stomach, small and large intestine: form and position. Primatologia 3:139-207.

Hill, W.C.O. 1962. Primates: Comparative anatomy and taxonomy, IV. Intersciences: New York, NY.

Hill, W.C.O. 1964. The maintenance of langurs (Colobidae) in captivity: Experiences and some suggestions. Folia Primatologica 2:222-231.

Hintz, H.F. 1969. Effect of coprophagy on digestion and mineral excretion in the guinea pig. Journal of Nutrition 99:375-378.

Hintz, H.F., A.J. Williams, J. Rogoff, and H.F. Schryvwer. 1973. Availability of phosphorus in wheat bran when fed to ponies. Journal of Animal Science 36:522-525.

Hladik, C.M. 1978. Adaptive strategies of primates in relation to leaf-eating. In: The Ecology of Arboreal Folivores (G.G. Montgomery, ed.). Pp. 373-395. Smithsonian Institution Press, Washington, DC.

Hladik, A., and C.M. Hladik. 1969 Rapports tróphiques entre végétation et Primates dans la fôret de Barro Colorado (Panama). La Terre et la Vie 1:25-117.

Hladik, C.M., and L. Guegen. 1974. Geophagie et nutrition minerale chez les primates sauvages. Comptes Rendus hebdomadaires des Seances de L'Academie des Sciences Paris Series D 279:1393-1396.

Hoffmann, R.R. 1973. The Ruminant Stomach: Stomach structure and feeding habits of East African game ruminants. East African Literature Burea, Nairobi, Kenya.

Hollihn, U. 1973. Remarks on the breeding and maintenance of Colobus monkeys (*Colobus guereza*), Proboscis monkeys (*Nasalis larvatus*), and Douc langurs (*Pygathrix nemaeus*) in Zoos. International Zoo Yearbook 13:185-188.

Hungate, R.E., G.D. Phillips, A. McGregor, D.P. Hungate, and H.K. Buechner. 1959. Microbial fermentation in certain mammals. Science 130:1192-1194.

Izawa, K. 1975. Foods and feeding behavior of monkeys in the upper Amazon basin. Primates 16:295-316.

Izawa, K. 1993. Soil eating by *Alouatta* and *Ateles*. International Journal of Primatology 14:229-242.

Izawa, K., and H.M. Lozano. 1990. Frequency of soil-eating by group of wild howler monkeys (*Alouatta seniculus*) in La Macarena. Field Studies of New World Monkeys, La Macarena, Columbia 4:47-56.

Jaffe, W.G. 1976. Selenium-tellurium in the environment. Pp. 88. Industrial Health Foundation. Pittsburgh, PA.

Janssen, D.L. 1994. Morbidity and mortality of douc langurs (*Pygathrix nemaeus*) at the San Diego Zoo. In: AAZV Annual Conference Proceedings. St. Louis, MO.

Jolly, A. 1972. The Evolution of Primate Behavior. MacMillan Company. New York, NY.

Jones, M.L. 1995. Personal communication.

Julliot, C., and D. Sabatier. 1993. Diet of red howler monkey (Alouatta seniculus) in French Guiana. International Journal of Primatology 14:527-550.

Kawabe, M., and T. Mano. 1972. Ecology and behavior of the wild proboscis monkey, *Nasalis larvatus* (Wurmb), in Sabah, Malaysia. Primates 13:213-228.

Kay, R.N.B., and A.G. Davies. 1994. Digestive physiology. In: Colobine monkeys: their ecology, behaviour, and evolution (A.G. Davies and J.F. Oates, eds.). Pp 229-250. Cambridge University Press, Cambridge.

Kay, R.N.B., P. Hoppe, and G.M.O. Maloiy. 1976. Fermentative digestion of food in the colobus monkey, *Colobus polykomos*. Experimentia 32(4):485-487.

Kern, J.A. 1964. Observations on the habits of the proboscis monkey, *Nasalis larvatus*. (Wurmb), made in the Brunai Bay area, Borneo. Zoologica 49:183-192.

Kleiber, M. 1961. The Fire of Life. John Wiley & Sons Inc. New York, NY.

Koontz, F.W., E.S. Dierenfeld, and R. Goldstein. 1988. Nutrition management of captive proboscis monkeys (*Nasalis larvatus*) at the New York Zoological Park. AAZPA Regional Conference Proceedings. Pp. 701-711. Wheeling, WV.

Kotb, A.K., and T.D. Luckey. 1972. Markers in nutrition. Nutrition Abstract Review 42:813-845.

Kuhn, H.J. 1964. Zur Kenntnis van Bau und Funktion des Magens der Schlankaffen (Colobinae). Folia Primatologica 2:193.

Langer, P. 1988. The Mammalian Herbivore Stomach: Comparative Anatomy, Function and Evolution. Gustav Fischer Veralg, New York, NY.

Levin, D. 1971. Plant phenolics: an ecological perspective. American Naturalist 105:157-181.

Levin, D. 1976. The chemical defenses of plants to pathogens and herbivores. Annual Review of Ecological Systematics 7:121-159.

Lindbergh, S. 1976. Natural social structures and feeding procedure in the acclimatization of South American primates. International Zoo Yearbook 16:146-149.

Lippold, L.K. 1977. The douc langur: A time for conservation. In: Primate Conservation (G. Bourne and Prince Rainier, eds). Pp. 513-538. Academic Press. New York, NY.

Lohan, O.P., D. Lall, H.P.S. Makkar, and S.S. Negi. 1981. Inhibition of rumen urease activity by tannins in oak leaves. Indian Journal of Animal Science 51:279-281.

Ludwig, V.W. 1989. "On nutrition and artificial diets of *Colobus guereza* at the Dresden Zoological Garden." Zoological Garten 59(3):173-187 (in German).

Mabry, T.J., and J.E. Gill. 1979. Sesquiterpene lactones and other terpenoids. In: Herbivores: Their Interactions with Secondary Plant Metabolites (G.A. Rosenthal and D.H. Janzen, eds.). Pp. 502-537. Academic Press. New York, NY.

MacKey, D. 1978. Soil, vegetation and seed-eating by black colobus monkeys. In: The Ecology of Arboreal Folivores (G.C. Montgomery, ed.). Pp. 423-439. Smithsonian Institution Press, Washington, DC.

Mandels, M., and E.T. Reese. 1963. Advances in Enzymic Hydrolysis of Cellulose and Related Materials. (E.T. Reese, ed.) Pergamon Press. New York, NY.

Mannchen, K. 1989. Feeding exotic tropical browse to lowland gorillas (*Gorilla gorilla gorilla*) at the Miami Metro Zoo. AAZPA Regional Conference Proceedings. Pp. 493-499.

Martz, F.A., P.J. Van Soest, J.R. Vogt, and E.S. Hildebrand. 1974. Use of elemental tracers and activation analysis in digestion, rate of digest flow and food particle tracking studies in cattle. Proceedings of the 6th Symposium on Energy Metabolism. Stuttgardt, Germany.

McCandlish, A.C. 1937. Wiss. Ber. Milchwiltsch. Weltkongr., 11th, vol. 1. Pp. 410-412.

McClure, H.M., W.L. Chapman, and B.E. Hooper. 1978. The Digestive System. In: Laboratory Animals (K. Benirschke, F.M. Garner, and Jones, eds.) Pp. 203-204. Springer-Verlag. New York, NY.

Merrit, D.A., Jr. 1982. Behavioural considerations in the acclimatization and nutrition of young black howlers (*Alouatta caraya*). International Zoo Yearbook 22:128-131.

Milton, K. 1978a. Behavioral adaptations to leaf-eating by the mantled howler monkey (Alouatta palliata). In: The Ecology of

Arboreal Folivores (G.G. Montgomery, ed.). Pp. 205-229. Smithsonian Institution Press, Washington, DC.

Milton, K. 1978b. The quality of diet as a possible limiting factor on the Barro Colorado Island howler monkey population. In: Proceedings of the Sixth Congress of the International Primate Society. Academic Press, Cambridge.

Milton, K. 1980. The Foraging Strategy of Howler Monkeys: a Study in Primate Economics. Columbia University Press, New York, NY.

Milton, K. 1981. Food choice and digestive strategies of two sympatric primate species. The American Naturalist 117:496-505.

Milton, K. 1984. The role of food-processing factors in primate food choice. In: Adaptations for foraging in nonhuman primates (P.S. Rodman and J.G.H. Cant, eds.). Pp. 249-279. Columbia University Press, NY.

Milton, K., and R.H. McBee. 1983. Rates of fermentative digestion in the howler monkey, *Alouatta palliata* (Primates: Ceboidea). Comparative Biochemistry and Physiology 74A:29-31.

Milton, K., P.J. Van Soest, and J.B. Robertson. 1980. Digestive efficiencies of wild howler monkeys. Physiological Zoology 53:402-409.

Mitchell, P.C. 1911. On longevity and relative viability in mammals and birds; with a note on the theory of longevity. Proceedings of the Zoological Society of London. Pp. 425-548

Moir, R.J. 1968. Ruminant digestion and evolution. In: Handbook of Physiology. Pp. 2673-2694. American Physiology Society. Washington, DC.

Moreno-Black, G. 1978. The use of scat samples in primate diet analysis. Primates 19:215-221.

Morin, M.L., D.M. Renquist, J. Knapka, and F.J. Judge. 1978. The effect of dietary crude fiber levels on rhesus monkeys during quarantine. Laboratory Animal Science 28:405-411.

Nagy, K.A., and K. Milton. 1979. Energy metabolism and food consumption by wild howler monkeys. Ecology 60:475-480.

National Research Council. 1978. Nutrient Requirements of Nonhuman Primates. National Academy of Sciences, Washington, DC.

- National Research Council. 1988. Nutrient Requirements of Swine. National Academy of Sciences, Washington, DC.
- National Research Council. 1989. Nutrient Requirements of Horses. National Academy of Sciences, Washington, DC.
- Nevens, W.B. 1928. Effects of fasting and the method of preparation of feed upon the digestive process in dairy cattle. Journal of Agricultural Research 36:777-794.
- Newton, P. 1992. Feeding and ranging patterns of forest Hanuman langurs (*Presbytis entellus*). International Journal of Primatology 13:245-285.
- Nhat, P., X. Mai, and H. Tay. 1993. First results on the diet of the red-shanked douc langur, *Pygathrix nemaeus*. Australian Primatology 8.
- Nimz, H. 1974. Angewande Chemie International Edition in English 13:313-322.
- Oates, J., T. Swain, and J. Zantovska. 1977. Secondary compounds and food selection by colobus monkeys. Biochemical Systematics and Ecology 5:317-321.
- Oftedal, O.T. 1991. The nutritional consequences of foraging in primates: the relationship of nutrient intakes to nutrient requirements. Philosophical Transcripts of the Royal Society of London B. 334:161-170.
- Oftedal, O.T., M. Jakubasz, and P. Whetter. 1982. Food intake and diet digestibility by captive black and white colobus (*Colobus guereza*) at the National Zoological Park. In: AAZV Annual Conference Proceedings. Pp. 33.
- Ohwaki, K., R.E. Hungate, L. Lotter, R.R. Hoffman, and G. Maloiy. 1974. Stomach fermentation in East African colobus monkeys in the natural state. Applied Microbiology 27(1):713-723.
- Olsson, N.O. 1969. The nutrition of the horse. In: International Encyclopedia of Food and Nutrition, Vol 17, part 2 (D. Cuthbertson, ed.). Pp. 921-960. Pergamon Press, Oxford.
- Oxnard, C.E. 1966. Vitamin B_{12} nutrition in some primates in captivity. Folia Primatologica 4:424-431.
- Oxnard, C.E. 1969. A note on the ruminant-like digestion of langurs. Laboratory Primate Newsletter 8:24-25.

Parfitt, A.M., B.A. Higgins, J.R. Nassim, J.A. Collins, and A. Hilb. 1964. Metabolic studies in patients with hypercalciuria. Clinical Science 27:463-482.

Parra, R. 1978. Comparison of foregut and hindgut fermentation in herbivores. In: The Ecology of Arboreal Folivores (G.G. Montgomery, ed.). Pp. 205-229. Smithsonian Institution Press. Washington, DC.

Parra, R., J. Combellas, and E. Gonzalez. 1972. Composicion y valor nutritivo de forrajes producidos en el trópico. 2. Fracciones quimicas que afectan la disponibilidad de los componenetes fibrosos. Agronomiá Tropical 22:219-230.

Pereira, M.E., M.L. Seeligson, and J.M. Macedonia. 1988. The behavioral repertoire of the black-and-white ruffed lemur, *Varecia variegata variegata* (Primates: Lemuridae). Folia Primatologica 51:1-32.

Petter, J.J., R. Albinac, and Y. Rumpler. 1977. Faune de Madagascar, vol. 44. Mammiferes lemuriens. Orstom. Paris.

Pope, B.L. 1966. The population characteristics of howler monkeys (*Alouatta caraya*) in northern Argentina. American Journal of Physical Anthropology 24:361-370.

Preston, R.C. 1977. Phosphorus in beef cattle and sheep nutrition. In: NFIA Literature Review on Phosphorus in Ruminant Nutrition. Pp. 1-44. National Feed Ingredients Association. West Des Moines, IA.

Quin, J.I. 1943. Studies on the alimentary tract of Merino sheep in South Africa VIII.—the pathogenesis of acute tympanites. Onderstepoort Journal of Veterinary Science and Animal Industry 18:113-117.

Racenis, J. 1952. Some observations on the red howling monkey (Alouatta seniculus) in Venezuela. Journal of Mammalogy 33:114-115.

Ramia, M. 1974. Plantas de las Sabanas Llaneras. Monte Avila, Caracas, Venezuela.

Recommended Dietary Allowances. 1989. 10th ed. National Research Council. National Academy Press. Washington, D.C.

Rhoades, D.F., and R.G. Cates. 1976. Toward a general theory of plant anti-herbivore chemistry. Recent Advances in Phytochemistry 10:168-213.

Richard, A. 1970. A comparative study of the activity patterns and behavior of *Alouatta villosa* and *Ateles geoffroyi*. Folia Primatologica 12:241-263.

Robbins, C.T., and A.N. Moen. 1975. Composition and digestibility of several deciduous browses in the northeast. Journal of Wildlife Management 39:337-341.

Robbins, C.T., P.J. Van Soest, W.W. MAutz, and A.N. Moen. 1975. Feed analysis and digestion with reference to white-tailed deer. Journal of Wildlife Management 39:67-79.

Robertson, J.B., and P.J. Van Soest. 1977. Dietary fiber estimation in concentrate feedstuffs. Journal of Animal Science 45:456.

Robinson, P.T., T.A. Reichard, P.A. Whetter and D.E. Ullrey. 1982. Evaluation of diets of leaf-eating monkeys. p. 36. AAZV Annual Proceedings. New Orleans, LA.

Robinson, T. 1979. The evolutionary ecology of alkaloids. In: Herbivores: Their Interactions with Secondary Plant Metabolites (G.A. Rosenthal and D.H. Janzen, eds.). Pp. 413-448. Academic Press, New York, NY.

Rosenberger, A.L. 1992. Ecolution of feeding niches in New World monkeys. American Journal of Physical Anthropolgy 88:525-562.

Rotruck, J.T., A.L. Pope, H.E. Ganther, A.B. Swanson, D.G. Hafeman, and W.G. Hoekstra. 1973. Selenium: Biochemical role as a compound of glutathione peroxidase. Science 179:588.

Rudran, R. 1979. The demography and social mobility of a red howler (*Alouatta seniculus*) population in Venezuela. In: Vertebrate Ecology in the Northern Neotropics (J.F. Eisenberg, ed.). Pp. 107-126. Smithsonian Institution Press. Washington, DC.

Sakaguchi, E., K. Suzuki, S. Kotera, and A. Ehara. 1991. Fiber digestion and digesta retention time in macaque and colobus monkeys. In: Primatology Today: Proceedings of the 13th Congress International Primatology Society. (A. Ehara, T. Kuimura, O. Takenaka, and M. Iwamoto, eds.). Pp. 671-674. Elsevier, England.

Salter, R.E, N.A. Mackenzie, N. Nightingale, K.M. Aken and P.K. Chai. 1985. Habitat use, ranging behaviour and food habits of the proboscis monkey, *Nasalis larvatus* (van Wurmb) in Sarawak. Primates 26:436-451.



- Santi, C., and D. Santi. 1989. Toxic plants: tropical and temperate. AAZPA Regional Conference Proceedings. Pp. 990-997.
- Shaver, T.N., and M.J. Lukefahr. 1969. Journal of Economic Entomology 62:643-646.
- Slavin, J.L., P.M. Brauer, and J.A. Marlett. 1981. Neutral detergent fiber, hemicellulose and cellulose digestibility in human subjects. Journal of Nutrition 111:287-297.
- Smith, C.C. 1977. Feeding behaviour and social organization in howling monkeys. In: Primate Ecology (T.H. Clutton-Brock, ed.) Pp. 97-126. Academic Press, London.
- Smith, J.D. 1970. The systematic status of the black howler monkey, *Alouatta pigra* Lawrence. Journal of Mammalogy 51:358-369.
- Stanyon, R., S. Tofanello, M.A. Morescalchi, G. Agoramoorthy, O.A. Ryder, and J. Wienberg. 1995. Cytogenetic analysis shows extensive genomic rearrangements between red howler (*Alouatta seniculus* Linnaeus) subspecies. American Journal of Primatology 35:171-183.
- Stevens, C.E. 1982. Comparative anatomy and physiology of the herbivore digestive tract. In: Proceedings of the Second Annual Dr. Scholl Conference on the Nutrition of Captive Wild Animals (T.P. Meehan, B.A. Thomas, and K. Bell). Pp. 8-16. Lincoln Park Zoological Society. Chicago, IL.
- Stevens, C.E. 1988. Comparative Physiology of the Vertebrate Digestive System. Cambridge University Press, New York, NY.
- Stowe, H.D. 1967. Serum selenium and related parameters of naturally and experimentally fed horses. Journal of Nutrition 93:60-64.
- Sullivan, J.T. 1966. Studies of the hemicelluloses of forage plants. Journal of Animal Science 25:83-86.
- Swain, T. 1977. Secondary compounds as protective agents. Annual Review of Plant Physiology 28:479-501.
- Swain, T. 1978. Phenolics in the environment. Recent Advances in Phytochemistry 12:617-640.
- Taff, M.A., and P. Dolhinow. 1989. Langur monkeys (*Presbytis entellus*) in captivity. In: Housing, Care and Psychological Wellbeing of Captive and Laboratory Primates (E.F. Segal, ed.). Pp. 291-304. Noyes Publications, Park Ridge, NJ.

- Tattersall, I. 1982. The Primates of Madagascar. Columbia University Press, New York, NY.
- Troth, R.G. 1979. Vegetational types on a ranch in the central llanos of Venezuela. In: Vertebrate Ecology in the Northern Neotropics (J.F. Eisenberg, ed.) Smithsonian Institution Press, Washington, DC.
- Udén, P. 1978. Comparative studies on rate of passage, particle size and rate of digestion in ruminants, equines, rabbits and man. Ph.D. Dissertation. Department of Animal Science. Cornell University, NY.
- Udén, P., P.E. Colucci, and P.J. Van Soest. 1980. Investigation of chromium, cerium and cobalt as markers in digesta. Rate of passage studies. Journal of the Science of Food and Agriculture 31:625-632.
- Ullrey, D.E. 1986. Nutrition of primates in captivity. In: Primates: The Road to Self-Sustaining Populations (K. Benirschke, ed.) Pp. 823-835. Springer-Verlag, New York, NY.
- Ullrey, D.E. 1987. If the diet is deficient will the animal let us know? Proceedings of the Seventh Annual Dr. Scholl Conference on the Nutrition of Captive Wild Animals.
- Ullrey, D.E., J.T. Nellist, J.P. Duvendeck, E.A. Whetter and L.D. Fay. 1987. Digestibility of vegetative rye for white-tailed deer. Journal of Wildlife Management 51:51-53.
- Ullrey, D.E., P.T. Robinson, T.A. Reichard, P.A. Whetter, D.K. Brockman. 1982. Dietary studies with colobids at the San Diego Zoo. Unpublished Research Report. Michigan State University, East Lansing, MI.
- Urbain, A., J. Nouvel, P. Bullier, and J. Rinjard. 1955. Rapport sur la mortalité et la natalité enregistrées au Parc Zoologique du Bois de Vincennes pendant l'année 1954. Bull. Mus. Hist. Nat., Paris, 2nd ser., 27(2):117-134.
- Van Dyne, G.M. 1968. Measuring quantity and quality of the diet of large herbivores. In: Productivity of Large Herbivores. Pp. 54-94. Blackwell. Oxford, England.
- Van Soest, P.J. 1967. Development of a comprehensive system of feed analysis and its application to forages. Journal of Animal Science 26:119-128.
- Van Soest, P.J. 1977. Plant fiber and its role in herbivore nutrition. Cornell Veterinarian 67:307-326.

Van Soest, P.J. 1982. Nutritional Ecology of the Ruminant. Cornell University Press. Ithaca, NY.

Van Soest, P.J. 1994. Nutritional Ecology of the Ruminant. Cornell University Press. Ithaca, NY

Van Soest, P.J. 1995. Personal Communication.

Van Soest, P.J., P. Uden, and K.F. Wrick. 1983. Critique and evaluation of markers for use in humans and farm and laboratory animals. Nutrition Reports International 27:17-28.

Van Soest, P.J., and R.H. Wine. 1967. Use of detergents in the analysis of fibrous feeds. IV. Determination of plant cell wall constituents. Journal of Association of Official Agricultural Chemists 50:50.

Warnell, K.J. 1988. Feed intake, digestibility, digesta passage, and fecal microbial ecology of the red pande (Ailurus fulgens). M.S. Thesis. Michigan State University, MI.

Warnell, K.J., S.D. Crissey, and O.T. Oftedal. 1989. Utilization of bamboo and other fiber sources in red panda diets. In: Red Panda Biology (A.R. Glatson, ed). Pp. 51-56. SPB Academic Publishing. The Hague, Netherlands.

Watkins, B.E., D.E. Ullrey, and P.A. Whetter. 1985. Digestibility of a high-fiber biscuit based diet by black and white colobus (*Colobus guereza*). American Journal of Primatology 9:137-144.

Whetter, P.A., and D.E. Ullrey. 1978. Improved fluorometric method for determining selenium. Journal of Association of Official Analytical Chemists 61:927-930.

White, F.J. 1991. Social organization, feeding ecology, and reproductive strategy of ruffed lemurs, *Varecia variegata*. In: Primatology Today: Proceedings of the 13th Congress International Primatology Society. (A. Ehara, T. Kuimura, O.Takenaka, and M Iwamoto, eds.). Pp. 81-84. Elsevier, England.

Yeager, C.P. 1989. Feeding ecology of the proboscis monkey (Nasalis larvatus) International Journal of Primatology 10:497-530.




Table Al. Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta seniculus) in the central llanos region of Venezuela.

Family/ Species	Species	Part*	Season	п	Dry matter (%)	ter (%)
	no.				Mean	SD
Anacardiaceae						
Mangifera indica	(1)	YL	Wet	1	20.6	
Mangifera indica	(2)	UF	Dry	4	22.5	0.00
Mangifera indica	(3)	UF	Wet	1	19.0	
Mangifera indica	(4)	RF	Dry	8	22.1	2.09
Mangifera indica	(5)	FUL	Dry	2	31.8	0.48
Annonaceae						
Annona jahnii	(9)	T	Wet	1	28.3	
Annona jahnii	(7)	UE	Wet	-1	30.2	1
Annona jahnii	(8)	RF	Wet	1	27.2	,
Annona jahnii	(6)	FL	Wet	-1	19.1	
Annona punicifolia	(10)	UF	Wet	1	16.0	
Annona punicifolia	(11)	RF	Wet	8	36.8	25.90
Arecaceae						
Copernicia tectorum	(12)	(St.)	Wet	2	28.2	2.90
. Copernicia tectorum	(13)	UF	Dry	3	33.1	3.45
Copernicia tectorum	(14)	UF	Wet	1	Q	1
Copernicia tectorum	(15)	RF	Wet	1	27.8	1

211 Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta 0.05 0.07 0.12 0.16 0.33 0.05 ı ı ı GE (kcal/g) 4.58 4.05 4.02 4.24 4.06 5.03 5.35 5.06 5.22 4.80 5.15 4.53 4.58 4.61 Mean 0.07 0.62 0.18 0.28 0.29 0.08 SD ı ı ı Ash (%) 4.4 5.9 2.5 2.8 11.0 4.5 4.0 5.9 3.4 3.2 8.0 8.0 8.5 Mean 0.14 0.22 0.31 1.00 3.36 0.35 ı SD (8) seniculus) in the central llanos region of Venezuela. 1.5 1.6 2.5 3.2 6.9 3.6 13.3 9.0 3.1 10.7 12.5 0.5 Mean 0.45 0.12 1.05 80.0 0.21 0.01 ı ı 1 SD CP (8) 2.5 2.5 4.5 5.6 17.5 10.8 7.6 6.7 7.2 6.0 14.9 10.1 14.7 Mean Table Al (cont'd). (Species no.) Anacardiaceae Annonaceae (13) (10) (11) Arecaceae (12)(14) (4) (2) (9) (8) 6 (1) (2) (3) Family

4.33

6.3

0.7

5.0

(15)

Table A1 (cont'd). Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta seniculus) in the central llanos region of Venezuela.

Family	NDF	(%)	ADF (%)	(8)	ADL (%)	(%)	Cellulose)Se (%)
(species no.)	Mean	SD	Mean	SD	Mean	SD	Mean	
Anacardiaceae								de
(1)	37.0	-	14.1	1	3.3		10.7	
(2)	55.7	2.09	8.0	0.26	1.5	0.18	6.2	0 33
(3)	15.8	1	9.3	,	4.5	-	5.0	
(4)	15.9	4.01	13.5	4.24	7.9	2.82	5.8	1.06
(5)	38.3	1.80	29.0	1.16	9.5	09.0	19.3	1.23
Annonaceae								24:1
(9)	57.5	-	37.4	-	13.9	-	22.7	
(7)	67.9	-	46.9	1	30		16.7	
(8)	59.0	-	41.5	1	15.8		25.7	
(6)	57.5	-	45.8	-	24.7	-	20.8	
(10)	8.69	-	52.1	,	23.5		28.2	,
(11)	52.1	1.40	39.4	2.14	19.5	4.11	21.4	5.14
Arecacaea								
(12)	70.2	12.49	50.8	17.33	32.8	14.41	17.9	2 97
(13)	80.8	1.93	45.9	1.42	23.2	1.11	22.6	0.31
(14)	68.3	-	33.2	-	16.6	,	16.5	
(15)	55.7	,	35.5	1	20.4		14 0	

Wet Wet Dry

(16)(11) (18)

Marsdenia undulata Marsdenia undulata Ceiba pentandra

Bombacaceae Cactaceae

FLBU UF н

Table A2. Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta seniculus)

in the central llanos region of Venezuela.

Family/ Species Asclepidaceae

u

Season

Part*

Species

no.

1	1	1 1	-	1 1	1 1
(%)	SD	21.84	-		
Dry matter (%)		Н		Н	Ц
Dry m	Mean	33.4	10.0	20.2	25.3

2	21	.3
	1	
		ľ
	1	
	1	_

Pereskia guamacho	(19)	RF	Wet	1	25.3	1
Chrysobalanaceae						
Licania pyrifolia	(20)	FL/BU	Wet	1	30.6	
Licania pyrifolia	(21)	NL	Dry	1	23.9	,
Licania pyrifolia	(22)	ī	Dry	2	40.8	0.50
Cucurbitaceae						
Melothria trilobata	(23)	I	Wet	. 2	17.2	0.08
Dilleniaceae						
Tetracera volubris	(24)	FL	Wet	1	22.8	,
Euphorbiaceae						
Sapium aucuparium	(25)	ı	Wet	е	67.0	2.41

Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta 0.11 0.07 0.54 0.04 1 GE (kcal/g) 4.30 4.36 4.96 4.39 4.48 4.70 4.31 4.61 4.99 4.21 Mean 0.10 0.43 0.11 0.29 SD ı 1 ı Ash (%) 14.0 9.5 3.3 8.2 4.1 3.6 5.8 3.9 15.3 8.5 Mean 2.78 0.40 0.20 0.23 SD EE (8) seniculus) in the central llanos region of Venezuela. 3.5 6.3 5.0 2.4 2.5 5.2 5.7 6.2 Mean 0.08 0.39 6.95 2.00 SD CP (%) 9.6 19.2 6.6 15.6 13.6 15.1 12.6 5.7 13.0 9.1 Mean rable A2 (cont'd). Chrysobalanaceae Euphorbiaceae Cucurbitaceae Asclepidaceae (Species no.) Dilleniaceae (24) (22) Bombaceae (18) Cactaceae (13) (20) (21) (22) (23) (16) Family.

Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta RF=ripe fruit; FL=flower; PET=petiole; 1.85 1.06 0.50 0.90 Cellulose (%) SD 1 20.2 14.7 20.0 23.8 20.4 19.0 24.1 20.5 20.4 16.0 Mean 3.41 1.74 4.00 0.70 SD (%) ADL 12.9 6.2 30.3 12.3 14.5 8.6 40.1 33.7 40.1 Mean L=mature leaf; NL=new leaf; YL=young leaf; F=fruit; UF=unripe fruit; 5.49 0.76 2.48 5.30 SD ADF seniculus) in the central llanos region of Venezuela. 33.3 20.3 50.4 30.1 35.1 58.7 53.3 49.5 38.5 Mean 4.70 4.81 0.60 4.84 ı SD NDF (%) 74.0 50.1 28.3 64.6 54.7 6.79 43.0 62.5 36.8 Mean Table A2 (cont'd). Chrysobalanaceae Cucurbitaceae Euphorbiaceae (Species no.) Asclepidaceae Dilleniaceae Bombacaceae PUL=pulvinus (16) (11) (18) Cactaceae (19) (20) (22) (23) (24) (22) Family

		-				
Family/ Species	Species	Part*	Season	п	Dry matter (%)	ter (%)
					Mean	SD
Fabaceae (Caesalpinoideae)						
Cassia grandis	(26)	1	Wet	8	51.6	12.13
Cassia grandis	(27)	UF	Wet	6	10.3	1.63
Cassia moschata	(28)	1	Wet	s	45.8	4.06
Cassia moschata	(29)	T	Dry	1	48.3	
Cassia moschata	(30)	ХГ	Wet	7	32.9	
Cassia moschata	(31)	YL	Dry	2	36.7	2.88
Cassia moschata	(32)	UF	Wet	3	20.3	1.49

1	
0	

Wet Wet Dry Wet Wet Wet Wet Wet Wet

н н ы

(33) (34) (32) (36) (31) (38) (38) (40) (41)

Albizia columbiana Albizia guachapele Albizia guachapele

Fabaceae (Mimosoldeae)

35.4 37.1 28.7 31.0 45.3 22.2 38.3

216	
6	1

1.98

.

1 7

Enterolobium cyclocarpum Pithecellobium dualense Pithecellobium dualense Pithecellobium saman

Albizia guachapele Albizia guachapele

XI ž 1

0.74 2.21

FL (BU)

ы

217 Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta 0.11 0.26 0.24 0.03 0.29 0.04 0.24 0.25 , ı 1 1 GE (kcal/g) 5.19 4.62 5.17 4.82 5.39 5.03 4.76 5.69 5.06 5.10 5.17 5.49 4.89 Mean ı 0.99 0.29 0.24 0.18 1.08 0.25 0.11 0.32 ı ı SD (%) Ash 3.0 3.5 3.6 4.0 4.5 4.3 4.2 4.4 6.3 8.3 5.2 6.8 3.7 Mean 0.53 2.67 1.57 0.12 0.14 0.25 0.40 0.32 ı ı i 1 1 (8) seniculus) in the central llanos region of Venezuela. EE 8.3 4.4 5.4 1.9 2.7 3.2 1.5 1.2 4.7 0.8 5.3 5.1 ı Mean 1.66 1.81 2.76 0.26 1.63 0.81 1.62 ı ı ı 1 SD CP (%) 26.4 22.3 25.3 33.0 12.8 7.1 18.8 12.5 9.01 12.1 14.8 16.6 Mean 1 Table A3 (cont'd).

Species no.)

Family

(27) (28) (53) (30) (31) (32) (33)

(26)

Fabaceae

0.11

5.61

0.44

2.38

7.0

2.38

(41)

0.17

5.19

0.09

5.4 3.1

0.94

1.4

27.7 23.2

0.80 0.08

16.9

(31) (38) (38) (40)

(34) (32) (38)

Fabaceae

22.6

4.88

15.4 33.2

7.48

38.6 50.0

16.42

46.2

Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta 0.94 1.87 1.26 1.59 Cellulose (%) SD 17.6 9.1 17.3 22.1 18.5 13.3 Mean 3.69 3.57 10.43 SD ADL (%) 22.0 20.2 19.9 21.1 29.4 19.2 Mean 66.7 3.51 3.31 8.64 SD ADF (%) seniculus) in the central llanos region of Venezuela. 40.9 38.4 39.3 34.0 45.0 46.7 Mean 5.13 4.13 1.90 6.43 SD (%) NDF 50.5 21.8 56.3 53.6 65.0 54.1 Mean

Table A3 (cont'd).

Species no.)

Family

(56) (27) (28) (58) 30) (31) (32) (33)

Fabaceae

-	
15.0	

3.03

15.9

1.19

17.5 15.3

1.83

33.6 33.4

8.40

(34) (32) (31) (38)

Fabaceae

(38)

70.9 55.7 55.4 72.3 49.7

ı

18.2 19.2 11.5 18.6 9.5 18.8

0.83 1.17 0.31 6.36

1.70 4.68 3.70

1.59 5.59 0.52 3.85

10.56 3.95 2.81

20.8 11.3 31.5

40.2 23.5 ı

0.49

7.4 22.3

49.9 16.8 41.6

74.6 44.2 59.4

(38) (40)

ı

L-mature leaf; NL-new leaf; YL-young leaf; F=fruit; UF=unripe fruit; RF=ripe fruit; FL=flower; PET=petiole;

4.54

(41)

Table A4. Proximate and fiber composition (dry basis) in the central llanos region of Venezuela.		nts consu	med by re	d howl	ers (Alouat	of plants consumed by red howlers (Alouatta seniculus)
Family/ Species	Species	Part	Season	R	Dry matter	tter (%)
	: OII				Mean	SD
Fabaceae (Mimosoideae)						
Pithecellobium saman	(42)	L	Dry	2	36.9	1.07
Pithecellobium saman	(43)	NT	Dry	1	Ą	
Pithecellobium saman	(44)	YL	Wet	3	24.5	
Pithecellobium saman	(45)	FLBU	Dry	1	£	• •
Pithecellobium saman	(46)	FL	Dry	2	19.0	1 78
Pithecellobium saman	(47)	RF	Dry	-	44.0	• •
Pithecellobium saman	(48)	UF	Drv	2	22.5	
Pithecellobium tortum	(49)	1	Wet	, v	28.0	
Pithecellobium tortum	(20)	YL	Wet	-	22.2	0/.5
Pithecellobium tortum	(51)	FL	Wet	-	18.8	
Pithecellobium tortum	(52)	LBU, NL	Dry	2	41.8	110
Pithecellobium tortum	(53)	UF	Dry		71.4	
Fabaceae (Papilionoideae)						
Entada polystachya	(54)	1	Wet	1	43.7	
Machaerium moritzianum	(55)	FL	Dry		₽ Q	
Platymiscium diadelphum	(56)	ı	Wet	-	37.5	
Platymiscium diadelphum	(57)	YL, NL	Wet	1	31.2	-

220 90.0 0.12 0.01 0.15 0.18 0.03 1 ı ı GE (kcal/g) 5.51 5.78 5.53 5.89 4.74 4.61 4.82 5.29 5.19 5.20 4.68 5.30 4.82 5.07 5.11 Mean Q 0.03 1.57 0.08 1.05 0.32 0.37 ı ì ı SD (%) Ash 4.0 4.9 3.3 3.6 6.2 5.0 3.5 3.2 3.9 5.6 5.5 3.3 5.2 Mean Q ND 9 1.88 0.15 0.23 0.43 0.18 0.29 1 ı SD seniculus) in the central llanos region of Venezuela. 1.4 3.9 1.6 5.2 5.3 1.8 1.5 2.8 1.8 1.8 1.5 5.2 2.2 2.1 Mean 3.73 2.15 0.19 0.52 0.41 0.85 ı SD 16.6 22.8 19.3 17.3 28.2 35.7 32.8 23.2 17.6 22.8 16.4 36.1 29.3 Mean Q g ND (Species no.) (23) (54) (22) (99) (57) (21) (52)(44) (42) (46) (41) (48) (48) Fabaceae (42)(43) Fabaceae Family.

Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta

rable A4 (cont'd).

Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta 2.26 9.70 0.87 3.38 0.88 SD ı Cellulose (%) 14.7 19.2 14.8 10.8 13.0 15.6 21.0 17.6 17.4 13.3 13.0 17.4 17.7 30.4 20.4 11.1 Mean 1.84 0.40 0.45 1.70 1.48 0.99 SD ADL (%) 12.6 17.6 8.5 9.3 23.9 15.8 8.0 14.5 18.7 19.7 40.0 40.8 15.2 10.8 14.6 18.2 Mean 3.84 4.14 2.45 3.65 0.64 1.42 ı SD ADF (%) seniculus) in the central llanos region of Venezuela. 23.6 45.9 33.2 39.7 34.9 31.4 20.7 55.6 30.2 59.6 34.4 24.9 35.8 31.3 19.5 41.9 Mean 6.26 6.28 6.16 3.85 1.22 1.05 ı ı ı SD 61.9 57.9 43.7 44.6 48.0 64.4 49.5 48.5 44.6 55.3 48.2 68.2 54.1 36.6 61.1 57.3 Mean rable A4 (cont'd). (Species no.) (96) (57) (55)(54) (51) (52)(53)(49)(20) (47) (48) (44) (45)(46) (42)(43)Fabaceae Fabaceae Family

* L-mature leaf; NL-new leaf; YL-young leaf; F-fruit; UF-unripe fruit; RF-ripe fruit; FL-flower; PET-petiole; PUL=pulvinus

Table A5. Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta seniculus) in the central llanos region of Venezuela.

family/ Species	Species	Part	Season	п	Dry matter (%)	cer (%)	
	по.				Mean	SD	
Fabaceae (Papilionoideae) (Cont'd)							
Platymiscium polystachyum	(58)	YL	Wet	1	21.2		
Platymiscium polystachyum	(69)	FLBU	Wet	1	17.9	,	
Pterocarpus acapulcensis	(09)	ı	Wet	4	33.3	8.49	
Pterocarpus acapulcensis	(61)	NL	Wet	4	27.6	1.77	
Moraceae							
Ficus pertusa	(62)	ı	Dry	3	26.5	16.32	
Ficus pertusa	(63)	I	Wet	4	29.6	3.17	
Ficus pertusa	(64)	UF	Wet	2	23.1	0.28	_
Ficus pertusa	(65)	RF	Wet	3	22.8	1.09	
Ficus trigonata	(99)	UF	Wet	1	16.3	-	
Poylgonaceae							
Coccoloba sp.	(67)	YL	1	'	-		
Rhamnaceae							
Zizyphys saeri	(89)	ı	Wet	1	43.5	-	
Zizyphys saeri	(69)	I	Dry	1	35.0	-	
Zizyphys saeri	(70)	[a ₄	Wet	1	39.8	1	

Table A5 (cont'd). Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta seniculus) in the central llanos region of Venezuela.

									2	23								
1/4)	an a	3		,	0.23	0.14		0.03	0.15	0.04	0.02							
GE (kcal/q)	Mean		5.31	5.13	5.09	5.13		4.29	4.23	4.89	4.50	4.43		-		4.51	4.59	4.59
(8)	SD		-	-	0.05	0.23		0.20	1.43	0.17	0.36	,				-	-	-
Ash (%)	Mean		6.9	7.4	4.8	4.3		9.6	11.4	6.9	4.9	5.9		-		10.6	8.6	6.1
(8)	SD		-	-	0.37	0.04		0.85	0.45	0.41	1.39	,		-		-		,
EE (Mean		2.7	2.5	1.5	1.2		2.7	3.9	2.0	3.3	1.8		,		2.2	1.6	0.8
(8	SD		'	-	0.61	5.23		5.02	1.47	0.17	0.26	-		,		-	-	-
CP (%)	Mean		25.0	25.5	19.1	24.9		11.2	12.3	7.1	4.9	10.5		-		14.1	22.6	5.9
Family	(Species no.)	Fabaceae	(58)	(59)	(60)	(61)	Moraceae	(62)	(63)	(64)	(65)	(99)	Polygonaceae	(67)	Rhamnaceae	(89)	(69)	(10)

224 Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta J-mature leaf; NL-new leaf; YL-young leaf; F-fruit; UF-unripe fruit; RF-ripe fruit; FL-flower; PET-petiole; 0.67 2.12 3.85 3.19 1.38 1.02 ı Cellulose (%) ı 11.3 15.4 19.0 17.3 21.2 21.3 25.8 17.9 8.02 18.6 13.3 21.7 Mean 1.15 1.72 3.54 3.05 1.87 1 ı SD ADL (%) 9.92 6.6 13.3 20.3 12.7 9.6 16.3 6.5 14.7 12.1 27.3 22.8 Mean 4.08 3.69 6.38 3.08 2.60 1 SD ADF (8) seniculus) in the central llanos region of Venezuela. 38.3 34.2 30.2 30.3 42.4 36.1 23.6 39.0 37.4 53.2 23.4 24.5 Mean 2.71 2.47 5.64 2.46 4.70 0.42 SD NDF (%) 58.1 61.7 38.8 54.6 55.4 48.0 71.6 47.5 60.7 57.0 43.3 Mean rable A5 (cont'd). (Species no.) Polygonaceae Rhamnaceae (64) (65) (99) (67) (89) (69) (10) (63) (88) (88) (09) (61) (62) Moraceae Fabaceae Family

PUL-pulvinus

Table A6.

Family/ Species	Species	Part	Season	E	Dry matter (%)	ter (%)	
	no.				Mean	SD	
Rubiaceae							
Randia venezuelensis	(11)	YL	Wet	2	18.0	3.79	
Randia Venezuelensis	(72)	NL	Wet	1	23.0	1	
Randia venezuelensis	(73)	UF	Wet	1	19.9	-	
Unidentified #01	(74)	I	Wet	3	29.9	6.84	
Unidentified #01	(75)	YL	Wet	1	18.7	1	
Rutaceae							
Zanthoxylum culantrillo	(16)	ı	Wet	1	74.9	1	22
Zanthoxylum culantrillo	(77)	YL, L	Wet	2	33.4	0.55	5
Zanthoxylum culantrillo	(78)	YL	Wet	2	31.2	4.06	
Zanthoxylum culantrillo	(19)	FLBU	Wet	2	31.7	1	

	ı	
9	ı	

2.00

41.3 18.6 25.0 58.3 35.2

4

Wet

н

(80) (81) 11.48

Dry

(84) (82) (83)

Dry Wet Dry

> UF RF

Guazuma ulmifolia Guazuma ulmifolia Guazuma ulmifolia Guazuma ulmifolia

Paullinia cururu

Sterculiaceae Sapindaceae

40.60

4.34 4.63

2.13 0.77

66.0

2.13 0.91

(83) (84)

13.2

Table A6 (

Family	CP ((%)	EE ((8)	Ash (%)	(8)	GE (kcal/g)	a1/g)
(Species no.)	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Rubiaceae								
(71)	20.8	0.01	2.8	0.46	4.5	0.10	4.87	0.10
(72)	QN		2.1	1	QN	,	4.75	
(73)	14.2		1.3	1	4.6	,	4.54	
(74)	16.6	1.56	4.1	2.57	7.5	06.0	4.61	0.07
(75)	QN	-	3.1	1	QN	-	4.99	-
Rutaceae								
(76)	12.5	1	1.3	,	5.9	-	4.78	
(77)	12.6	0.07	2.7	0.21	5.1	0.12	4.95	0.25
(78)	14.1	0.76	1.9	1.01	4.1	0.13	4.81	0.11
(46)	15.8	0.42	2.1	0.12	5.8	0.81	5.01	0.20
Sapindaceae								
(80)	12.5	1.20	3.4	0.81	7.8	0.97	4.49	0.09
Sterculiaceae								
(81)	12.1	2.60	1.1	0.12	7.4	1.68	4.10	0.10
(82)	8.8	,	2.5	1	5.8	1	4.32	,

Family	NDF	(8)	ADF	(8)	ADL	(8)	Cellulose	se (%)
(Species no.)	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Rubiaceae								
(71)	42.8	6.43	28.4	1.09	10.4	1.43	17.3	0.77
(72)	44.4	-	26.4		9.7	-	16.7	,
(73)	45.9	,	35.5	1	19.4	,	15.3	,
(74)	60.3	2.24	52.2	2.52	23.2	1.56	27.6	1.50
(75)	59.7	-	48.5	-	22.2	1	25.7	-
Rutaceae								
(16)	QN	1	55.5	,	40.0	-	15.7	
(77)	48.0	2.95	41.5	7.37	25.5	3.64	16.3	1.31
(78)	44.5	3.33	34.6	2.15	14.2	2.80	19.6	4.80
(42)	48.0	5.24	36.9	5.47	21.4	5.12	15.2	0.61
Sapindaceae								
(80)	59.1	5.17	43.9	1.89	19.7	2.93	22.4	3.67
Sterculiaceae								
(81)	37.6	20.84	36.6	4.94	19.7	6.70	15.4	0.31
(82)	67.9	1	48.6	1	17.8	1	30.2	

29.0 20.1

15.8 22.2

18.80 6.74

43.4 45.1

5.67 16.93

55.5 59.0

(83) (84)

6.28 11.16

Table A7. Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta seniculus)

in the central llanos region of Venezuela.	•		^- F		on protect of the	(STINGTER SETTINGT OF THE SETT
Family/ Species	Species	Part*	Season	u	Dry matter (8)	ter (%)
	no.				Mean	SD
Sterculiaceae (Cont'd)						
Guazuma ulmifolia	(85)	FL	Wet	1	19.0	•
Sterculia apetala	(86)	FL	Dry	1	QN.	ı
Sterculia apetala	(87)	FLBU	Dry	2	20.7	4.48
Sterculia apetala	(88)	FL (BU)	Dry	1	21.3	•
Sterculia apetala	(88)	ST	Dry	1	Q.	ŧ
UNIDENTIFIED PLANTS						
Unidentified #02	(06)	ı	Dry	2	34.0	0.95
Unidentified #03	(91)	L	Dry	1	21.4	•

229 Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta 0.04 90.0 í 1 ı GE (kcal/g) 4.57 4.25 4.55 4.43 4.98 Mean NF ND 0.15 0.52 ı ı ı SD Ash (%) 6.5 8.1 6.9 7.2 4.7 Mean ND N 0.13 0.31 Ether Extract (%) ı ı SD ı seniculus) in the central llanos region of Venezuela. 1.6 0.8 3.1 2.4 2.2 Mean R N 0.71 0.21 Crude Protein (%) , SD 1 9.3 11.7 18.1 15.1 20.1 Mean ND Q Table A7 (cont'd). Family/ Sample Sterculiaceae UNIDENTIFIED (06) (16) (88) (88) (88) (88) (81)

Table A7 (cont'd). Proximate and fiber composition (dry basis) of plants consumed by red howlers (Alouatta seniculus) in the central llanos region of Venezuela.

Family/Sample	NDF (%)	(8)	ADF (%)	(%)	TON	ADL (%)	Cellulose (%)	se (%)
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Sterculiaceae								
(85)	57.8	,	44	-	16.6	-	26.4	'
(86)	QN	-	QN	1	ND	,	ND	,
(87)	47.7	0.49	36.8	2.82	13.6	1.68	22.5	1.66
(88)	70.7	'	51.9	-	21.1	,	30.4	,
(88)	QN	-	QN	-	QN	-	N	
UNIDENTIFIED								
(06)	55.3	1.86	31.8	3.68	14.1	0.62	13.2	8.26
(61)	46.8	1	45.8	1	23.7	1	19.1	1

Family/ Species	Species	Part*	Season	п	Dry matter (%)	ter (%)
	no.				Mean	SD
Anacardiaceae						
Mangifera indica	(1)	YL	Wet	1	20.6	
Mangifera indica	(2)	UF	Dry	4	22.5	0.00
Mangifera indica	(3)	UF	Wet	1	19.0	,
Mangifera indica	(4)	RF	Dry	3	22.1	2.09
Mangifera indica	(5)	PUL	Dry	2	31.8	0.40

23	1	
-		,
28.3	30.2	27.2

30.2 27.2 19.1 16.0 36.8

Wet

(9)

Wet Wet Wet Wet Wet Wet Dry Wet

UF RF FL UF RF

(8) (6) (7)

(10) (11) (12)(13)

Annona punicifolia Annona punicifolia

Arecaceae

Annona jahnii Annona jahnii Annona jahnii Annona jahnii 2.90 3.45

28.5

2 3

33.1 g 27.8

Wet

RF OF UF

(12)

(14)

Copernicia tectorum Copernicia tectorum Copernicia tectorum

Copernicia tectorum

25.90

2

Macro mineral composition (dry basis) of plants consumed by red howlers (Alouatta 0.03 0.01 90.0 0.02 0.13 0.21 SD , ı (8) Mg 0.27 60.0 0.04 0.24 0.25 0.29 0.20 0.17 0.38 0.26 0.20 0.15 0.42 0.29 0.23 Mean 0.37 1.36 0.47 90.0 0.47 0.25 SD ı ı 1 (8) 1.46 1.56 3.06 2.39 × 2.10 1.49 1.21 2.75 1.67 1.72 2.01 1.30 1.38 2.28 3.11 Mean 9.48 2.40 31.79 10.68 51.09 30.26 SD ı (mdd) 184 94 53 122 146 118 117 39 52 29 48 30 53 52 87 Na Mean 2.36 0.50 1.59 0.76 2.42 2.44 0.61 2.00 3.00 1.83 16.52 11.33 0.36 1.40 2.00 Ca: P seniculus) in the central llanos region of Venezuela. 0.03 0.04 0.02 0.00 0.01 00.0 SD 1 1 0.12 0.11 60.0 0.33 90.0 0.21 0.18 0.14 0.22 0.29 0.15 0.21 0.12 Д 90.0 0.02 Mean 0.03 0.04 0.02 0.03 0.01 0.27 1 ı (%) 0.16 0.29 0.24 0.26 0.22 3.47 0.11 0.46 0.21 90.0 0.11 2.04 0.05 0.20 0.12 Mean (cont'd). Anacardiaceae no.) Annonaceae (11) (12)(13) (14)(15)(10) Arecaeae Species (2) (9) (7) (8) 6) (3) (4) (2) rable A8 Family

Wet Wet Dry

(16)(11) (18)

Marsdenia undulata Marsdenia undulata

Asclepidaceae

Ceiba pentandra

Bombacaeae Cactaceae Chrysobalanaceae

UF н

FLBU

Season

Part*

Species

the central llanos region of Venezuela.

Table A9.

Family/ Species

1 1	1.1.1	11	1 1
SD SD	21.84		-
Dry matter (%) Mean SD	33.4	20.2	25.3

	2	3	33	3
		1		
				1
п		1	_	

Pereskia guamacho	(19)	RF	Wet	1	25.3	1
sobalanaceae						
Licania pyrifolia	(20)	FL (BU)	Wet	1	30.6	1
Licania pyrifolia	(21)	NE	Dry	1	23.9	1
Licania pyrifolia	(22)	1	Dry	2	40.8	0.44
urbitaceae						
Melothria trilobata	(23)	П	Wet	2	17.2	0.08
leniaceae						
Tetracera volubris	(24)	FL	Wet	1	22.8	-
horbiaceae						
Sapium aucuparium	(25)	п	Wet	т	67.0	2.41

Cucurbitaceae

Euphorbiaceae Dilleniaceae

Family	Ca ((%)	P ((%)	Ca:P	Na ((mdd)	K ((8)	Mg ((8)
(Species no.)	Mean	SD	Mean	SD		Mean	SD	Mean	SD	Mean	SD
Asclepidaceae											
(16)	0.47	0.43	0.16	0.05	2.94	96	80.08	3.13	1.53	0.50	0.26
(17)	0.11	-	0.08	,	1.38	89	1	1.66	ı	0.18	1
Bombacaceae											
(18)	1.05	1	0.33	1	3.18	82	-	3.05	1	0.39	1
Cactaceae											
(19)	0.17	1	0.21	1	0.81	57	-	1.39	-	0.29	-
Chrysobalanaceae											
(20)	0.17	1	0.18	,	0.94	93	1	1.37	1	0.19	1
(21)	0.17	ı	0.37	,	0.46	154	1	2.57	-	0.36	-
(22)	0.31	0.05	0.15	00.00	2.07	118	9.62	1.38	90.0	0.28	0.01
Cucurbitaceae											
(23)	2.95	0.79	0.29	90.0	10.17	96	8.28	2.63	1.40	0.45	0.14
Dilleniaceae											
(24)	0.97	1	0.31	1	3.13	98	1	1.99	1	0.27	1
Euphorbiaceae											
(25)	2.17	0 13	0.15	10 0	14 47	545	25.02	1 46	0.16	1.13	0.07

Table AlO. Macro mineral composition (dry basis) c the central llanos region of Venezuela.	of plants	plants consumed by	by red ho	red howlers	(Alouatta seniculus)	eniculus) in
Family/ Species	Species	Part.	Season	u	Dry matter	:ter (%)
	no.				Mean	SD
Fabaceae (Caesalpinoideae)						
Cassia grandis	(26)	L	Wet	3	51.6	12.13
Cassia grandis	(27)	UF	Wet	3	10.3	1.63
Cassia moschata	(28)	L	Wet	5	45.8	4.06
Cassia moschata	(53)	L	Dry	1	48.3	ı
Cassia moschata	(30)	YL	Wet	1	32.9	1
Cassia moschata	(31)	YL	Dry	2	36.7	2.88
Cassia moschata	(32)	UF	Wet	3	20.3	1.49
Fabaceae (Mimosoideae)						
Albizia columbiana	(33)	L	Wet	1	81.0	3
Albizia guachapele	(34)	Ţ	Wet	2	35.4	1
Albizia guachapele	(35)	T	Dry	1	37.1	ı
Albizia guachapele	(36)	YL	Wet	1	28.7	1
Albizia guachapele	(37)	NL	Wet	3	31.0	1.98
Enterolobium cyclocarpum	(38)	ē	Wet	1	1	1
Pithecellobium dualense	(38)	1	Wet	5	45.3	2.21
Pithecellobium dualense	(40)	FL/BU	Wet	2	22.2	0.74
Pithecellobium saman	(41)	Ţ	Wet	7	38.3	7.44

Table A10 (cont'd). Macro mineral composition (dr. seniculus) in the central llanos region of Venezuela.	Macro m	ineral c os regio	composit;	ion (dry	basis)	of plan	Macro mineral composition (dry basis) of plants consumed by red howlers (Alouatta ral llanos region of Venezuela.	med by	red how]	lers (Al	ouatta	
Family	Ca (%)	(8)	Ъ ((8)	Ca:P	Na (Na (ppm)	K (%)	8)	Mg (%)	(8)	
(Species no.)	Mean	SD	Mean	SD		Mean	SD	Mean	SD	Mean	SD	
Fabaceae										1		
(26)	1.91	0.28	0.11	0.30	17.36	50	6.20	0.54	0.17	0.40	0.50	
(27)	0.26	0.02	0.29	0.01	06.0	53	4.93	2.62	1.36	0.28	0.03	
(28)	1.34	0.62	0.11	0.02	12.18	47	25.31	0.51	0.21	0.51	0.09	
(29)	2.21	1	0.13	1	17.00	48	,	0.56	,	0.38	,	
(30)	0.31	1	0.14	-	2.21	33	1	1.17	,	0.15	,	
(31)	0.18	0.04	0.24	0.03	0.75	80	25.37	1.30	0.07	0.48	0.61	
(32)	0.16	0.02	0.16	0.02	1.00	38	15.08	1.65	0.47	0.14	0.01	2
Fabaceae												36
(33)	1.01	1	0.14	-	7.21	160	-	0.74	-	0.21	-	
(34)	0.65	0.34	0.19	0.05	3.42	66	33.19	1.40	0.03	0.39	0.01	
(35)	0.31	1	0.22	-	1.41	114	-	1.18	-	0.25	-	
(36)	0.27	-	0.30	-	0.90	25	-	1.81	-	0.25	-	
(37)	0.16	0.03	0.42	0.05	0.38	45	3.32	1.69	0.07	0.26	0.01	
(38)	N		ND			ND		ND		ND		
(39)	0.99	0.09	0.11	0.03	9.00	42	11.38	0.61	0.04	0.61	1.12	
(40)	0.36	90.0	0.44	0.01	0.82	47	20.85	2.30	0.13	0.21	0.04	
(41)	09.0	0.31	0.14	0.04	4.29	161	65.35	0.83	0.28	0.24	0.31	
' 1-mature leaef; NI-new leaf; YL-young leaf; F-fruit; UF-unripe fruit; RF-ripe fruit; El-flower; PET-petiole; PUI-pulvinus	ew leaf;	YL=young	leaf; F	-fruit, U	F=unripe	fruit;	RF=ripe	fruit; FI	i=flower	; PET=pe	tiole;	

Table A-11. Macro mineral composition of plants cor in the central llanos region of Venezuela.	consumed by red howler monkeys	red howle		(Alouatta		seniculus seniculus)
Family/ Species	Species	Part*	Season	и	Dry matter	ter (%)
	.110.				Mean	SD
Fabaceae (Mimosoideae)						
Pithecellobium saman	(42)	Ţ	Dry	2	36.9	1.06
Pithecellobium saman	(43)	NL	Dry	1	QN	QN
Pithecellobium saman	(44)	YL	Wet	3	24.5	1.29
Pithecellobium saman	(45)	FLBU	Dry	1	QN	QN
Pithecellobium saman	(46)	FL	Dry	2	19	1.00
Pithecellobium saman	(47)	RF	Dry	1	44	•
Pithecellobium saman	(48)	UF	Dry	2	22.5	1
Pithecellobium tortum	(49)	L	Wet	5	38	3.77
Pithecellobium tortum	(50)	YL	Wet	1	22.2	•
Pithecellobium tortum	(51)	FL	Wet	1	18.8	l
Pithecellobium tortum	(52)	BU, NI	Dry	2	41.8	0.11
Pithecellobium tortum	(53)	UF	Dry	1	71.4	,
Fabaceae (Papilionoideae)						
Entada polystachya	(54)	Ţ	Wet	1	43.7	1
Machaerium moritzianum	(55)	FL	Dry	1	ND	J
Platymiscium diadelphum	(56)	Ţ	Wet	1	37.5	ı
Platymiscium diadelphum	(57)	YL, NL	Wet	1	31.2	ê

Table A-11 (cont'd). seniculus) in the ce	. Macro mineral central llanos r		l composition c region of Vene	of	ints	consumed by	red	howler monkeys		(Alouatta se	seniculus
Family/ Sample	Ca	(8)	P ((8)	Ca:P	Na ((wdd)	Ж ((8)	Mg	(8)
	Mean	SD	Mean	SD		Mean	SD	Mean	SD	Mean	SD
Fabaceae					. :						
(42)	0.47	0.03	0.22	0.01	2.14	72	24.93	0.93	0.11	0.19	0.02
(43)	0.16	•	0.61	ı	0.26	59	. 1	2.96	1	0.19	,
(44)	0.17	0.01	0.28	0.04	0.61	48	17.66	1.79	0.24	0.32	0.63
(45)	0.67		0.66	-	1.02	48	1	2.72	-	0.38	ı
(46)	0.19	0.03	0.31	0.02	0.61	90	10.72	1.82	0.11	0.17	0.03
(47)	0.27		0.15	_	1.80	36	1	1.21	_	0.12	ı
(48)	0.21	0.05	0.20	0.02	1.05	41	10.37	1.99	0.15	0.15	0.05
(49)	0.79	0.23	0.13	0.03	6.08	68	7.54	1.37	0.56	0.30	0.05
(20)	0.48	1	0.33	1	1.45	26	1	2.13	1	0.22	ı
(51)	0.33	1	0.38	1	0.87	48	1	2.24	ı	90.0	,
(52)	0.48	0.18	0.35	0.09	1.37	34	17.53	1.35	0.13	0.25	0.01
(53)	0.34	-	0.14	•	2.43	40	-	1.27		0.18	1
Fabaceae											
(54)	0.98	_	0.08	-	12.25	27	1	0.70	_	0.15	1
(55)	1.28	1	0.27	ı	4.74	35	ł	2.08	1	0.10	,
(56)	0.48	•	0.26	1	1.85	112	ı	2.44	1	0.10	1
(57)	0.29	1	0.39	ı	0.74	42	ı	2.33	1	0.16	,
* L=mature leaf; NL=new	new leaf;	YL≖young	leaf;	F=fruit;	UF=unripe	e fruit;	RF=ripe	fruit;	FL=flower;		PET=petiole;

Se	
seniculus	
(Alouatta	
monkeys (
d howler	
red	
by	
sition of plants consumed by red f Venezuela.	
plants	
ion of	
of V	
compo qion	
acro mineral compos	
acro r	
. Ma	
le Al2. Mac	
Table A	
E- 44	44

eniculus)

Family/ Species	Species	Part	Season	п	Dry matter (%)	er (%)
	no.				Mean	SD
Fabaceae (Papilionoideae) (cont'd)						
Platymiscium polystachyum	(58)	YL	Wet	1	21.2	
Platymiscium polystachyum	(59)	FLBU	Wet	1	17.9	
Pterocarpus acapulcensis	(60)	I	Wet	4	33.3	8.49
Pterocarpus acapulcensis	(61)	NL	Wet	4	27.6	1.77
Moraceae						
Ficus pertusa	(62)	I	Dry	8	26.5	16.32
Ficus pertusa	(63)	I	Wet	4	29.6	3.17
Ficus pertusa	(64)	UF	Wet	2	23.1	0.28
Ficus pertusa	(65)	RF	Wet	е	22.8	1.09
Ficus trigonata	(99)	UF	Wet	1	16.3	,
Polygonaceae						
Coccoloba sp.	(67)	YL	-	,	1	1

239

35.0

Wet Dry Wet

(69)

Zizyphys saeri Zizyphys saeri Zizyphys saeri

Rhamnaceae

Macro mineral composition of plants consumed by red howler monkeys (Alouatta seniculus Table A12 (cont'd).

Family/ Sample	Ca ((8)	Ъ ((8)	Ca:P	Na (ppm)	(mdd	X	(%)	Mg	(%)
	Mean	SD	Mean	SD		Mean	SD	Mean	SD	Mean	SD
Fabaceae											
(58)	0.39	-	0.48	1	0.81	53	1	2.79	1	0.18	,
(59)	0.27	1	0.48	-	0.56	46	1	3.38	1	0.18	,
(09)	1.19	0.22	0.18	0.09	6.61	83	41.20	1.48	0.26	0.35	0.16
(61)	0.36	0.12	0.32	0.07	1.13	80	5.13	1.48	0.12	0.18	0.14
Moraceae											
(62)	4.45	3.69	0.13	0.04	34.23	310	33.46	0.93	0.35	08.0	0.11
(63)	2.89	0.16	0.11	0.02	26.27	226	141.1	1.67	0.42	0.50	0.21
(64)	1.14	0.16	0.16	0.01	7.12	146	30.39	1.96	0.26	1.52	2.12
(65)	0.74	0.03	0.12	0.02	6.17	108	19.61	1.52	0.24	0.31	0.01
(99)	0.74	1	0.23	1	3.22	377	1	1.95	1	0.27	
Polygonaceae											
(67)	1	1	ı	1	1	1	1	1	1	1	1
Rhamnaceae											
(89)	1.09	ı	0.22	1	4.95	42	,	2.55	1	0.43	1
(69)	1.69	1	0.15	1	11.27	114	1	2.71	1	0.38	,
(70)	1.17	1	0.15	1	7.80	55	1	1.34	1	0.12	1

11.50 40.60

58.3 35.2

7

Dry Dry

FL RF

(84) (83)

Guazuma ulmifolia Guazuma ulmifolia

Family/ Species	Species	Part*	Season	п	Dry matter	ter (%)	
	no.				Mean	SD	
Rubiaceae							
Randia venezuelensis	(71)	YL	Wet	2	18.0	3.79	
Randia venezuelensis	(72)	NL	Wet	1	23.0		
Randia venezuelensis	(73)	UF	Wet	1	19.9		
Unidentified #01	(74)	ī	Wet	е	29.9	6.84	
Unidentified #01	(75)	YL	Wet	1	18.7		
Rutaceae							
Zanthoxylum culantrillo	(16)	ч	Wet	1	74.9	,	24
Zanthoxylum culantrillo	(77)	YL, L	Wet	2	33.4	0.55	
Zanthoxylum culantrillo	(78)	YL	Wet	2	31.2	4.06	
Zanthoxylum culantrillo	(79)	FLBU	Wet	2	31.7		
Sapindaceae							
Paullinia cururu	(80)	ı	Wet	. 4	41.3	2.00	
Sterculiaceae							
Guazuma ulmifolia	(81)	Eu	Wet	2	18.6		
Guazuma ulmifolia	(82)	UE	Dry	1	25.0	1	

Table A13 (cont'd). Macro mineral composition of plants consumed by red howler monkeys (Alouatta seniculus seniculus) in the central llanos region of Venezuela

Family/ Sample	Ca	(%)	Ь ((%)	Ca:P	Na ()	(mdd)	×	(8)	Mg	(%)
	Mean	SD	Mean	SD		Mean	SD	Mean	SD	Mean	SD
Rubiaceae											
(71)	0.80	0.35	0.38	0.05	2.11	58	3.1	1.82	0.58	0.27	0.03
(72)	0.53	ı	0.27	1	1.96	49	1	2.43	1	0.32	1
(73)	0.36	ı	0.19	í	1.89	58	1	2.12	ī	0.18	,
(74)	1.16	0.19	0.14	0.02	8.29	101	39.3	2.24	0.18	0.41	0.04
(75)	0.84	1	0.31	-	2.71	92	1	3.15	1	0.33	1
Rutaceae											
(76)	0.82	1	0.12	1	6.83	46	1	0.98	1	0.72	1
(77)	0.68	0.01	0.16	0.02	4.25	93	11.8	06.0	0.02	0.88	0.17
(78)	0.40	0.01	0.16	0.04	2.50	195	161.9	1.47	0.44	0.39	0.01
(79)	0.66	-1	0.24	1	2.75	119	1	1.85	1	0.47	,
Sapindaceae											
(80)	2.30	0.49	0.17	0.04	13.53	152	91.2	0.99	0.29	0.47	0.12
Sterculiaceae											
(81)	1.56	1	0.25	-	6.24	37	1	1.95	ı	0.48	1
(82)	0.99	1	0.21	-	4.71	28	1	2.31	ì	0.29	1
(83)	0.80	0.08	0.18	0.07	4.44	75	23.6	2.00	0.72	0.24	0.02

Table A14. Macro mineral composition of plants consumed by red howler monkeys (Alouatta seniculus seniculus) in the central llanos region of Venezuela.

Family/ Species	Species	Part*	Season	п	Dry matter (%)	ter (%)
	no.				Mean	SD
Sterculiaceae (Cont'd)						
Guazuma ulmifolia	(82)	FL	Wet	1	19.0	-
Sterculia apetala	(88)	FL	Dry	1	ND	,
Sterculia apetala	(87)	FLBU	Dry	2	20.7	4.48
Sterculia apetala	(88)	FL (BU)	Dry	1	21.3	-
Sterculia apetala	(68)	ST	Dry	1	ND	,
UNIDENTIFIED PLANTS						
Unidentified #02	(06)	1	Dry	2	34.0	0.95
Unidentified #03	(91)	T	Dry	1	21.4	,

Macro mineral composition of plants consumed by red howler monkeys (Alouatta seniculus 0.11 0.15 SD 1 i Mq (8) 2.53 0.47 0.40 0.59 0.39 Mean Ð Q 0.53 69.0 SD K (8) 3.33 2.59 2.53 2.32 2.57 Mean Q 8.47 20.83 SD ı Na (ppm) 66 68 149 105 41 Mean Q Ð 1.89 2.12 90.6 1.38 1.81 Ca: P Q 0.02 0.01 seniculus) in the cntral lanos rgion of Venezuela. SD ı P (8) 0.29 0.46 0.31 0.26 0.16 Mean QN ND 0.19 0.32 SD Ca (8) 1.45 0.40 0.87 0.56 0.55 Mean R R Table A14 (cont'd). Family/ Sample Sterculiaceae UNIDENTIFIED (16) (87) (88) (88) (06) (88) (88)

1-mature leaf; NL-new leaf; YL-young leaf; F-fruit; UF-unripe fruit; RF-ripe fruit; FL-flower; PET-petiole; PUL=pulvinus

Family/ Species	Species	Part	Season	п	Dry matter	ter (%)
	no.				Mean	SD
Anacardiaceae						
Mangifera indica	(1)	YL	Wet	1	20.6	1
Mangifera indica	(2)	UF	Dry	4	22.5	00.00
Mangifera indica	(3)	UF	Wet	1	19.0	,
Mangifera indica	(4)	RF	Dry	е	22.1	2.09
Mangifera indica	(5)	PUL	Dry	2	31.8	0.40
Annonaceae						
Annona jahnii	(9)	ı	Wet	1	28.3	,
Annona jahnii	(7)	UF	Wet	1	30.2	,
Annona jahnii	(8)	RF	Wet	1	27.2	,
Annona jahnii	(6)	FL	Wet	1	19.1	,
Annona punicifolia	(10)	UF	Wet	1	16.0	1
Annona punicifolia	(11)	RF	Wet	ю	36.8	25.90
Arecaceae						
Copernicia tectorum	(12)	Ĺ	Wet	2	28.2	2.90
Copernicia tectorum	(13)	UF	Dry	е	33.1	3.45
Copernicia tectorum	(14)	UF	Wet	1	QN	-
minotopo injunction	13.67			,		

Family/ Sample	Fe (I	(mdd)	(mdd) no	(mdc	Mn ()	(mdd)	Se ((mdd)	(mdd) uz	(mdc
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Anacardiaceae										
(1)	35.19	1	13.26	1	113.40	1	0.11	1	30.82	1
(2)	10.57	1.26	4.60	0.55	21.28	1.40	0.03	0.01	5.45	0.40
(3)	28.70	1	6.10	1	14.78	1	0.04	1	6.89	1
(4)	16.56	3.00	4.99	0.82	28.53	8.51	0.02	0.01	9.56	4.26
(5)	28.37	2.08	4.54	0.65	348.56	6.94	0.08	00.00	57.94	5.46
Annonaceae										
(9)	154.90	1	16.59	1	690.03	1	0.10	1	21.99	1
(7)	31.88	1	13.28	1	193.40	1	0.01	1	19.40	,
(8)	37.02	1	19.11	1	29.64	1	0.03	1	16.34	1
(6)	76.82	,	29.70	1	223.99	1	0.01	1	35.65	1
(10)	21.34	1	17.07	1	13.60	1	0.06	1	17.84	1
(11)	28.23	4.14	14.73	3.15	10.74	1.07	0.05	0.02	14.14	1.19
Arecaceae										
(12)	34.66	13.80	8.81	2.54	67.01	35.65	1.33	0.77	25.70	15.16
(13)	29.25	6.00	10.95	1.39	51.34	11.36	0.68	0.34	15.75	0.94
(14)	235.49	1	11.08	1	32.71	1	0.20	1	17.88	1
(15)	50.44	1	7.64	,	18.84	1	0.18	,	16.62	1

Table A16. Trace mineral composition (dry basis) of plants consumed by red howler monkeys (Alouatta seniculus

Family/ Species	Species	Part	Season	п	Dry matter (%)	er (%)
	no.				Mean	SD
Asclepidaceae						
Marsdenia undulata	(16)	ц	Wet	3	33.4	21.84
Marsdenia undulata	(17)	UF	Wet	1	10.0	
Вотрасаеае						
Ceiba pentandra	(18)	FLBU	Dry	1	20.2	1
Cactaceae						
Pereskia guamacho	(19)	RF	Wet	1	25.3	1
Chrysobalanaceae						
Licania pyrifolia	(20)	FL(BU)	Wet	1	30.6	
Licania pyrifolia	(21)	NL	Dry	1	23.9	,
Licania pyrifolia	(22)	ı	Dry	2	40.8	0.44
Cucurbitaceae						
Melothria trilobata	(23)	ı	Wet	2	17.2	0.08
Dilleniaceae						
Tetracera volubris	(24)	FL	Wet	1	22.8	1
Euphorbiaceae						
Sapium aucuparium	(25)	I	Wet	т	67.0	2.41

Table A16 (cont'd). Trace mineral composition (dry basis) of plants consumed by red howler monkeys (Alouatta seniculus seniculus) in the central llanos region of Venezuela.

Family/ Sample	Fe ()	(mdd)	(mdd) no	(mdd	Mn	Mn (ppm)	Se	(mdd)	(mdd) uz	(md
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Asclepidaceae										
(16)	199.64	66.62	16.63	3.49	944.92	443.52	0.04	0.02	27.77	1.82
(17)	25.65	1	8.13	1	66.97	1	0.04	,	14.64	1
Bombaceae										
(18)	30.92	1	15.79	1	22.33	1	0.05	1	17.41	,
Cactaceae										
(19)	26.51	-	9.19	1	42.38	1	0.03	-	30.09	1
Chrysobalanaceae										
(20)	49.97	1	17.68	1	125.49	1	0.04	1	18.62	1
(21)	133.92	1	16.03	1	66.91	1	0.21	1	25.88	1
(22)	52.37	6.58	11.02	0.63	200.27	8.99	0.15	0.04	16.84	0.88
Cucurbitaceae										
(23)	142.80	40.11	13.64	0.81	104.64	70.20	0.05	0.01	34.25	4.75
Dilleniaceae										
(24)	42.29	1	13.91	1	83.52	1	0.08	1	21.55	1
Euphorbiaceae										
(25)	93.23	16.40	10.07	1.76	86.08	4.26	0.06		0.00 100.90	6.41

Table A17. Trace mineral composition (dry basis) of plants consumed by red howler monkeys (Alouatta seniculus seniculus) in the central llanos region of Venezuela.

Family/ Species	Species	Part.	Season	r,	Dry matter	ter (%)
	no.				Mean	SD
Fabaceae (Caesalpinoideae)						
Cassia grandis	(26)	ı	Wet	ю	51.6	12.13
Cassia grandis	(27)	UF	Wet	3	10.3	1.63
Cassia moschata	(28)	ı	Wet	2	45.8	4.06
Cassia moschata	(29)	ı	Dry	1	48.3	
Cassia moschata	(30)	YL	Wet	1	32.9	1
Cassia moschata	(31)	YL	Dry	2	36.7	2.88
Cassia moschata	(32)	UF	Wet	8	20.3	1.49
Fabaceae (Mimosoideae)						
Albizia columbiana	(33)	ı	Wet	1	81.0	
Albizia guachapele	(34)	I	Wet	2	35.4	1
Albizia guachapele	(35)	I	Dry	1	37.1	
Albizia guachapele	(36)	YL	Wet	1	28.7	1
Albizia guachapele	(37)	NL	Wet	8	31.0	1.98
Enterolobium cyclocarpum	(38)	[by	Wet	1	-	-
Pithecellobium dualense	(39)	T	Wet	2	45.3	2.21
Pithecellobium dualense	(40)	FL/BU	Wet	2	22.2	0.74
Pithecellobium saman	(41)	ц	Wet	7	38.3	7.44

Table A17 (cont'd). Trace mineral composition (dry basis) of plants consumed by red howler monkeys (Alouatta seniculus seniculus) in the central llanos region of Venezuela.

Family/ Sample	Fe ((mdd)	Cu (F	(mdd)	Mn ()	(mdd)	Se (I	(mdd)	d) uz	(mdd)
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Fabaceae										
(26)	220.03	158.28	4.77	0.85	66.52	47.53	0.23	0.15	12.95	3.06
(27)	34.03	4.00	10.23	0.46	21.41	1.86	0.21	0.27	18.03	0.87
(28)	251.62	95.87	5.63	0.28	331.70	109.88	0.11	0.01	23.21	6.99
(29)	178.50	-	5.32	1	224.64	-	0.05	-1	5.32	1
(30)	29.55	1	8.27	1	123.24	,	0.08	1	14.25	1
(31)	174.72	4.89	9.96	1.95	50.19	25.98	0.08	0.03	15.64	3.68
(32)	28.48	2.76	7.01	0.43	51.01	6.53	0.04	0.02	11.00	2.82
Fabaceae										
(33)	197.41	1	10.00	1	458.82	1	0.15	1	15.29	1
(34)	80.56	21.89	7.79	3.26	42.19	10.94	0.10	00.00	23.18	10.61
(35)	146.78	1	8.02	,	41.14	1	0.41	1	18.19	ı
(36)	76.24	1	6.94	1	27.72	1	0.87	1	32.03	,
(37)	92.59	18.94	15.02	1.81	25.72	15.23	0.40	0.29	50.60	9.51
(38)	1	1	1	1	1	1	1	1	1	ı
(39)	147.57	37.41	13.52	2.93	505.70	45.15	0.31	0.13	12.48	1.72
(40)	48.53	0.97	27.80	0.58	59.56	2.21	0.32	0.01	53.86	0.83
(41)	118.02	22.87	9.59	2.47	33.19	7.27	0.19	0.06	9.03	3.48
- Lemature leaf; NL-new leaf; YL-young leaf; F-fruit; UF-unripe fruit; RF-ripe fruit; PUF-pulvinus	new leaf;	YL-young 1	.eaf; F=fr	uit; UF=	unripe fr	uit; RF=1	ripe frui		wer; PET	=petiole;
(41) L=mature leaf; NL=r PUL=pulvinus	118.02 new leaf;	22.87 YL=young 1	9.59 eaf; F=fr	2.47	33.19 unripe fr	7.27 uit; RF=1		0.19		0.06 FL-flower

Trace mineral composition of plants consumed by red howler monkeys (Alouatta seniculus seniculus) Table A18.

in the central llanos region of Venezuela.			e kaumam	(Atouatea	ra senicains	Sentcatus
Family/ Species	Species	Part	Season	n	Dry matter	ter (%)
	no.				Mean	SD
Fabaceae (Mimosoideae)						
Pithecellobium saman	(42)	L	Dry	2	36.9	1.06
Pithecellobium saman	(43)	NL	Dry	1	ND	ND
Pithecellobium saman	(44)	YL	Wet	3	24.5	1.29
Pithecellobium saman	(45)	FLBU	Dry	1	ND	ND
Pithecellobium saman	(46)	FL	Dry	2	19.0	1.00
Pithecellobium saman	(47)	RF	Dry	1	44.0	1
Pithecellobium saman	(48)	UF	Dry	2	22.5	•
Pithecellobium tortum	(49)	L	Wet	5	38.0	3.77
Pithecellobium tortum	(20)	YL	Wet	1	22.2	-
Pithecellobium tortum	(51)	FL	Wet	1	18.8	1
Pithecellobium tortum	(52)	BU, NL	Dry	2	41.8	0.11
Pithecellobium tortum	(53)	UF	Dry	1	71.4	-
Fabaceae (Papilionoideae)						
Entada polystachya	(54)	L	Wet	1	43.7	1
Machaerium moritzianum	(55)	FL	Dry	1	ND	1
Platymiscium diadelphum	(56)	L	Wet	1	37.5	1
Platymiscium diadelphum	(57)	YL, NL	Wet	1	31.2	-

Table Al8 (cont'd). Trace mineral composition of plants consumed by red howler monkeys (Alouatta seniculus seniculus) in the central llanos region of Venezuela.

Fe ((wdd	I) no	(mdc	Mn ()	(mdd	Se ()	(mdc	Zu (F	(mdi
Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
74.71	4.65	10.59	0.67	17.48	0.29	0.23	0.09	11.26	0.34
162.04	-	22.54	_	17.49	-	0.25	-	45.81	-
67.44	9.10	14.20	2.63	18.23	1.33	0.37	0.12	21.86	5.89
88.31	-	21.45	-	26.71	-	0.37	-	47.64	-
163.05	105.71	13.53	1.19	19.62	1.02	0.43	0.33	23.19	1.05
19.54	1	6.58	-	11.45	-	0.12		7.04	,
26.98	3.13	7.67	0.65	10.21	4.68	0.14	0.02	14.20	0.54
241.98	248.43	8.12	1.04	179.79	45.94	0.09	0.03	8.01	1.41
85.53	-	24.21	-	108.27	-	0.06	1	27.77	1
47.87	-	23.36	-	59.23	-	0.09	-	29.22	-
51.69	8.39	21.47	7.81	56.28	20.67	0.23	0.17	28.75	10.42
1.27	-	12.58	-	40.39	-	0.79	1	10.44	-
	Fe (Mean 74.71 162.04 67.44 88.31 163.05 19.54 24.38 85.53 47.87 51.69	100 10 24	(ppm) SD Me SD Me 4.65 10 - 22 9.10 14 - 21 105.71 13 248.43 8 8.39 21 - 23	(ppm) Cu (ppm) SD Mean 4.65 10.59 - 22.54 - 22.54 - 13.53 - 6.58 - 6.58 - 6.58 - 248.43 8.12 - 248.21 - 23.36 8.39 21.47 - 12.58	(ppm) Cu (ppm) Mean SD	CPPM CU (PPM) Mn (PPPM SD Mean SD SD SD SD SD SD SD S	(Ppm) Cu (Ppm) Mn (Ppm) Mn (Ppm) Me Mn (Ppm) Mn (Ppm	Cppm Cu (ppm Nean Ppm Ppm SD Nean SD SD Nean SD SD SD SD SD SD SD S	Cpm Cu (ppm) Mn (ppm) Se (ppm) Se (ppm) Se (ppm) Sp Mean Mean Sp Mean Mean

L-mature leaf; NL-new leaf; YL-young leaf; F-fruit; UF-unripe fruit; RF-ripe fruit; FL-flower; PET-petiole; 49.75 31.38 0.07 0.10 111.66 53.67 20.94 23.40 57.03 128.87 PUL=pulvinus (99) (57)

13.63

0.03

0.52

22.06

86.69

11.44

102.20

(24)

(52)

59.62

20.72

Trace mineral composition of plants consumed by red howler monkeys (Alouatta seniculus seniculus) in the central llanos region of Venezuela. Table A19.

Banks

Choose

Family/ Species

rainty/ species	Species		Fart Season	п	Dry matter (%)	ter (%)	
	no.				Mean	SD	
Fabaceae (Papilionoideae) (cont'd)							
Platymiscium polystachyum	(58)	YL	Wet	1	21.2		
Platymiscium polystachyum	(59)	FLBU	Wet	1	17.9		
Pterocarpus acapulcensis	(60)	r	Wet	4	33.3	8.49	
Pterocarpus acapulcensis	(61)	NE	Wet	4	27.6	1.77	
Moraceae							
Ficus pertusa	(62)	T	Dry	3	26.5	16.32	
Ficus pertusa	(63)	ı	Wet	4	29.6	3.17	2.

Ficus pertusa Ficus pertusa Ficus pertusa Ficus pertusa

⊣ 1	0
- 1	

23.1 22.8 16.3

2

Wet Wet Wet

GE RF GE

(64)

(69) (99)

Ficus trigonata

1.09

43.5 35.0 39.8

Wet Dry

п ы [su

(89) (69) (20)

Zizyphys saeri Zizyphys saeri Zizyphys saeri

Coccoloba sp.

Polygonaceae Rhamnaceae

XL

(67)

Wet

Trace mineral composition of plants consumed by red howler monkeys (Alouatta seniculus Table A19 (cont'd).

Family/ Sample	Fe	(mdd)	Cu ((mdd) no	Mn ((mdd)	Se	Se (ppm)	Zn	(mdd) uz
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Fabaceae										
(58)	60.14	1	28.84	1	34.70	1	0.24	1	57.23	,
(59)	65.11	1	35.49	1	30.58	1	0.37	1	58.52	
(09)	320.78	159.92	16.37	5.62	273.45	55.72	0.27	0.02	24.78	9.29
(61)	97.33	14.49	20.05	2.14	145.92	122.50	0.08	0.01	37.12	2.35
Moraceae										
(62)	162.08	14.32	5.89	0.98	34.27	16.61	0.31	0.06	13.62	2.37
(63)	238.13	102.87	7.20	1.00	105.37	19.42	0.04	0.01	16.54	1.13
(64)	81.30	5.76	13.09	1.22	43.69	1.57	0.05	0.02	21.14	1.26
(65)	56.62	4.33	9.15	1.59	30.14	0.68	0.03	0.01	14.17	1.96
(99)	49.93	1	12.98	-1	42.48	1	0.10	1	26.88	,
Polygonaceae										
(67)	1	-	1	1	1	1	1	1	1	-
Rhamnaceae										
(68)	90.09	1	20.22	1	119.85	1	17.59	1	40.70	,
(69)	215.32	1	9.91	-	66.19	1	1.02	,	20.70	,
(70)	51.54	1	8.86	,	154.10	1	77.40	,	10.43	,

Table A20. Trace mineral composition of plants consumed by red howler monkeys (Alouatta seniculus seniculus) 3.79 6.84 Dry matter (%) SD 23.0 18.0 19.9 29.9 18.7 Mean C Н 2 Season Wet Wet Wet Wet Wet Part XL XL N UF н Species (71) (72) (74) (75) (73) no. in the central llanos region of Venezuela. Randia venezuelensis Randia venezuelensis Randia venezuelensis Unidentified #01 Unidentified #01

Family/ Species

Rubiaceae

2.	55	
	0.55	
74.9	33.4	
1	2	
Wet	Wet	
L	YL, L	
(16)	(77)	

Zanthoxylum culantrillo

Rutaceae

Paullinia Sterculiaceae Sapindaceae

Zanthoxylum culantrillo	(77)	YL, L	Wet	2	33.4	0.55	5
Zanthoxylum culantrillo	(78)	YL	Wet	2	31.2	4.06	
Zanthoxylum culantrillo	(79)	FLBU	Wet	2	31.7	,	
ndaceae							
Paullinia cururu	(80)	I	Wet	4	41.3	2.00	
culiaceae							

11.50

35.2

40.60

18.6

7 2 4

Wet

(81) (82) (83) (84)

Guazuma ulmifolia Guazuma ulmifolia Guazuma ulmifolia Guazuma ulmifolia

Dry Dry Dry

UF

RF FL

25.0 58.3

Family/ Sample	Fe (I	(mdd)) no	(mdd)	Mn (ppm)	(mdd	Se ()	(mdd)	(mdd) uz	(mdd
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Rubiaceae										
(71)	100.67	10.59	19.50	5.53	85.26	12.93	0.14	00.00	24.94	6.04
(72)	91.60	1	13.33	1	79.99	1	0.20	1	19.15	1
(73)	37.68	1	7.94	1	42.27	1	0.05	1	8.48	1
(74)	172.23	69.88	18.04	1.33	52.78	7.68	0.09	0.05	26.20	2.21
(75)	94.92	-	26.08	1	61.19	1	0.20	1	30.99	,
Rutaceae										
(16)	323.75	1	13.49	,	721.64	1	0.06	1	17.83	,
(77)	141.16	4.65	13.07	2.96	372.52	7.20	0.18	0.21	41.69	30.90
(78)	126.01	4.45	10.99	5.24	274.87	7.79	0.07	0.02	13.50	4.74
(19)	64.01	1	20.97	1	390.85	-	0.04	1	27.22	1
Sapindaceae										
(80)	358.11	134.34	14.15	0.67	112.57	34.78	0.16	0.14	16.48	1.89
Sterculiaceae										
(81)	40.03	1	16.05	1	59.89	1	0.03	1	23.90	1
(82)	42.07	1	12.50	1	31.17	1	0.03	1	18.41	
(83)	46.04	5.15	21.40	3.20	42.35	4.69	0.04	0.02	15.88	2.05
(84)	341.55	78.99	17.86	1.66	74.21	6.17	0.15	0.18	24.85	2.34

Table A21. Trace mineral composition of plants consumed by red howler monkeys (Alouatta seniculus seniculus) in the central llanos region of Venezuela.

Family/ Species	Species	Part.	Season	и	Dry matter (%)	ter (%)
	no.				Mean	SD
Sterculiaceae (Cont'd)						
Guazuma ulmifolia	(85)	FL	Wet	1	19.0	-
Sterculia apetala	(86)	FL	Dry	1	ND	1
Sterculia apetala	(87)	FLBU	Dry	2	20.7	4.48
Sterculia apetala	(88)	FL(BU)	Dry	1	21.3	-
Sterculia apetala	(89)	ST	Dry	1	ND	1
UNIDENTIFIED PLANTS						
Unidentified #02	(06)	I	Dry	2	34.0	0.95
Unidentified #03	(91)	ı	Dry	1	21.4	1

Trace mineral composition of plants consumed by red howler monkeys (Alouatta seniculus Table A21 (cont'd).

Family/ Sample Fe (ppm) Mean SD Sterculiaceae (85) 145.24 -	Ω .	Cu (ppm)		,	1	,	•		
Mean 145.24	$\dashv H$	fean		(mdd) um	Duit)	Se (ppm)	(mdd	(mdd) uz	(mdc
145.24 ND			SD	Mean	SD	Mean	SD	Mean	SD
145.24 ND									
CN		18.02	1	36.45	1	0.05	-	31.00	1
		ND	1	ND	-	ND	1	ND	-
(87) 82.42 4.6	4.64	13.04	0.92	70.50	30.56	0.08	0.01	41.74	4.51
- 73.07 -		16.18	1	17.37	1	0.07	_	48.52	1
- QN (68)		ND	1	ND	1	ND	_	ND	1
UNIDENTIFIED									
(90) 159.24 7.	7.30	13.20	0.77	42.99	6.96	0.34	0.01	44.31	8.84
(91) 97.55	1	19.89	1	42.81	ı	0.49	ı	39.77	ı

* L-mature leaf; NL-new leaf; YL-young leaf; F-fruit; UF-unripe fruit; RF-ripe fruit; FL-flower; PET-petiole; PUL=pulvinus

APPENDIX B

5.80

6.55

6.85

6

Waldo

Table B1. Body weights (BW) of study subjects prior to, during, and following feeding 4.05 3.60 4.25 4.45 3.80 7.60 Post-study Body weight (kg) 4.05 3.70 4.25 4.40 3.80 8.55 7.90 6.05 4.30 Crossover 5.30 4.95 5.00 9.15 7.50 5.15 4.50 4.30 4.65 4.72 8.94 7.44 10.66 Pre-study Sex ъ ъ Jalapeno Pilgrim Michele Jessie Shadow Phoebe Athena Rinda Katie Howie Spica Night Ino ΩI Varecia variegata variegata Alouatta villosa palliata Alouatta seniculus sara Varecia variegata rubra Alouatta caraya Species trials.

Table B1 (cont'd). Body weights (BW) of study subjects prior to, during, and following 5.76 7.25 8.90 10.65 7.71 5.72 3.99 12.40 10.30 Post-study Body weight (kg) 10.90 6.75 6.05 12.10 10.10 10.00 Crossover 10.70 6.67 3.76 5.75 5.85 11.75 9.60 10.20 11.34 5.67 Pre-study Sex ъ ъ O+ ъ Chrissie GooNgoot Stanley Fungar Flower Susie Rhett Heine Gomez Janet Jack Cercopithecus g. chrysogaster Colobus guereza kikuyuensis Trachypithecus f. francoisi Pygathrix nemaeus nemaeus Mandrillus 1. leucophaeus feeding trials. Species

Table B2. Body weight (BW) and dry matter intake (DMI) of ruffed lemurs (Varecia spp.) fed diet 15ADF and apparent digestibility of dry matter (DM) and fiber fractions in that diet.

Species	ID	BW	DMI		Dige	Digestibility (%)		ofª	
		(kg)	(g/kg BW)	DM	NDF	ADF	υ	HC	ADL
Varecia variegata rubra	Athena	4.50	32.5	51.9	22.2	8.9	13.3	43.7	23.2
	Michele	4.30	21.5	56.7	24.2	17.6	15.2	34.8	33.8
	Rinda	5.30	17.8	56.2	26.9	20.4	17.0	37.3	22.8
	Spica	4.95	36.0	51.2	22.4	9.6	17.3	42.4	48.1
	Mean	4.76	26.9	54.0	23.9	14.2	15.7	39.5	32.0
	SD	0.45	8.7	2.9	2.2	5.7	1.9	4.2	11.9
Varecia variegata variegata	Ino	4.65	29.8	43.7	15.6	5.2	8.7	35.6	27.2
	Night	5.00	25.2	51.6	17.7	5.5	3.1	37.3	83.1
	Mean	4.83	27.5	47.7	16.6	5.3	5.9	36.5	55.1
	SD	0.25	3.3	5.6	1.5	0.2	4.0	1.2	39.5
Varecia variegata ssp.	Mean	4.78	27.1	51.9	21.5	11.3	12.4	38.5	39.7
	SD	0.37	6.9	4.7	4.2	6.3	5.5	3.7	23.3
*DM=drv matter; NDF=neutral-detergent fiber;	ergent fiber;		ADF=acid-detergent fiber;	er; C=C	C=cellulose;	H	HC=hemicellulose;	li li	ADL=acid-

detergent lignin

Table B3. Body weight (BW) and dry matter intake (DMI) of ruffed lemurs (Varecia spp.) fed diet 30ADF and apparent digestibility of dry matter (DM) and fiber fractions in that diet.

Species	ID	ВЖ	DMI		Dige	Digestibility (%)		of*	
		(kg)	(g/kg BW)	DM	NDF	ADF	ນ	нс	ADL
Varecia variegata rubra	Athena	4.05	45.2	43.5	23.3	14.7	12.6	42.6	62.3
	Michele	3.70	38.7	45.2	23.3	15.2	7.6	41.4	59.2
	Spica	4.25	53.0	34.8	10.5	1.8	-2.6	30.0	62.8
	Mean	4.00	45.6	41.2	19.0	10.6	9.9	38.0	61.4
	SD	0.28	7.2	5.6	7.4	7.6	8.1	7.0	2.0
Varecia variegata variegata	Ino	4.40	63.8	48.8	29.3	24.1	20.0	40.8	28.0
	Night	3.80	73.9	36.2	17.3	7.4	3.8	39.8	37.1
	Mean	4.10	8.89	42.5	23.3	15.8	11.9	40.3	32.5
	SD	0.42	7.1	8 6.	8.5	11.8	11.5	0.7	6.4
Varecia variegata ssp.	Mean	4.04	54.9	41.7	20.7	12.6	8.7	38.9	49.9
	SD	0.30	14.2	6.0	7.1	8.5	8.6	5.1	16.2
*DM=dry matter; NDF=neutral-detergent fiber; ADF=acid-detergent fiber; C=cellulose;	ergent fiber;	ADF=aci	d-detergent fib	er; C=c	llulose	HC=hen	HC=hemicellulose;		ADL=acid

2 -6 detergent lignin

Table B4. Body weight (BW) of ruffed lemurs (*Varecia* spp.) fed diets 15ADF and 30ADF and the apparent digestibility of dry matter and gross energy in those diets.*

Species	ID	BW		15ADF			30ADF	
		(kg)	DM (%)	GE (%)	DE (kcal/g)	DM (%)	GE (%)	DE (kcal/g)
Varecia variegata rubra	Athena	4.50	51.9	43.0	1.92	43.5	41.6	1.82
	Michele	4.30	56.7	54.5	2.43	45.2	44.9	1.97
	Rinda	5.30	56.2	52.1	2.37	ı	ı	I
	Spica	4.95	51.2	48.9	2.18	34.8	30.7	1.35
	Mean	4.76	54.0	49.6	2.23	41.2	39.1	1.71
	SD	0.45	2.85	4.98	0.23	5.58	7.43	0.32
Varecia variegata variegata	Ino	4.65	43.7	43.0	1.92	48.8	46.9	2.06
	Night	5.00	51.6	45.7	2.57	36.2	34.6	1.52
	Mean	4.83	47.7	44.4	2.25	42.5	40.8	1.79
	SD	0.25	5.59	1.91	0.46	8.91	8.70	0.38
Varecia variegata ssp.	Mean	4.78	51.9	47.9	2.23	41.7	39.7	1.74
	SD	0.37	4.68	4.80	0.27	5.99	6.88	0.30
		l						

*DM=dry matter; GE=gross energy; DE=digestible energy

Table B5. Body weight (BW) and dry matter intake (DMI) of howler monkeys (Alouatta spp.) fed diet 15ADF and 68.0 28.7 57.4 67.2 55.3 18.4 58.2 76.7 67.5 8.5 60.8 13.1 59.1 65.1 14.7 71.1 ADL 48.6 6.9 54.8 49.0 61.8 7.6 49.4 54.3 3.5 39.0 43.8 50.1 6.8 HC Digestibility (%) of 49.5 42.5 36.6 57.4 39.0 10.8 36.4 37.4 37.9 8.8 39.5 4.2 24.1 40.8 21.2 28.1 ပ 45.9 56.7 29.5 42.0 29.6 37.9 40.2 19.2 54.1 40.5 10.2 41.7 43.8 3.0 43.1 10.1 ADF 47.3 53.6 44.7 43.5 57.0 40.1 33.2 43.4 14.4 7.9 38.2 44.4 45.8 8.0 43.7 5.1 NDF and fiber fractions in that diet. 69.4 9.04 76.7 65.2 71.0 69.4 70.0 67.9 70.2 70.7 68.3 6.0 8.1 3. 8 2.8 Z 18.5 26.0 11.8 20.4 21.7 5.73 15.6 10.5 23.8 3.05 16.1 24.9 17.1 6.12 17.10 (g/kg BW) DMI 7.50 2.03 5.15 6.55 5.85 0.99 9.15 8.32 1.17 7.51 8.94 7.44 7.94 2.52 4.72 10.66 (kg) dry matter (DM) Jalapeno Pilgrim Jessie Shadow Phoebe Howie Waldo Katie Mean Mean Mean Mean SD SD ΩI Alouatta villosa palliata digestibility of Alouatta seniculus sara Alouatta caraya Alouatta spp. apparent Species

Alouatta caraya Ratie Shadow Mean SD Allouatta villosa palliata Waldo Mean	(kg) 4.72 8.94 10.66 8.11 3.06	(g/kg BW) 38.4 18.7 19.4 25.5 11.18	61.0 63.7 61.4 62.0	NDF 41.2 50.6 45.5 45.8	ADF 34.2	ບ	HC	
, , , , , , , , , , , , , , , , , , , ,	4.72 8.94 10.66 8.11 3.06	38.4 18.7 19.4 25.5 11.18	61.0 63.7 61.4 62.0	41.2 50.6 45.5 45.8	34.2)	ADL
, , , , , ,	8.94 10.66 8.11 3.06	18.7 19.4 25.5 11.18	63.7 61.4 62.0	50.6 45.5 45.8	1	33.0	56.8	38.3
, , , , , ,	10.66 8.11 3.06	19.4 25.5 11.18	61.4	45.8	43.3	42.9	66.7	57.3
7 7 7	3.06	25.5	62.0	45.8	35.7	34.5	67.3	56.2
1 - 1	3.06	11.18	1.5	7 7	37.7	36.8	63.6	50.6
		α-		· r	4.9	5.3	5.9	10.7
	<u></u>		2 62	ر بر	ر د	4	æ	7
Mean) (• ()) () (• L		•
Mean	6.85	7.77	91.6	49.2	42.1	38.5	64.9	60.3
US	6.45	20.3	62.1	52.5	46.2	44.5	8.99	62.3
	0.57	2.67	0.7	4.7	5.7	8.4	5.6	2.8
Alouatta seniculus sara Pilgrim	8.55	19.0	61.0	47.0	43.6	42.4	54.5	56.2
Howie	7.50	20.4	60.3	42.5	35.3	32.1	58.3	59.7
Mean	8.03	19.7	9.09	44.8	39.5	37.3	56.4	58.0
QS	0.74	0.95	0.5	3.2	5.9	7.3	2.7	2.5
Alonatta spp	7.61	2.2.4	61.7	47 4	40.6	39.1	62.4	56.0
	1.97	7.2	1.1	5.0	5.8	9.9	5.7	8.3

Body weight (BW) howler monkeys (Alouatta spp.) fed diets 15ADF and 30ADF and the apparent (kcal/g) 2.66 2.83 2.70 2.62 2.70 0.11 2.71 2.71 2.71 0.00 2.71 2.65 2.68 0.04 0.07 30ADF (&) 61.6 2.53 61.7 61.8 61.7 60.5 0.85 61.5 64.5 59.7 1.53 0.07 61.1 60.7 GE GE 1.46 62.6 61.6 (& 62.0 62.1 0.71 63.4 60.3 61.8 2.19 61.0 62.0 61.4 1.27 63.7 ¥ (kcal/q) 3.06 3.45 3.08 2.93 3.19 3.19 3.06 3.02 0.01 2.79 0.17 0.37 0.19 3.03 3.07 3.07 15ADF (%) 68.6 68.6 0.69 71.6 3.75 68.89 0.28 77.3 65.7 71.5 8.20 68.89 4.29 62.6 68.0 67.7 GE digestibility of dry matter and gross energy in those diets. DM (%) 9.01 71.0 64.5 67.9 68.3 70.0 0.85 76.7 65.2 8.13 69.4 3.80 70.2 70.7 2.83 69.4 9.15 2.03 8.94 7.44 10.66 6.55 5.85 0.99 7.50 4.72 7.94 5.15 8.32 1.17 7.51 2.52 (kg) BW Jalapeno Pilgrim Phoebe Jessie Shadow Katie Waldo Howie Mean Mean Mean Mean ID SD SD SD SD Alouatta villosa palliata Alouatta seniculus sara Alouatta caraya Alouatta spp. Table B7. Species

Table B8. Body weight (BW) and dry matter intake (DMI) of colobines fed diet 15ADF and apparent digestibility

of dry matter (DM) and fi	fiber fractions in that	in that di	diet.					_	
Species	ID	BW	DMI		Dige	Digestibility (%)	ty (%)	ofa	
		(kg)	(g/kg BW)	MO	NDF	ADF	ບ	нс	ADL
Colobus g. kikuyuensis	Flower	11.34	16.3	73.5	39.7	30.2	40.2	54.9	-16.3
	Gomez	10.90	14.2	80.3	77.0	80.1	81.4	67.9	84.5
	Mean	11.12	15.3	76.9	58.4	55.1	8.09	61.4	34.1
	SD	0.31	1.48	4.8	26.4	35.3	29.1	9.5	71.3
Pygathríx n. nemaeus	Rhett	12.10	35.5	76.2	66.5	9.99	59.7	66.4	78.4
Trachypithecus f.	Jack	7.71	23.7	84.6	80.5	81.4	82.2	79.1	83.7
francoisi	Janet	5.67	24.8	79.7	77.7	81.4	90.4	71.8	51.8
	Chrissie	6.67	32.7	82.9	83.4	88.2	88.3	71.9	83.6
	Stanley	3.99	42.3	85.9	83.9	85.0	87.6	82.2	87.8
	Fungar	5.75	34.7	78.9	70.9	75.4	73.1	63.7	71.8
	Mean	5.96	31.6	82.4	79.3	82.3	84.3	73.7	75.7
	SD	1.38	7.65	3.0	5.3	4.8	7.0	7.2	14.7
Colobines	Mean	8.02	28.0	80.3	72.5	73.5	75.4	69.7	65.7
	SD	3.04	9.88	4.1	14.5	18.7	17.4	8.6	35.1

*DM=dry matter; NDF=neutral-detergent fiber; ADF=acid-detergent fiber; C=cellulose; HC=hemicellulose; ADL=acid-detergent lignin

Table B9. Body weight (BW) and dry matter intake (DMI) of colobines fed diet 30ADF and apparent digestibility of dry matter (DM) and fiber fractions in that diet.

	ID	ВМ	DMI		Dig	Digestibility (%)	ity (%)	of*	
		(kg)	(g/kg BW)	DM	NDF	ADE	ນ	нс	ADL
Colobus g. kikuyuensis	Gomez	10.70	14.4	78.7	74.3	56.2	76.1	78.0	77.2
Pygathríx n. nemaeus	Rhett	11.75	33.3	73.9	8.69	67.6	66.1	74.8	82.5
${\it Trachypithecus}$ f.	Jack	7.78	23.1	79.9	82.6	84.5	87.6	78.2	61.9
francoisi	Janet	5.72	18.1	74.4	81.5	83.2	89.8	7.77	49.4
	Chrissie	5.76	38.2	77.0	74.8	72.3	72.7	80.4	78.4
	Stanley	3.77	37.8	80.1	76.6	76.0	76.5	9.77	71.4
	Fungar	6.75	35.2	72.0	63.2	59.2	57.9	72.2	70.1
	GooNgoot	5.85	38.1	81.1	76.3	72.2	73.7	85.3	60.1
	Mean	5.94	31.8	77.4	75.8	74.6	76.4	78.6	65.2
	SD	1.33	8.8	3.6	6.9	9.5	11.6	4 .	10.2
Colobines	Mean	7.26	29.8	77.1	74.9	71.4	75.1	78.1	68.9
	SD	2.71	7.6	3.4	6.2	10.2	10.4	3.8	11.1

"DM=dry matter; NDF=neutral-detergent lignin

3.35 3.50 3.11 3.36 3.18 3.43 3.23 3.54 0.17 3.35 0.16 3.42 3.42 Body weight (BW) of colobines fed diets 15ADF and 30ADF and digestibility of apparent dry matter (kcal/g) A A 30ADF 73.5 79.8 70.9 80.6 9.91 72.5 76.4 3.76 76.2 78.0 78.0 78.1 3.55 (%) KZ GE 73.9 80.0 77.4 77.2 78.7 78.7 74.4 77.0 80.1 72.0 3.63 3.37 (%) Z Z ¥ 3.56 0.14 3.26 3.50 3.49 3.79 3.62 3.38 0.17 3.64 3.72 3.64 3.47 0.17 (kcal/q) 15ADF 81.6 79.9 73.0 75.8 3.89 83.3 78.3 81.7 85.1 7.77 3.18 3.8 78.5 (%) ŧ GE GE 84.6 82.9 85.9 78.9 73.5 80.3 76.9 76.2 79.7 65.5 80.3 4.2 3.00 4.80 (%) M 3.99 5.75 5.85 5.59 7.78 5.67 6.67 1.38 2.94 11.12 7.71 12.10 10.90 0.31 11.34 (kg) BW Chrissie GooNgoot Stanley Fungar Flower Janet Rhett Gomez Jack Mean Mean Mean SD SD SD QI in those diets.a Trachypithecus f. francoisi Colobus g. kikuyuensis Pygathrix n. nemaeus and gross energy Colobines Table B10. Species

5.0 8.8 3.5 5.6 1.9 1.2 2.8 1.96 5.41 3.07 Digesta transit time (\mathtt{TT}_1) and total retention time ($\mathtt{R}_{\mathtt{GIT}}$) in hours of ruffed lemurs (Varecia spp.) $R_{\rm GIT}$ Cr-fiber 1.5 1.50 0.00 4.5 3.0 1.5 1.5 2.5 1.55 1.5 TT_1 2.45 5.5 3.9 1.1 6.4 2.31 1.97 RGIT Marker type CO-EDTA 3.46 3.10 1.5 4.50 1.5 00.0 3.5 $\operatorname{TT}_{_{1}}$ 5.8 3.76 12.3 1.53 3.80 4.5 6.7 3.3 2.2 5.2 $R_{\rm GIT}$ Bead 4.5 1.5 3.0 1.73 2.12 4.5 1.5 4.5 3.0 3.0 1.64 $\mathrm{TT}_{_{1}}$ 4.95 4.76 0.45 4.65 5.00 0.25 4.78 5.30 4.50 4.83 0.37 4.30 (kg) BW Michele Athena Rinda Spica Night Mean Mean Mean Ino SD SD SD ID Varecia variegata variegata Varecia variegata rubra Varecia variegata spp fed diet 15ADF Table B11. Species

3.6 1.6 3.4 2.3 1.03 2.0 0.33 3.1 0.37 $R_{\rm GIT}$ 1.4 1.4 Digesta transit time (TT $_1$) and total retention time ($R_{
m GIT}$) in hours of ruffed lemurs (Varecia.spp.) Cr-fiber 1.73 0.00 4.5 3.5 1.5 1.5 2.7 4.5 1.64 1.5 1 $\mathrm{TT}_{_{1}}$ 1.0 1.0 0.09 2.4 1.37 3.3 0.77 1.1 RGIT 2.5 4.1 Marker type CO-EDTA 1.73 4.5 3.0 2.12 1.5 1.5 2.7 1.64 $\mathrm{TT}_{_{1}}$ ١ 5.5 5.7 2.5 2.3 4.9 2.35 4.8 8.5 0.31 6.1 1.63 2.1 Acetate Bead **7** 3.0 3.00 1.5 7.5 1.5 1.5 1.5 1.5 00.0 2.5 2.45 TT_1 3.80 0.42 0.28 4.25 4.08 0.27 4.40 4.10 4.08 4.05 3.70 4.30 (kg) BW Michele Athena Spica Night Rinda Mean Mean Mean Ino SD SD SD ID Varecia variegata variegata Varecia variegata rubra Varecia variegata ssp. fed diet 30ADF Table B12. Species

39.9

34.5

58.2

22.5

34.5

46.5

8.32

Mean

32.10

16.97

41.15

0.00

0.11

33.94

1.17

SD

30.5

28.0

32.9

14.6

27.1

30.5

7.51

Mean

Alouatta spp

SD

22.3

15.92

23.22

8.48

12.15

20.20

2.03

Digesta transit time (TT,) and total retention time (RGIT) in hours of howler monkeys (Alouatta spp.) 19.9 13.9 $R_{\rm GIT}$ 8.0 37.8 62.6 17.2 54.4 8.41 21.1 23.55 Cr-fiber 28.5 37.5 7.5 16.5 46.5 46.5 12.73 12.0 6.36 22.5 TT_1 1 19.9 28.7 38.5 5.19 33.6 87.3 29.1 22.3 24.7 12.7 $R_{\rm GIT}$ 20.1 6.87 Marker type CO-EDTA 22.5 14.3 22.5 22.5 4.5 9.60 7.5 7.5 $\mathrm{TT}_{_{1}}$ 22.5 7.5 7.5 0.00 33.9 20.9 34.6 34.4 15.4 8.0 25.7 14.66 18.30 $R_{\rm GIT}$ 36.1 Acetate Bead 27.0 6.36 19.5 16.5 18.0 22.5 22.5 31.5 2.12 70.5 $\mathrm{TT}_{_{1}}$ 9.15 7.50 10.66 5.15 6.55 5.85 0.99 2.52 4.72 8.94 7.44 7.94 (kg) BW Jalapeno Pilgrim Phoebe Jessie Shadow Howie Katie Waldo Mean Mean SD SD ID Alouatta villosa palliata Alouatta seniculus sara Alouatta caraya fed diet 15ADF. Table B13. Species

Table B14. Digesta transit time (TT_1) and total retention time (R_{GIT}) in hours of howler monkeys (Alouatta spp.) fed diet 30ADF.

Species								
	ID	BW			Marker type	type		
		(kg)	Bead	ָּסַ	CO-EDTA	DTA	Cr-fiber	ber
			\mathbf{TT}_1	$R_{\rm GIT}$	\mathtt{TT}_1	RGIT	TT_1	RGIT
Alouatta caraya	Phoebe	4.72	19.5	8.0	7.5	27.8	19.5	41.5
Ж	Katie	8.94	10.5	17.0	4.5	29.3	46.5	10.8
ω 	Shadow	10.66	1	1	16.5	58.1	١	1
Σ	Mean	8.10	15.0	12.5	9.5	38.4	33.0	26.1
S	SD	3.06	6.36	6.36	6.24	17.08	19.09	27.71
Alouatta villosa palliata	Jalapeno	6.05	7.5	25.3	4.5	33.3	4.5	35.8
4	Waldo	6.85	19.5	25.5	10.5	27.9	4.5	37.8
	Mean	6.45	13.5	25.4	7.5	30.6	4.5	36.8
0,7	SD	0.57	8.49	0.14	4.24	3.82	00.00	1.41
Alouatta seniculus sara	Pilgrim	8.55	46.5	32.0	46.5	82.0	28.5	69.5
	Howie	7.50	25.5	13.9	22.5	24.6	25.5	12.2
	Mean	8.03	36.0	22.9	34.5	53.3	27.0	40.8
	SD	0.74	14.85	12.80	16.97	40.59	2.12	40.52
•								
Alouatta sp.	Mean	7.61	21.5	20.3	16.1	40.4	21.5	34.6
	SD	1.97	13.90	8.86	14.94	21.53	15.95	21.66

Table Bl5. Digesta transit time (TT ₁) 15ADF.	and total	retention time	time (R	(R _{GIT}) in hours of		olobine	colobine primates	fed diet
Species	ID	BW			Marker type	type		
		(kg)	Bead	ซ	CO-EDTA	ota	Cr-fiber	ber
			TT_1	$R_{\rm GIT}$	\mathtt{TT}_1	$R_{\rm GIT}$	\mathtt{TT}_1	$R_{\rm GIT}$
Colobus guereza kikuyuensis	Flower	11.34	I	I	28.5	41.2	ı	ı
	Gomez	10.90	31.5	20.7	43.5	46.9	46.5	37.6
	Mean	11.12	31.5	20.7	36.0	44.0	46.5	37.6
	SD	0.31	NA	NA	10.6	4.0	NA	ď
Pygathrix nemaeus nemaeus	Rhett	12.10	31.5	I	1.5	35.3	1.5	33.8
Trachypithecus francoisi francoisi	Janet	5.67	31.5	44.9	10.5	37.7	31.5	22.3
	Chrissie	6.67	22.5	48.4	10.5	50.6	22.5	38.8
	Stanley	3.99	١	ı	22.5	39.2	1	ı
	Fungar	5.75	١	ı	10.5	37.0	1	1
	Mean	5.96	27.0	46.7	13.5	41.1	27.0	30.5
	SD	1.38	6.36	2.48	6.00	6.38	6.36	11.65
Colobines	Mean	8.02	29.3	38.0	18.2	41.1	25.5	33.1
	SD	3.04	4.50	15.08	14.27	5.62	18.81	7.52

Species	ID	BW			Marker type	type		
		(kg)	Bead	ט	CO-EDTA	DTA	Cr-fiber	ber
			\mathtt{TT}_1	RGIT	\mathtt{TT}_1	$R_{\rm GIT}$	TT_1	RGIT
Colobus guereza kikuyuensis	Gomez	10.70	31.5	37.6	43.5	53.4	43.5	9.89
Pygathrix nemaeus nemaeus	Rhett	11.75	43.5	I	19.5	32.5	1	ı
Trachypithecus francoisi francoisi	Jack	7.78	22.5	17.4	10.5	36.6	t	1
	Janet	5.72	ı	1	19.5	27.3	1	1
	Chrissie	5.76	ı	1	19.5	40.5	ı	1
	Stanley	3.77	19.5	15.6	4.5	47.8	10.5	42.5
	Fungar	6.75	28.5	95.7	19.5	28.6	I	1
	Goo Ngoot	5.85	43.5	42.7	19.5	32.5	19.5	35.6
	Mean	5.94	28.5	42.8	15.5	35.5	15.0	39.0
	SD	1.33	10.68	37.34	6.48	7.76	6.36	4.85
Colobines	Mean	7.26	31.8	41.8	19.5	37.4	24.5	48.9
	SD	2.71	10.58	32.42	11.22	9.27	17.06	17.41

Table B17. Body weight (BW) and dry matter intake (DMI) of two cercopithecine monkeys fed diet 15ADF and apparent digestibility of dry matter (DM) and fiber fractions in that diet.

	ADL	81.5
	4	8
of.	нс	77.
gestibility (%) of*	υ	23.8 85.7 76.5 75.7 75.4 77.9
stibili	ADF	75.7
Dige	NDF	76.5
	MQ	85.7
IMG	(g/kg BW) DM NDF	23.8
BW	(kg)	9.6
ID		Heine
		ercocebus g. chrysogaster
Species		Cercocebus g.

Mandrillus 1. leucophaeus Susie 10.20 11.7 85.6 76.1 83.2 77.4 77.5 82.1 TM-dry matter; NDF-neutral detergent fiber; ADF-acid detergent fiber; ADF-a detergent lignin.

Table B18. Body weight (BW) and dry matter intake (DMI) of two cercopithecine monkeys fed diet 30ADF and apparent digestibility of dry matter (DM) and fiber fractions in that diet.	W) and dry dry matter	matter intake (DM) and fiber	(DMI) of two fractions in	cercopit that die	hecine t.	monkeys	fed o	liet 30	ADF 8	pui
Species	ID	BW	DMI		Dige	Digestibility (%) of	ty (%)	of,		
		(kg)	(g/kg BW) DM NDF ADF C HC ADL	DM	NDF	ADF	υ	НС	A	IC
Cercocebus g. chrysogaster Heine	r Heine	10.10	23.5	23.5 82.0 80.5 78.0 78.7 86.0 70.7	80.5	78.0	78.7	86.	7 0	7.0

Mandrillus 1 leucophaeus Susie 10.00 7.2 81.1 79.2 79.0 77.6 84.9 46.2 The Theorem of the Theore

Table B19. Body weight (BW) of two cercopithecine monkeys fed diets 15ADF and 30ADF and the apparent

digestibility of dry matter and groa	and gross en	ss energy in those diets.	hose die	ts.				
Species	ΙD	BW		15ADF			30ADF	
		(kg)	DM (%) GE (%)	GE (%)	DE (kcal/g)	DM (%) GE (%)	(%) ES	DE (kcal/g)
Cercocebus g. chrysogaster	Heine	9.60	9.60 85.7	89.1	3.97	82.0	82.5	3.62
Mandrillus 1. leucophaeus Susie 10.20 85 *DM=dry matter; GE=gross energy; DE=digestible energy	Susie rgy; DE=diges	10.20 stible en	10.20 85.6 83.7 ble energy	83.7	3.73	81.1	78.9	3.46

